

DIMENSION OF SELF-AFFINE SETS FOR FIXED TRANSLATION VECTORS

BALÁZS BÁRÁNY, ANTTI KÄENMÄKI, AND HENNA KOIVUSALO

ABSTRACT. An affine iterated function system is a finite collection of affine invertible contractions and the invariant set associated to the mappings is called self-affine. In 1988, Falconer proved that, for given matrices, the Hausdorff dimension of the self-affine set is the affinity dimension for Lebesgue almost every translation vectors. Similar statement was proven by Jordan, Pollicott, and Simon in 2007 for the dimension of self-affine measures. In this article, we have an orthogonal approach. We introduce a class of self-affine systems in which, given translation vectors, we get the same results for Lebesgue almost all matrices.

1. INTRODUCTION

For a non-singular $d \times d$ matrix $A \in GL_d(\mathbb{R})$ and a translation vector $v \in \mathbb{R}^d$, let us denote the affine map $x \mapsto Ax + v$ by $f = f(A, v)$. Let $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$ be a tuple of contractive non-singular $d \times d$ matrices and let $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^d)^N$ be a tuple of translation vectors. Here and throughout we assume that $N \geq 2$ is an integer. The tuple $\Phi_{\mathbf{A}, \mathbf{v}} = (f_1, \dots, f_N)$ obtained from the affine mappings $f_i = f(A_i, v_i)$ is called the *affine iterated function system (IFS)*. Hutchinson [15] showed that for each $\Phi_{\mathbf{A}, \mathbf{v}}$ there exists a unique non-empty compact set $E = E_{\mathbf{A}, \mathbf{v}}$ such that

$$E = \bigcup_{i=1}^N f_i(E).$$

The set $E_{\mathbf{A}, \mathbf{v}}$ associated to an affine IFS $\Phi_{\mathbf{A}, \mathbf{v}}$ is called *self-affine*. In the special case where each of the linear maps A_i is a scalar multiple of an isometry, we call $\Phi_{\mathbf{A}, \mathbf{v}}$ a *similitude iterated function system* and the set $E_{\mathbf{A}, \mathbf{v}}$ *self-similar*.

The dimension theory of self-similar sets satisfying a sufficient separation condition was completely resolved by Hutchinson [15]. Without separation, i.e. when the images $f_i(E)$ and $f_j(E)$ can have severe overlapping, the problem is more difficult. The most recent progress in this direction is by Hochman [13, 14]. Among other things, he managed to calculate the Hausdorff dimension of a self-similar set on the real line under very mild assumptions. The dimension theory of self-affine sets and measures is far from being fully understood. Hence, while working on the topic, it is common to focus on specific subclasses of self-affine sets, for which more methods are available, even though recently the general case is becoming more accessible as well; see [2, 3, 8]. One very popular subclass being studied is that of self-affine carpets. In this class the structure of self-affine set has been made more tractable by imposing special relations between the affine maps. For recent results in this direction, see [11, 12, 21]. Another standard subclass of self-affine sets being considered is introduced by Falconer [6]. He proved that for a fixed matrix tuple $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$, in which the operator norms $\|A_i\|$ are strictly less than $1/2$, the Hausdorff dimension of the self-affine set $E_{\mathbf{A}, \mathbf{v}}$, $\dim_H(E_{\mathbf{A}, \mathbf{v}})$, is the affinity dimension of \mathbf{A} , $\dim_{\text{aff}}(\mathbf{A})$, for \mathcal{L}^{dN} -almost all $\mathbf{v} \in (\mathbb{R}^d)^N$. Here \mathcal{L}^d is the d -dimensional Lebesgue measure and the affinity dimension is a number depending only

Date: November 27, 2016.

2010 Mathematics Subject Classification. Primary 37C45; Secondary 28A80.

Key words and phrases. Self-affine set, self-affine measure, Hausdorff dimension.

BB acknowledges the support from the grant OTKA K104745 and the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, and HK from EPSRC EP/L001462 and Osk. Huttunen foundation.




FIGURE 1. The picture illustrates three example cases where some other covering is more optimal than the one obtained from (1.1). From left to right: ellipses have severe overlapping, the ellipsis does not contain E all the way, and ellipses are badly aligned.

on \mathbf{A} defined below. A similar result also holds for self-affine measures, due to Jordan, Pollicott and Simon [16].

Let us next give an intuitive explanation for Falconer's result. It is easy to see that

$$E = \bigcap_{n=1}^{\infty} \bigcup_{i_1, \dots, i_n \in \{1, \dots, N\}} f_{i_1} \circ \dots \circ f_{i_n}(B(0, R)), \quad (1.1)$$

where $B(0, R)$ is the closed ball centered at the origin with radius $R = \max_{i \in \{1, \dots, N\}} |v_i| / (1 - \max_{i \in \{1, \dots, N\}} \|A_i\|) > 0$. Since we are interested in the dimension of E we may, by rescaling if necessary, assume that $R = 1$. We immediately see from (1.1) that for each fixed n the sets $f_{i_1} \circ \dots \circ f_{i_n}(B(0, 1))$ form a cover for the self-affine set. For any $A \in GL_d(\mathbb{R})$, let $1 > \alpha_1(A) \geq \dots \geq \alpha_d(A) > 0$ be the lengths of the principal semiaxes of the ellipse $A(B(0, 1))$. Observe that $f_{i_1} \circ \dots \circ f_{i_n}(B(0, 1))$ is a translated copy of $A_{i_1} \cdots A_{i_n}(B(0, 1))$. To find the Hausdorff dimension of E , it is necessary to find optimal covers for E . Natural candidates for such covers come immediately from (1.1). In \mathbb{R}^2 , we need approximately $\alpha_1(A_{i_1} \cdots A_{i_n}) / \alpha_2(A_{i_1} \cdots A_{i_n})$ many balls of radius $\alpha_2(A_{i_1} \cdots A_{i_n})$ to cover $f_{i_1} \circ \dots \circ f_{i_n}(B(0, R))$. By the definition of the s -dimensional Hausdorff measure \mathcal{H}^s , it follows that

$$\begin{aligned} \mathcal{H}^s(E) &\lesssim \lim_{n \rightarrow \infty} \sum_{i_1, \dots, i_n \in \{1, \dots, N\}} \frac{\alpha_1(A_{i_1} \cdots A_{i_n})}{\alpha_2(A_{i_1} \cdots A_{i_n})} \alpha_2(A_{i_1} \cdots A_{i_n})^s \\ &= \lim_{n \rightarrow \infty} \sum_{i_1, \dots, i_n \in \{1, \dots, N\}} \alpha_1(A_{i_1} \cdots A_{i_n}) \alpha_2(A_{i_1} \cdots A_{i_n})^{s-1}. \end{aligned}$$

The *singular value pressure* $P_{\mathbf{A}}$ of \mathbf{A} in this case is

$$P_{\mathbf{A}}(s) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{i_1, \dots, i_n \in \{1, \dots, N\}} \alpha_1(A_{i_1} \cdots A_{i_n}) \alpha_2(A_{i_1} \cdots A_{i_n})^{s-1}.$$

For the complete definition, see (2.3). The function $s \mapsto P_{\mathbf{A}}(s)$ is strictly decreasing and it has a unique zero. If $P_{\mathbf{A}}(s) < 0$, then the sum above is strictly less than one for all large enough n . Therefore, defining $\dim_{\text{aff}}(\mathbf{A})$ to be the minimum of 2 and s for which $P_{\mathbf{A}}(s) = 0$, we have $\dim_{\text{H}}(E_{\mathbf{A}, \mathbf{v}}) \leq \dim_{\text{aff}}(\mathbf{A})$ for all $\mathbf{v} \in (\mathbb{R}^2)^N$. The question then becomes, when are the covers obtained in this way optimal. It is easy to find situations in which some other cover is more efficient; see Figure 1. Intuitively, since the role of the translation vector is to determine the placement of the ellipses, Falconer's result asserts that one never encounters these situations with a random choice of translation vectors.

Our approach is opposite to that of Falconer's. We fix the tuple of translation vectors and study what can be said about the dimension for different choices of matrix tuples. We remark that the first result in this direction is given by Bárány and Rams [4]. Their proof relies on the dimension of the Furstenberg measure and on the exceptional sets of the dimension of orthogonal projections. In the same vein as with the intuitive explanation of Falconer's result, one expects that, keeping the centers of the ellipses fixed, a small random change in the shape of the ellipses guarantees that the covers obtained from (1.1) are optimal. Indeed, in the main results of the paper, Theorems A and B, we fix the tuple of distinct translation vectors $\mathbf{v} \in (\mathbb{R}^d)^N$, and show that $\dim_{\text{H}}(E_{\mathbf{A}, \mathbf{v}}) = \dim_{\text{aff}}(\mathbf{A})$

for \mathcal{L}^{d^2N} -almost all \mathbf{A} in a large open set of matrix tuples. An interesting feature of our results is that the planar case is different from the higher dimensional case. The planar case is stated in Theorem A and the higher dimensional case in Theorem B. This reflects the fact that the proofs in these cases have major differencies; see Remark 2.1.

The remainder of the paper is organized as follows. In §2, we give a detailed explanation of the setting and state our main results. We explore in §3 how the Hausdorff dimension of an ergodic measure satisfying the Ledrappier-Young formula can be calculated under a modified self-affine transversality condition. In §4, we prove an analogous result for self-affine sets. Finally, the proof of Theorem A is given in §5 and Theorem B is proved in §6.

2. PRELIMINARIES AND STATEMENTS OF MAIN RESULTS

Let Σ be the set of one-sided words of symbols $\{1, \dots, N\}$ with infinite length, i.e. $\Sigma = \{1, \dots, N\}^{\mathbb{N}}$. Let us denote the left-shift operator on Σ by σ . Let the set of words with finite length be $\Sigma_* = \bigcup_{n=0}^{\infty} \{1, \dots, N\}^n$ with the convention that the only word of length 0 is the empty word \emptyset . The set $\Sigma_n = \{1, \dots, N\}^n$ is the collection of words of length n . Denote the length of $\mathbf{i} \in \Sigma \cup \Sigma_*$ by $|\mathbf{i}|$, and for finite or infinite words \mathbf{i} and \mathbf{j} , let $\mathbf{i} \wedge \mathbf{j}$ be their common beginning. The concatenation of two words \mathbf{i} and \mathbf{j} is denoted by \mathbf{ij} . We define the cylinder sets of Σ in the usual way, that is, by setting

$$[\mathbf{i}] = \{\mathbf{j} \in \Sigma : \mathbf{i} \wedge \mathbf{j} = \mathbf{i}\} = \{\mathbf{ij} \in \Sigma : \mathbf{j} \in \Sigma\}$$

for all $\mathbf{i} \in \Sigma_*$. For a word $\mathbf{i} = (i_1, \dots, i_n)$ with finite length let $f_{\mathbf{i}}$ be the composition $f_{i_1} \circ \dots \circ f_{i_n}$ and $A_{\mathbf{i}}$ be the product $A_{i_1} \cdots A_{i_n}$. For $\mathbf{i} \in \Sigma \cup \Sigma_*$ and $n < |\mathbf{i}|$, let $\mathbf{i}|_n$ be the first n symbols of \mathbf{i} . Let $\mathbf{i}|_0 = \emptyset$, A_{\emptyset} be the identity matrix, and f_{\emptyset} be the identity function. Finally, we define the *natural projection* $\pi = \pi_{\mathbf{A}, \mathbf{v}} : \Sigma \rightarrow E_{\mathbf{A}, \mathbf{v}}$ by setting

$$\pi(\mathbf{i}) = \sum_{k=1}^{\infty} A_{\mathbf{i}|_{k-1}} v_{i_k} \quad (2.1)$$

for all $\mathbf{i} \in \Sigma$. Note that $E = \bigcup_{\mathbf{i} \in \Sigma} \pi(\mathbf{i})$.

Denote by $\alpha_i(A)$ the i -th largest (counting with multiplicity) *singular value* of a matrix $A \in GL_d(\mathbb{R})$, i.e. the positive square root of the i -th eigenvalue of AA^T , where A^T is the transpose of A . We note that $\alpha_1(A)$ is the usual operator norm $\|A\|$ induced by the Euclidean norm on \mathbb{R}^d and $\alpha_d(A)$ is the mininorm $\mathbf{m}(A) = \|A^{-1}\|^{-1}$. We say that A is *contractive* if $\|A\| < 1$. For a given tuple $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$ we also set $\|\mathbf{A}\| = \max_{i \in \{1, \dots, N\}} \|A_i\|$ and $\mathbf{m}(\mathbf{A}) = \min_{i \in \{1, \dots, N\}} \mathbf{m}(A_i)$. Following Falconer [6], we define the *singular value function* φ^s of a matrix A by setting

$$\varphi^s(A) = \begin{cases} \alpha_1(A) \cdots \alpha_{[s]}(A) \alpha_{[s]}(A)^{s-[s]}, & \text{if } 0 \leq s \leq d, \\ |\det A|^{s/d}, & \text{if } s > d. \end{cases}$$

The singular value function satisfies

$$\varphi^s(AB) \leq \varphi^s(A) \varphi^s(B)$$

for all $A, B \in GL_d(\mathbb{R})$. Moreover, if $(A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$, then

$$\varphi^s(A_i) \mathbf{m}(\mathbf{A})^{\delta|\mathbf{i}|} \leq \varphi^{s+\delta}(A_i) \leq \varphi^s(A_i) \|\mathbf{A}\|^{\delta|\mathbf{i}|} \quad (2.2)$$

for all $\mathbf{i} \in \Sigma_*$ and $s, \delta \geq 0$.

For a tuple $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$ of contractive non-singular $d \times d$ matrices the function $P_{\mathbf{A}} : [0, \infty) \rightarrow \mathbb{R}$ defined by

$$P_{\mathbf{A}}(s) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \sum_{\mathbf{i} \in \Sigma_n} \varphi^s(A_{\mathbf{i}}) \quad (2.3)$$

is called the *singular value pressure*. It is well-defined, continuous, strictly decreasing on $[0, \infty)$, and convex between any two integers. Moreover, $P_{\mathbf{A}}(0) = \log N$ and $\lim_{s \rightarrow \infty} P_{\mathbf{A}}(s) = -\infty$. Let us

denote by $\dim_{\text{aff}} \mathbf{A}$ the minimum of d and the unique root of the singular value pressure function and call it the *affinity dimension*.

If μ is a Radon measure on \mathbb{R}^d , then the *upper* and *lower local dimensions* of μ at x are defined by

$$\overline{\dim}_{\text{loc}}(\mu, x) = \limsup_{r \downarrow 0} \frac{\log \mu(B(x, r))}{\log r} \quad \text{and} \quad \underline{\dim}_{\text{loc}}(\mu, x) = \liminf_{r \downarrow 0} \frac{\log \mu(B(x, r))}{\log r},$$

respectively. The measure μ is *exact-dimensional* if

$$\text{ess inf}_{x \sim \mu} \underline{\dim}_{\text{loc}}(\mu, x) = \text{ess sup}_{x \sim \mu} \overline{\dim}_{\text{loc}}(\mu, x).$$

In this case, the common value is denoted by $\dim \mu$. The above quantities are naturally linked to set dimensions. For example, the *lower Hausdorff dimension* of the measure μ is

$$\underline{\dim}_{\text{H}} \mu = \text{ess inf}_{x \sim \mu} \underline{\dim}_{\text{loc}}(\mu, x) = \inf \{ \dim_{\text{H}} A : A \text{ is a Borel set with } \mu(A) > 0 \}.$$

Here $\dim_{\text{H}} A$ is the Hausdorff dimension of the set A . For more detailed information, the reader is referred to [7].

Fix a probability vector $\mathbf{p} = (p_1, \dots, p_N) \in (0, 1)^N$ and denote the product $p_{i_1} \cdots p_{i_n}$ by $p_{\mathbf{i}}$ for all finite words $\mathbf{i} = (i_1, \dots, i_n)$. Let $\nu_{\mathbf{p}}$ be the corresponding *Bernoulli measure* on Σ . It is uniquely defined by setting $\nu_{\mathbf{p}}([\mathbf{i}]) = p_{\mathbf{i}}$ for all $\mathbf{i} \in \Sigma_*$. It is easy to see that $\nu_{\mathbf{p}}$ is σ -invariant and ergodic. We say that ν on Σ is a *step- n Bernoulli measure* if it is a Bernoulli measure on $(\Sigma_n)^{\mathbb{N}}$ for some probability vector from $(0, 1)^{N^n}$. Furthermore, we say that a measure ν on Σ is *quasi-Bernoulli* if there is a constant $C \geq 1$ such that

$$C^{-1} \nu([\mathbf{i}]) \nu([\mathbf{j}]) \leq \nu([\mathbf{ij}]) \leq C \nu([\mathbf{i}]) \nu([\mathbf{j}])$$

for all $\mathbf{i}, \mathbf{j} \in \Sigma_*$. The *entropy* of a σ -invariant measure ν on Σ is

$$h_{\nu} = - \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{\mathbf{i} \in \Sigma_n} \nu([\mathbf{i}]) \log \nu([\mathbf{i}]). \quad (2.4)$$

Note that the entropy of a Bernoulli measure $\nu_{\mathbf{p}}$ is given by $h_{\mathbf{p}} = - \sum_{i=1}^N p_i \log p_i$.

If $\nu_{\mathbf{p}}$ is a Bernoulli measure and $\Phi_{\mathbf{A}, \mathbf{v}}$ is an affine iterated function system, then the push-down measure $\mu_{\mathbf{A}, \mathbf{v}, \mathbf{p}} = \pi_{\mathbf{A}, \mathbf{v}} \nu_{\mathbf{p}} = \nu_{\mathbf{p}} \circ (\pi_{\mathbf{A}, \mathbf{v}})^{-1}$ is called *self-affine*. It is well known that the self-affine measure $\mu = \mu_{\mathbf{A}, \mathbf{v}, \mathbf{p}}$ satisfies

$$\mu = \sum_{i=1}^N p_i f_i \mu.$$

We say that $\mathbf{A} \in GL_d(\mathbb{R})^N$ satisfies the *totally dominated splitting condition* if there exist constants $C \geq 1$ and $0 < \tau < 1$ such that for every $i \in \{1, \dots, d-1\}$ either

$$\frac{\alpha_{i+1}(A_{\mathbf{i}})}{\alpha_i(A_{\mathbf{i}})} \leq C \tau^{|\mathbf{i}|} \quad (2.5)$$

for every $\mathbf{i} \in \Sigma_*$ or

$$\frac{\alpha_{i+1}(A_{\mathbf{i}})}{\alpha_i(A_{\mathbf{i}})} > C^{-1}$$

for every $\mathbf{i} \in \Sigma_*$. By Bochi and Gourmelon [5, Theorem B], the set

$$\mathcal{D} = \{ \mathbf{A} \in GL_d(\mathbb{R})^N : (2.5) \text{ holds for every } i \in \{1, \dots, d\} \}. \quad (2.6)$$

is an open subset of $GL_d(\mathbb{R})^N$.

If ν is an ergodic measure on Σ , then, by Oseledets' theorem, there exist constants $0 < \chi_1(\mathbf{A}, \nu) \leq \dots \leq \chi_d(\mathbf{A}, \nu) < \infty$ such that

$$\chi_i(\mathbf{A}, \nu) = - \lim_{n \rightarrow \infty} \frac{1}{n} \log \alpha_i(A_{i_1} \cdots A_{i_n}) \quad (2.7)$$

for ν -almost every $\mathbf{i} \in \Sigma$. The numbers $\chi_i(\mathbf{A}, \nu)$ are called the *Lyapunov exponents* of \mathbf{A} with respect to ν . The Lyapunov exponents of \mathbf{A} with respect to a Bernoulli measure $\nu_{\mathbf{p}}$ are denoted by $\chi_i(\mathbf{A}, \mathbf{p})$. Furthermore, let us define the *Lyapunov dimension* of ν by

$$\dim_{\mathbb{L}} \nu = \min_{k \in \{0, \dots, d\}} \left\{ k + \frac{h_{\nu} - \sum_{j=1}^k \chi_j(\mathbf{A}, \nu)}{\chi_{k+1}(\mathbf{A}, \nu)}, d \right\}.$$

The Lyapunov dimension of the projected measure $\pi\nu$ on $E_{\mathbf{A}, \mathbf{v}}$ is defined by setting $\dim_{\mathbb{L}} \pi\nu = \dim_{\mathbb{L}} \nu$. Käenmäki [17, Theorems 2.6 and 4.1] proved the existence of ergodic equilibrium states. If $s = \dim_{\text{aff}} \mathbf{A} \leq d$, then an ergodic s -equilibrium state μ of \mathbf{A} on Σ is defined by the equality

$$h_{\mu} = -\lim_{\frac{1}{n}} \sum_{\mathbf{i} \in \Sigma_n} \mu([\mathbf{i}]) \log \varphi^s(A_{\mathbf{i}}) = \sum_{j=1}^{\lfloor s \rfloor} \chi_j(\mathbf{A}, \mu) + (s - \lfloor s \rfloor) \chi_{\lfloor s \rfloor}(\mathbf{A}, \mu). \quad (2.8)$$

Here the second equality follows from Kingman's ergodic theorem. It is easy to see that such an s -equilibrium state is a measure of maximal Lyapunov dimension.

Bárány and Käenmäki recently proved in [3, Theorem 2.3] that the self-affine measure $\mu_{\mathbf{A}, \mathbf{v}, \mathbf{p}}$ is exact-dimensional regardless of the choices of \mathbf{A} , \mathbf{v} , and \mathbf{p} provided that all the corresponding Lyapunov exponents are distinct. Furthermore, they showed that if \mathbf{A} satisfies the totally dominated splitting condition, then the image of any quasi-Bernoulli measure under $\pi_{\mathbf{A}, \mathbf{v}}$ is exact-dimensional regardless of the choice of \mathbf{v} ; see [3, Theorem 2.6].

Let us next state the main results of the article. Let \mathcal{L}^d be the d -dimensional Lebesgue measure and $\overline{\dim}_{\mathbb{M}}$ be the upper Minkowski dimension. We define $\|\mathbf{v}\| = \max_{i \in \{1, \dots, N\}} |v_i|$ and recall that $\|\mathbf{A}\| = \max_{i \in \{1, \dots, N\}} \|A_i\|$.

Theorem A. *Suppose that $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^2)^N$ is such that $v_i \neq v_j$ for $i \neq j$ and*

$$\mathcal{A}_{\mathbf{v}} = \left\{ \mathbf{A} \in GL_2(\mathbb{R})^N : 0 < \max_{i \neq j} \frac{\|A_i\| + \|A_j\|}{|v_i - v_j|} \cdot \frac{\|\mathbf{v}\|}{1 - \|\mathbf{A}\|} < \frac{\sqrt{2}}{2} \right\}. \quad (2.9)$$

Then

$$\dim_{\mathbb{H}} E_{\mathbf{A}, \mathbf{v}} = \overline{\dim}_{\mathbb{M}} E_{\mathbf{A}, \mathbf{v}} = \dim_{\text{aff}} \mathbf{A}$$

for \mathcal{L}^{4N} -almost all $\mathbf{A} \in \mathcal{A}_{\mathbf{v}}$. Moreover, for every probability vector $\mathbf{p} \in (0, 1)^N$ the corresponding self-affine measure $\mu = \mu_{\mathbf{A}, \mathbf{v}, \mathbf{p}}$ satisfies

$$\dim \mu = \dim_{\mathbb{L}} \mu = \min \left\{ \frac{h_{\mathbf{p}}}{\chi_1(\mathbf{A}, \mathbf{p})}, 1 + \frac{h_{\mathbf{p}} - \chi_1(\mathbf{A}, \mathbf{p})}{\chi_2(\mathbf{A}, \mathbf{p})} \right\}$$

for \mathcal{L}^{4N} -almost all $\mathbf{A} \in \mathcal{A}_{\mathbf{v}}$.

Note that matrix tuples in $\mathcal{A}_{\mathbf{v}}$ are contractive. In higher dimensions, our result is the following.

Theorem B. *Suppose that $d \in \mathbb{N}$ is such that $d \geq 3$, $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^d)^N$ is such that $v_i \neq v_j$ for $i \neq j$, and*

$$\mathcal{A}'_{\mathbf{v}} = \left\{ \mathbf{A} \in GL_d(\mathbb{R})^N : 0 < \max_{i \neq j} \frac{\|A_i\| + \|A_j\|}{|v_i - v_j|} \cdot \frac{\|\mathbf{v}\|}{1 - \|\mathbf{A}\|} < \frac{2}{\sqrt{3}} - 1 \right\}. \quad (2.10)$$

Then for every probability vector $\mathbf{p} \in (0, 1)^N$ the corresponding self-affine measure $\mu = \mu_{\mathbf{A}, \mathbf{v}, \mathbf{p}}$ satisfies

$$\dim \mu = \dim_{\mathbb{L}} \mu = \min_{k \in \{0, \dots, d\}} \left\{ k + \frac{h_{\mathbf{p}} - \sum_{j=1}^k \chi_j(\mathbf{A}, \mathbf{p})}{\chi_{k+1}(\mathbf{A}, \mathbf{p})}, d \right\}$$

for \mathcal{L}^{d^2N} -almost all $\mathbf{A} \in \mathcal{A}'_{\mathbf{v}}$. Moreover,

$$\dim_{\mathbb{H}} \mu_{\mathbf{A}} = \dim_{\mathbb{H}} E_{\mathbf{A}, \mathbf{v}} = \overline{\dim}_{\mathbb{M}} E_{\mathbf{A}, \mathbf{v}} = \dim_{\text{aff}} \mathbf{A}$$

for \mathcal{L}^{d^2N} -almost all $\mathbf{A} \in \mathcal{A}'_{\mathbf{v}} \cap \mathcal{D}$, where \mathcal{D} is as in (2.6) and $\mu_{\mathbf{A}}$ is an ergodic s -equilibrium state of \mathbf{A} for $s = \dim_{\text{aff}} \mathbf{A}$.




FIGURE 2. The condition (2.9) requires that the images of the unit ball under f_i are separated away from each other as in the picture: if the ellipses are rotated around their centers, they still stay inside a cone having an angle at most $\pi/2$.

Remark 2.1. We emphasize that the methods used to prove Theorems A and B are significantly different. At first, the higher dimensional exact dimensionality result (the Ledrappier-Young formula to be more precise) of Bárány and Käenmäki [3] requires the totally dominated splitting condition, which is why we restrict our matrix tuples to the set \mathcal{D} . Very recently, Feng [personal communication] has informed the authors that the Ledrappier-Young formula holds also without totally dominated splitting. By relying on this, one could improve Theorem B by replacing $\mathcal{A}'_{\mathbf{v}} \cap \mathcal{D}$ by $\mathcal{A}'_{\mathbf{v}}$. Secondly, the transversality argument used in the higher dimensional case is different from the two-dimensional case. Curiously, the higher dimensional transversality argument requires the dimension to be at least three, so it cannot be used in the two-dimensional case. This difference is also the reason why we use different upper bounds in the definitions of $\mathcal{A}_{\mathbf{v}}$ and $\mathcal{A}'_{\mathbf{v}}$. Currently we do not know if the upper bound $2/\sqrt{3} - 1$ used in (2.10) can be replaced by the upper bound $\sqrt{2}/2$ used in (2.9). The sharpness of the methods used in our proofs is discussed in Remark 3.6.

Remark 2.2. Many of the recent works on dimensions of self-affine sets and measures (see e.g. [3, 22, 24]) rely on properties of the Furstenberg measure (for definitions, see §3). However, Theorem A covers situations that cannot be addressed by using this approach. Even though the condition (2.9) is rather restrictive (for example, we will see in Lemma 2.4 that it implies that the images $f_i(E)$ are disjoint, i.e. $\Phi_{\mathbf{A}, \mathbf{v}}$ satisfies the *strong separation condition*) Theorem A introduces a checkable condition for an affine iterated function system to satisfy the desired dimension result. This is in contrast to, for example, [3, Corollaries 2.7 and 2.8] where the claim for the self-affine measure μ in Theorem A holds provided that the strong separation condition holds and the dimension of μ does not drop when projected to orthogonal complements of Furstenberg typical lines. The condition (2.9) can be illustrated via Lemma 3.7 as in Figure 2.

We will next exhibit an open set of matrices that satisfy the assumptions of Theorem A, but do not satisfy the projection condition of [3]. Consider an affine iterated function system consisting of three mappings $x \mapsto A_i x + v_i$. Let the translation vectors v_i be equidistributed on the unit circle, that is, $|v_i| = 1$ and $|v_i - v_j| = \sqrt{3}$ for all $i \neq j$. It is easy to see that if $\|A_i\| < \sqrt{6}/(4 + \sqrt{6}) =: \eta$ for all i , then $\mathbf{A} \in \mathcal{A}_{\mathbf{v}}$. Since $\eta > 1/3$ we find an open set of three matrices such that all elements are strictly positive, the Furstenberg measure is supported on a Cantor set having dimension less than $1/2$, and the affinity dimension is between 1 and $3/2$. Recalling [3, Corollary 2.9], we see that this case does not satisfy the projection condition.

We record a small lemma showing that the equilibrium states in Theorem B are quasi-Bernoulli.

Lemma 2.3. *For $\mathcal{L}^{d^2 N}$ -almost every $\mathbf{A} \in \mathcal{A}'_{\mathbf{v}} \cap \mathcal{D}$ the unique s -equilibrium state of \mathbf{A} is quasi-Bernoulli for $s = \dim_{\text{aff}}(\mathbf{A})$.*

Proof. By [19, Propositions 3.4 and 3.6], for $\mathcal{L}^{d^2 N}$ -almost every $\mathbf{A} \in GL_d(\mathbb{R})^N$, the s -equilibrium state of \mathbf{A} for $s = \dim_{\text{aff}} \mathbf{A}$ is unique and satisfies the following Gibbs property: there exists a constant $C \geq 1$ such that

$$C^{-1} \varphi^s(A_{\mathbf{i}}) \leq \mu([\mathbf{i}]) \leq C \varphi^s(A_{\mathbf{i}})$$

for all $\mathbf{i} \in \Sigma_*$. By [5, Theorem B] and [10, Lemma 2.1], for each $\mathbf{A} \in \mathcal{D}$, there exists a constant $C' > 0$ such that

$$C' \varphi^s(A_{\mathbf{i}}) \varphi^s(A_{\mathbf{j}}) \leq \varphi^s(A_{\mathbf{ij}})$$

for all $\mathbf{i}, \mathbf{j} \in \Sigma_*$. The statement of the lemma follows. \square

In the next lemma, we note that the strong separation condition follows from the conditions (2.9) and (2.10). Denote by $O(d)$ the orthogonal group of matrices $G \in GL_d(\mathbb{R})$ with $G^T G = I$. For $G \in O(d)$, we denote the vector (Gv_1, \dots, Gv_d) by $G(\mathbf{v})$.

Lemma 2.4. *Let $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^d)^N$ be such that $v_i \neq v_j$ for $i \neq j$ and $\mathbf{A} = (A_1, \dots, A_N) \in \mathcal{A}_{\mathbf{v}} \cup \mathcal{A}'_{\mathbf{v}}$. The affine iterated function system $\Phi_{\mathbf{A}, G(\mathbf{v})}$ satisfies the strong separation condition for all $G \in O(d)$.*

Proof. By definition, it is easy to see that

$$|\pi_{\mathbf{A}, G(\mathbf{v})}(\mathbf{i})| \leq \frac{\|\mathbf{v}\|}{1 - \|\mathbf{A}\|}$$

for every $\mathbf{i} \in \Sigma$ and $G \in O(d)$. If $\mathbf{i}, \mathbf{j} \in \Sigma$ are such that $\mathbf{i}|_1 \neq \mathbf{j}|_1$, then, by either (2.9) or (2.10),

$$\begin{aligned} |\pi_{\mathbf{A}, G(\mathbf{v})}(\mathbf{i}) - \pi_{\mathbf{A}, G(\mathbf{v})}(\mathbf{j})| &\geq |Gv_{i_1} - Gv_{j_1}| - |A_{i_1} \pi_{\mathbf{A}, G(\mathbf{v})}(\sigma(\mathbf{i})) - A_{j_1} \pi_{\mathbf{A}, G(\mathbf{v})}(\sigma(\mathbf{j}))| \\ &\geq |v_{i_1} - v_{j_1}| - (\|A_{i_1}\| + \|A_{j_1}\|) \frac{\|\mathbf{v}\|}{1 - \|\mathbf{A}\|} > 0. \end{aligned}$$

This shows that the strong separation condition holds in both cases, proving the claim. \square

3. MODIFIED SELF-AFFINE TRANSVERSALITY

This section is devoted to proving that for measures, which satisfy the Ledrappier-Young formula, the dimension of the measure is equal to the Lyapunov dimension for almost every matrix tuple whenever the modified self-affine transversality condition, defined below, holds.

Let us denote the Grassmannian of k -planes in \mathbb{R}^d by $G(d, k)$. If $V \in G(d, k)$, then $V^\perp \in G(d, d-k)$ is the subspace orthogonal to V . For $A \in GL_d(\mathbb{R})$ let $\|A|V\|$ be the operator norm of A restricted to V defined by $\|A|V\| = \sup_{v \in V} |Av|/|v|$ and $\mathbf{m}(A|V)$ the mininorm of A restricted to V defined by $\mathbf{m}(A|V) = \inf_{v \in V} |Av|/|v|$.

Let ν be an ergodic σ -invariant measure on Σ and let $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$. We say that $\mu_F^{d,k}$ is the (d, k) -Furstenberg-measure with respect to ν and \mathbf{A} if $\mu_F^{d,k} \times \nu$ is an ergodic T -invariant measure, where

$$T: G(d, k) \times \Sigma \rightarrow G(d, k) \times \Sigma, \quad (V, \mathbf{i}) \mapsto (A_{i_0}^{-1}V, \sigma(\mathbf{i})),$$

and furthermore,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbf{m}(A_{i_n}^{-1} \cdots A_{i_0}^{-1} | V) = \chi_{d-k+1}(\mathbf{A}, \nu) \quad (3.1)$$

for $\mu_F^{d,k} \times \nu$ -almost all $(V, \mathbf{i}) \in G(d, k) \times \Sigma$.

Lemma 3.1. *Let ν be an ergodic measure on Σ and let $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$ be a tuple of contractive matrices. Assume that the $(d, d-k)$ -Furstenberg measure $\mu_F^{d, d-k}$ with respect to ν and \mathbf{A} exists for some $k \in \{1, \dots, d-1\}$. Then*

$$-\lim_{n \rightarrow \infty} \frac{1}{n} \log \varphi^s(\text{proj}_{V^\perp} A_{\mathbf{i}|_n}) = \sum_{j=1}^{\lfloor s \rfloor} \chi_j(\mathbf{A}, \nu) + (s - \lfloor s \rfloor) \chi_{\lceil s \rceil}(\mathbf{A}, \nu)$$

for all $0 \leq s \leq k$ and for $\mu_F^{d-k} \times \nu$ -almost all $(V, \mathbf{i}) \in G(d, d-k) \times \Sigma$.

Proof. Notice that if $A \in GL_d(\mathbb{R})$ and $\ell < s \leq \ell + 1$, then $\varphi^s(A) = \|A^{\wedge \ell}\|^{\ell+1-s} \|A^{\wedge(\ell+1)}\|^{s-\ell}$ and $\varphi^s(A) = \varphi^s(A^T)$; for example, see [20, §3.4]. Thus, it is enough to show that

$$- \lim_{n \rightarrow \infty} \frac{1}{n} \log \|((A_{\mathbf{i}|_n})^T \text{proj}_{V^\perp})^{\wedge \ell}\| = \sum_{j=1}^{\ell} \chi_j(\mathbf{A}, \nu)$$

and the corresponding limit for $\ell + 1$ hold for $\mu_F^{d-k} \times \nu$ -almost all $(V, \mathbf{i}) \in G(d, d-k) \times \Sigma$. It is easy to see that $(\text{proj}_{V^\perp})^{\wedge \ell} = \text{proj}_{\wedge^\ell V^\perp}$ and $\|((A_{\mathbf{i}|_n})^T)^{\wedge \ell} \text{proj}_{\wedge^\ell V^\perp}\| = \|((A_{\mathbf{i}|_n})^T)^{\wedge \ell} | \wedge^\ell V^\perp\|$. Since $k \geq \ell + 1$, it suffices to show that

$$- \lim_{n \rightarrow \infty} \frac{1}{n} \log \|((A_{\mathbf{i}|_n})^T)^{\wedge k} | \wedge^k V^\perp\| = \sum_{j=1}^k \chi_j(\mathbf{A}, \nu) \quad (3.2)$$

for $\mu_F^{d-k} \times \nu$ -almost all $(V, \mathbf{i}) \in G(d, d-k) \times \Sigma$. By the Oseledets' decomposition, this implies that the corresponding limit holds for all values in $\{1, \dots, k\}$.

We denote the Hodge star operator between $\wedge^k \mathbb{R}^d$ and $\wedge^{d-k} \mathbb{R}^d$ by $*$. Let $\{e_1, \dots, e_d\}$ be the standard orthonormal basis of \mathbb{R}^d . The operator $*$ is the bijective linear map satisfying

$$*(e_{i_1} \wedge \dots \wedge e_{i_k}) = \text{sgn}(i_1, \dots, i_d) e_{i_{k+1}} \wedge \dots \wedge e_{i_d}$$

for all $1 \leq i_1 < \dots < i_k \leq d$, where $1 \leq i_{k+1} < \dots < i_d \leq d$ are such that $\{i_{k+1}, \dots, i_d\} = \{1, \dots, d\} \setminus \{i_1, \dots, i_k\}$, and $\text{sgn}(i_1, \dots, i_d) = 1$ if (i_1, \dots, i_d) is an even permutation of $\{1, \dots, d\}$ and $\text{sgn}(i_1, \dots, i_d) = -1$ otherwise. Recall that the inner product on $\wedge^k \mathbb{R}^d$ is defined by setting $\langle v, w \rangle = *(v \wedge *w)$ for all $v, w \in \wedge^k \mathbb{R}^d$. It is straightforward to see that $*(v) = (-1)^{k(d-k)} v$ and $v \wedge *w = (-1)^{k(d-k)} *v \wedge w$ for all $v, w \in \wedge^k \mathbb{R}^d$ and hence, $\|v\| = \|*v\|$ for all $v \in \wedge^k \mathbb{R}^d$. For a more detailed treatment, the reader is referred e.g. to [20, §3.2].

Observe first that if $A \in GL_d(\mathbb{R})$ and the vectors $v, w \in \mathbb{R}^d$ are perpendicular, then also the vectors $A^T v, A^{-1} w \in \mathbb{R}^d$ are perpendicular. Let v_1, \dots, v_{d-k} be an orthonormal basis of V and v_{d-k+1}, \dots, v_d be an orthonormal basis of V^\perp . Moreover, let g_1, \dots, g_{d-k} and g'_1, \dots, g'_k be the vectors obtained from the Gram-Schmidt orthogonalization of $(A_{\mathbf{i}|_n})^{-1} v_1, \dots, (A_{\mathbf{i}|_n})^{-1} v_{d-k}$ and $(A_{\mathbf{i}|_n})^T v_{d-k+1}, \dots, (A_{\mathbf{i}|_n})^T v_d$, respectively. This means that

$$\begin{aligned} g_1 &= (A_{\mathbf{i}|_n})^{-1} v_1, \\ g_2 &= (A_{\mathbf{i}|_n})^{-1} v_1 - c_{1,1} g_1, \\ &\vdots \\ g_{d-k} &= (A_{\mathbf{i}|_n})^{-1} v_{d-k} - c_{d-k,1} g_1 - \dots - c_{d-k,d-k-1} g_{d-k-1} \end{aligned}$$

and

$$\begin{aligned} g'_1 &= (A_{\mathbf{i}|_n})^T v_{d-k+1}, \\ g'_2 &= (A_{\mathbf{i}|_n})^T v_{d-k+2} - c'_{1,1} g'_1, \\ &\vdots \\ g'_k &= (A_{\mathbf{i}|_n})^T v_d - c'_{k,1} g'_1 - \dots - c_{k,k-1} g'_{k-1} \end{aligned}$$

with appropriate choices of the constants $c_{i,j}$ and $c'_{i,j}$. Hence,

$$(A_{\mathbf{i}|_n})^{-1} v_1 \wedge \dots \wedge (A_{\mathbf{i}|_n})^{-1} v_{d-k} = g_1 \wedge \dots \wedge g_{d-k}$$

and

$$(A_{\mathbf{i}|_n})^T v_{d-k+1} \wedge \dots \wedge (A_{\mathbf{i}|_n})^T v_d = g'_1 \wedge \dots \wedge g'_k,$$

and $\{g_1, \dots, g_{d-k}, g'_1, \dots, g'_k\}$ is an orthogonal basis of \mathbb{R}^d . Therefore,

$$0 \neq (g_1 \wedge \dots \wedge g_{d-k}) \wedge (g'_1 \wedge \dots \wedge g'_k) = \langle g_1 \wedge \dots \wedge g_{d-k}, *(g'_1 \wedge \dots \wedge g'_k) \rangle e_1 \wedge \dots \wedge e_d.$$

Since any g'_i is perpendicular to any g_j , we must have

$$*(g'_1 \wedge \cdots \wedge g'_k) = C g_1 \wedge \cdots \wedge g_{d-k}$$

for some constant $C > 0$. Since

$$\begin{aligned} \det(A_{\mathbf{i}|_n})^T v_1 \wedge \cdots \wedge v_d &= ((A_{\mathbf{i}|_n})^T v_1 \wedge \cdots \wedge (A_{\mathbf{i}|_n})^T v_{d-k}) \wedge ((A_{\mathbf{i}|_n})^T v_{d-k+1} \wedge \cdots \wedge (A_{\mathbf{i}|_n})^T v_d) \\ &= \langle (A_{\mathbf{i}|_n})^T v_1 \wedge \cdots \wedge (A_{\mathbf{i}|_n})^T v_{d-k}, *(A_{\mathbf{i}|_n})^T v_{d-k+1} \wedge \cdots \wedge (A_{\mathbf{i}|_n})^T v_d \rangle e_1 \wedge \cdots \wedge e_d \\ &= C \langle (A_{\mathbf{i}|_n})^T v_1 \wedge \cdots \wedge (A_{\mathbf{i}|_n})^T v_{d-k}, (A_{\mathbf{i}|_n})^{-1} v_1 \wedge \cdots \wedge (A_{\mathbf{i}|_n})^{-1} v_{d-k} \rangle e_1 \wedge \cdots \wedge e_d \\ &= C \langle v_1 \wedge \cdots \wedge v_{d-k}, v_1 \wedge \cdots \wedge v_{d-k} \rangle e_1 \wedge \cdots \wedge e_d \\ &= C e_1 \wedge \cdots \wedge e_d \end{aligned}$$

we have $|C| = |\det(A_{\mathbf{i}|_n})^T|$. Therefore, by (3.1) and the Oseledets' decomposition,

$$\begin{aligned} \sum_{j=k+1}^d \chi_j(\mathbf{A}, \nu) &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \|((A_{\mathbf{i}|_n})^{-1})^{\wedge(d-k)} | \wedge^{d-k} V \| \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \| (A_{\mathbf{i}|_n})^{-1} v_1 \wedge \cdots \wedge (A_{\mathbf{i}|_n})^{-1} v_{d-k} \| \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{\| *(A_{\mathbf{i}|_n})^T v_{d-k+1} \wedge \cdots \wedge (A_{\mathbf{i}|_n})^T v_d \|}{\det(A_{\mathbf{i}|_n})^T} \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \| (A_{\mathbf{i}|_n})^T v_{d-k+1} \wedge \cdots \wedge (A_{\mathbf{i}|_n})^T v_d \| + \sum_{j=1}^d \chi_j(\mathbf{A}, \nu) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \log \| ((A_{\mathbf{i}|_n})^T)^{\wedge k} | \wedge^k V \| + \sum_{j=1}^d \chi_j(\mathbf{A}, \nu) \end{aligned}$$

for $\mu_F^{d-k} \times \nu$ -almost every $(V, \mathbf{i}) \in G(d, d-k) \times \Sigma$. This completes the proof of the lemma. \square

Let $\mathbf{A} \in GL_d(\mathbb{R})^N$ be a tuple of contractive matrices and $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^d)^N$. Let \mathcal{U} be a parameter space equipped with a measure m such that each $u \in \mathcal{U}$ is a mapping $u: \mathbb{R}^d \rightarrow \mathbb{R}^d$. We will use this parametrised family of mappings to modify the self-affine iterated function system $\Phi_{\mathbf{A}, \mathbf{v}}$ by replacing it with $\Phi_{\mathbf{A}, u(\mathbf{v})}$, where $u(\mathbf{v}) = (u(v_1), \dots, u(v_N))$. We say that the pair (\mathcal{U}, m) satisfies the *modified self-affine transversality condition* for \mathbf{A} , if there exists a constant $C > 0$ such that for every proper subspace V of \mathbb{R}^d and $t > 0$ it holds that

$$m(\{u \in \mathcal{U} : |\text{proj}_V \pi_{\mathbf{A}, u(\mathbf{v})}(\mathbf{i}) - \text{proj}_V \pi_{\mathbf{A}, u(\mathbf{v})}(\mathbf{j})| < t\}) \leq C \prod_{i=1}^{\dim V} \min \left\{ 1, \frac{t}{\alpha_i(\text{proj}_V A_{\mathbf{i} \wedge \mathbf{j}})} \right\}$$

for all $\mathbf{i}, \mathbf{j} \in \Sigma$ with $\mathbf{i} \neq \mathbf{j}$. We note that if $V = \mathbb{R}^d$, then this condition is the self-affine transversality condition of Jordan, Pollicott and Simon [16].

Lemma 3.2. *Let $\mathbf{v} \in (\mathbb{R}^d)^N$ and $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$ be a tuple of contractive matrices. If (\mathcal{U}, m) satisfies the modified transversality condition for \mathbf{A} , then there exists a constant $C > 0$ such that*

$$\int \frac{dm(u)}{|\text{proj}_V(\pi_{\mathbf{A}, u(\mathbf{v})}(\mathbf{i}) - \text{proj}_V(\pi_{\mathbf{A}, u(\mathbf{v})}(\mathbf{j}))|^s} \leq \frac{C}{\varphi^s(\text{proj}_V A_{\mathbf{i} \wedge \mathbf{j}})}$$

for every proper subspace V of \mathbb{R}^d and for all $0 \leq s < \dim V$.

Proof. The proof is a slight modification of the proof of [16, Lemma 4.5] and hence omitted. \square

We will use the above lemma in the proof of the following proposition which is a key observation related to the modified self-affine transversality condition.

Proposition 3.3. *Let $\mathbf{v} \in (\mathbb{R}^d)^N$, $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$ be a tuple of contractive matrices, and ν be an ergodic measure on Σ . Assume that the $(d, d-k)$ -Furstenberg measure $\mu_F^{d,d-k}$ with respect to ν and \mathbf{A} exists for all $k \in \{1, \dots, d-1\}$. If (\mathcal{U}, m) satisfies the modified self-affine transversality condition for \mathbf{A} , then*

$$\underline{\dim}_{\mathbb{H}} \text{proj}_{V^\perp} \pi_{\mathbf{A}, u(\mathbf{v})} \nu = \min\{k, \dim_{\mathbb{L}} \nu\}$$

for $\mu_F^{d,d-k}$ -almost all $V \in G(d, d-k)$, for all $k \in \{1, \dots, d-1\}$, and for m -almost all $u \in \mathcal{U}$.

Proof. To simplify notation, we denote $\chi_j(\mathbf{A}, \nu)$ by χ_j and $\pi_{\mathbf{A}, u(\mathbf{v})}$ by π_u . Let $s < \min\{\dim_{\mathbb{L}} \nu, k\}$. By standard methods, it suffices to show that for ν -almost every $\mathbf{i} \in \Sigma$ it holds that

$$\int |\text{proj}_{V^\perp} \pi_u(\mathbf{i}) - \text{proj}_{V^\perp} \pi_u(\mathbf{j})|^{-s} d\nu(\mathbf{j}) < \infty$$

for $\mu_F^{d,d-k}$ -almost all $V \in G(d, d-k)$ and m -almost all $u \in \mathcal{U}$. For this to hold, it is enough to prove that for every $\varepsilon > 0$ small enough there exist sets $E_1 \subset \Sigma$ with $\nu(E_1) > 1 - \varepsilon$ and $E_2 \subset G(d, d-k) \times \Sigma$ with $\mu_F^{d,d-k} \times \nu(E_2) > 1 - \varepsilon$ such that

$$\iiint \frac{d\nu|_{E_1}(\mathbf{i}) d(\mu_F^{d,d-k} \times \nu)|_{E_2}(V, \mathbf{j}) dm(u)}{|\text{proj}_{V^\perp} \pi_u(\mathbf{i}) - \text{proj}_{V^\perp} \pi_u(\mathbf{j})|^s} < \infty.$$

Fix $0 < \varepsilon < \chi_{[s]}(\dim_{\mathbb{L}} \nu - s)/2$. By Egorov's Theorem, Shannon-McMillan-Breiman Theorem, and Lemma 3.1, there exist $C \geq 1$, $E_1 \subset \Sigma$ with $\nu(E_1) > 1 - \varepsilon$, and $E_2 \subset G(d, d-k) \times \Sigma$ with $\mu_F^{d,d-k} \times \nu(E_2) > 1 - \varepsilon$ such that

$$\begin{aligned} C^{-1} e^{-n(h_\nu + \varepsilon)} &\leq \nu([\mathbf{i}]_n) \leq C e^{-n(h_\nu - \varepsilon)}, \\ C^{-1} e^{-n(\chi_1 + \dots + \chi_{[s]} + (s - [s])\chi_{[s]} + \varepsilon)} &\leq \varphi^s(\text{proj}_{V^\perp} A_{\mathbf{j}|_n}) \leq C e^{-n(\chi_1 + \dots + \chi_{[s]} + (s - [s])\chi_{[s]} - \varepsilon)} \end{aligned}$$

for every $\mathbf{i} \in E_1$ and $(V, \mathbf{j}) \in E_2$. By Fubini's Theorem and Lemma 3.2, we have

$$\begin{aligned} \iiint \frac{d\nu|_{E_1}(\mathbf{i}) d(\mu_F^{d,d-k} \times \nu)|_{E_2}(V, \mathbf{j}) dm(u)}{|\text{proj}_{V^\perp} \pi_u(\mathbf{i}) - \text{proj}_{V^\perp} \pi_u(\mathbf{j})|^s} &= \iiint \frac{dm(u) d\nu|_{E_1}(\mathbf{i}) d(\mu_F^{d,d-k} \times \nu)|_{E_2}(V, \mathbf{j})}{|\text{proj}_{V^\perp} \pi_u(\mathbf{i}) - \text{proj}_{V^\perp} \pi_u(\mathbf{j})|^s} \\ &\leq C' \iint \frac{d(\mu_F^{d,d-k} \times \nu)|_{E_2}(V, \mathbf{j}) d\nu|_{E_1}(\mathbf{i})}{\varphi^s(\text{proj}_V A_{\mathbf{i} \wedge \mathbf{j}})} \end{aligned}$$

for some constant $C' > 0$. By decomposing the space $\Sigma \times G(d, d-k)$ into $\{\mathbf{i}\} \times G(d, d-k)$ and $\bigcup_{n=0}^{\infty} \{\mathbf{j} : |\mathbf{i} \wedge \mathbf{j}| = n\} \times G(d, d-k)$, we get

$$\begin{aligned} C' \iint \frac{d(\mu_F^{d,d-k} \times \nu)|_{E_2}(V, \mathbf{j}) d\nu|_{E_1}(\mathbf{i})}{\varphi^s(\text{proj}_V A_{\mathbf{i} \wedge \mathbf{j}})} &\leq CC' \sum_{n=0}^{\infty} \int e^{n(\chi_1 + \dots + \chi_{[s]} + (s - [s])\chi_{[s]} + \varepsilon)} (\mu_F^{d,d-k} \times \nu)|_{E_2}(G(d, d-k) \times [\mathbf{i}]_n) d\nu|_{E_1}(\mathbf{i}) \\ &= CC' \sum_{n=0}^{\infty} e^{n(\chi_1 + \dots + \chi_{[s]} + (s - [s])\chi_{[s]} + \varepsilon)} \sum_{|\mathbf{k}|=n} (\mu_F^{d,d-k} \times \nu)|_{E_2}(G(d, d-k) \times [\mathbf{k}]) \nu|_{E_1}([\mathbf{k}]) \\ &\leq C^2 C' \sum_{n=0}^{\infty} e^{n(\chi_1 + \dots + \chi_{[s]} + (s - [s])\chi_{[s]} + \varepsilon)} e^{-n(h_\nu - \varepsilon)} \sum_{|\mathbf{k}|=n} (\mu_F^{d,d-k} \times \nu)|_{E_2}(G(d, d-k) \times [\mathbf{k}]) \\ &\leq C^2 C' \sum_{n=0}^{\infty} e^{n(\chi_1 + \dots + \chi_{[s]} + (s - [s])\chi_{[s]} + \varepsilon)} e^{-n(h_\nu - \varepsilon)} \end{aligned}$$

Since $s\chi_{[s]} < \chi_{[s]} \dim_{\mathbb{L}} \nu - 2\varepsilon = [s]\chi_{[s]} + h_\nu - \sum_{j=1}^{[s]} \chi_j - 2\varepsilon$ we have finished the proof. \square

We say that an ergodic measure ν on Σ satisfies the *Ledrappier-Young formula* for a tuple $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})$ of contractive matrices if

$$\dim \pi_{\mathbf{A}, \mathbf{v}} \nu = \sum_{k=1}^{d-1} \frac{\chi_{k+1}(\mathbf{A}, \nu) - \chi_k(\mathbf{A}, \nu)}{\chi_d(\mathbf{A}, \nu)} \dim \text{proj}_{V_k^\perp} \pi_{\mathbf{A}, \mathbf{v}} \nu + \frac{h_\nu - H_{\mathbf{A}, \mathbf{v}, \nu}}{\chi_d(\mathbf{A}, \nu)}$$

for all $\mathbf{v} \in (\mathbb{R}^d)^N$ and for $\mu_F^1 \times \dots \times \mu_F^{d-1}$ -almost all $(V_1, \dots, V_{d-1}) \in G(d, 1) \times \dots \times G(d, d-1)$. Here $H_{\mathbf{A}, \mathbf{v}, \nu}$ denotes the conditional entropy defined in [3, §2]. We omit its definition since $H_{\mathbf{A}, \mathbf{v}, \nu} = 0$ in our considerations; see [3, Corollary 2.8].

Lemma 3.4. *If a Bernoulli measure ν on Σ has simple Lyapunov spectrum, then ν satisfies the Ledrappier-Young formula. Moreover, every ergodic quasi-Bernoulli measure ν on Σ satisfies the Ledrappier-Young formula for every tuple $\mathbf{A} \in GL_d(\mathbb{R})^N$ of contractive matrices satisfying the totally dominated splitting condition.*

Proof. The claims follow immediately from [3, Theorems 2.3 and 2.6]. \square

The next theorem shows that, under the strong separation condition, the Ledrappier-Young formula and the modified self-affine transversality condition together guarantee the desired dimension formula.

Theorem 3.5. *Let $\mathbf{v} \in (\mathbb{R}^d)^N$ and ν be an ergodic measure on Σ satisfying the Ledrappier-Young formula for a tuple $\mathbf{A} \in GL_d(\mathbb{R})^N$ of contractive matrices. Assume that (\mathcal{U}, m) satisfies the modified self-affine transversality condition for \mathbf{A} . If $\Phi_{\mathbf{A}, u(\mathbf{v})}$ satisfies the strong separation condition for all $u \in \mathcal{U}$ or $\dim_{\mathbb{L}} \nu \leq d-1$, then*

$$\dim \pi_{\mathbf{A}, u(\mathbf{v})} \nu = \dim_{\mathbb{L}} \nu$$

for m -almost all $u \in \mathcal{U}$.

Proof. By the Ledrappier-Young formula and Proposition 3.3, we have

$$\begin{aligned} \dim \pi_{\mathbf{A}, u(\mathbf{v})} \nu &= \sum_{k=1}^{d-1} \frac{\chi_{k+1}(\mathbf{A}, \nu) - \chi_k(\mathbf{A}, \nu)}{\chi_d(\mathbf{A}, \nu)} \dim \text{proj}_{V_k^\perp} \pi_{\mathbf{A}, u(\mathbf{v})} \nu + \frac{h_\nu - H_{\mathbf{A}, u(\mathbf{v}), \nu}}{\chi_d(\mathbf{A}, \nu)} \\ &= \sum_{k=1}^{d-1} \frac{\chi_{k+1}(\mathbf{A}, \nu) - \chi_k(\mathbf{A}, \nu)}{\chi_d(\mathbf{A}, \nu)} \min\{k, \dim_{\mathbb{L}} \nu\} + \frac{h_\nu - H_{\mathbf{A}, u(\mathbf{v}), \nu}}{\chi_d(\mathbf{A}, \nu)} \\ &= \dim_{\mathbb{L}} \nu - \frac{H_{\mathbf{A}, u(\mathbf{v}), \nu}}{\chi_d(\mathbf{A}, \nu)} \end{aligned}$$

for $\mu_F^1 \times \dots \times \mu_F^{d-1}$ -almost every $(V_1, \dots, V_{d-1}) \in G(d, 1) \times \dots \times G(d, d-1)$ and for m -almost every $u \in \mathcal{U}$. If $\Phi_{\mathbf{A}, u(\mathbf{v})}$ satisfies the strong separation condition, then [3, Corollary 2.8] implies that $H_{\mathbf{A}, u(\mathbf{v}), \nu} = 0$ and the claim follows. Furthermore, if $\dim_{\mathbb{L}} \nu \leq d-1$, then, by choosing a typical $V \in G(d, 1)$ in the sense of Proposition 3.3, we have $\dim_{\mathbb{L}} \nu = \dim \text{proj}_{V^\perp} \pi_{\mathbf{A}, u(\mathbf{v})} \nu \leq \dim \pi_{\mathbf{A}, u(\mathbf{v})} \nu = \dim_{\mathbb{L}} \nu - H_{\mathbf{A}, u(\mathbf{v}), \nu} / \chi_d(\mathbf{A}, \nu)$. Therefore $H_{\mathbf{A}, u(\mathbf{v}), \nu} = 0$ also in this case. \square

Remark 3.6. We indicate that some assumptions on the matrix norms in the statement of Theorem A are necessary, at least for our approach of proof. As we will see in §5, the strategy of the proof for Theorem A relies heavily on Theorem 3.5. In order to apply Theorem 3.5 on a planar self-affine system, we only need to check that the strong separation condition and the modified self-affine transversality condition hold, because by Lemma 3.4 the Ledrappier-Young formula holds for Bernoulli measures of planar self-affine systems.

We consider the example of Przytycki and Urbański [23, Theorem 8]. They investigate an IFS $\Phi_{\mathbf{A}, \mathbf{v}}$ with $\mathbf{A} = (A_1, A_2)$, where

$$A_i = \begin{pmatrix} \lambda & 0 \\ 0 & \gamma \end{pmatrix}$$

such that $\lambda > 1/2 > \gamma$ and λ^{-1} is a Pisot number, and $\mathbf{v} = \{(0, 0), (1 - \lambda, 1 - \gamma)\}$. They prove that in this case, for the equidistributed Bernoulli measure μ , $\dim_{\mathbb{H}} \pi\mu < \dim_{\mathbb{L}} \mu$. Notice that here, the strong separation condition holds for $\Phi_{\mathbf{A}, \mathbf{v}}$ and by varying λ and γ we can break the condition (2.9).

We will see in Lemma 5.2 below that for any system satisfying the condition (2.9) the modified self-affine transversality condition does hold for $O(d)$ equipped with the Haar measure (recall that $O(d) \subset GL_2(\mathbb{R})$ denotes the orthogonal group). However, this is not the case here. For $G \in O(d)$, the Furstenberg measure for the system $\Phi_{\mathbf{A}, G(\mathbf{v})}$ is the Dirac measure supported on $V = \text{span}\{(0, 1)\}$ and $\dim_{\mathbb{H}} \text{proj}_{V^\perp} \pi_G \mu < 1$. Therefore $\dim_{\mathbb{H}} \pi_G \mu < \dim_{\mathbb{L}} \mu$ for all $G \in O(d)$. Observe that therefore the claim of Theorem 3.5 does not hold, and consequently, the modified self-affine transversality condition does not hold.

We say that two affine iterated function systems Φ and Ψ are *equivalent* if the self-affine set of Ψ is an isometric copy of the self-affine set of Φ . This equivalence obviously preserves all the dimensional and separation properties of the self-affine set.

Let $\eta > 0$ and m be a probability measure on a group \mathcal{U} contained in $\{u \in GL_d(\mathbb{R}) : \eta < \alpha_d(u) \leq \alpha_1(u) < \eta^{-1}\}$. Notice that all the eigenvalues of the matrices in \mathcal{U} are one in modulus. We will introduce a method which allows us to handle matrices as parameters when using the modified self-affine transversality condition. If $u \in \mathcal{U}$ and $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$, then we define $u(\mathbf{A}) = (u^{-1}A_1u, \dots, u^{-1}A_Nu)$. Recall that $u(\mathbf{v}) = (uv_1, \dots, uv_N)$ for $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^d)^N$.

Lemma 3.7. *If $\mathbf{v} \in (\mathbb{R}^d)^N$ and $\mathbf{A} \in GL_d(\mathbb{R})^N$, then the affine iterated function systems $\Phi_{u(\mathbf{A}), \mathbf{v}}$ and $\Phi_{\mathbf{A}, u(\mathbf{v})}$ are equivalent for all $u \in \mathcal{U}$.*

Proof. Recall that $f(A, v)$ is the affine mapping $x \mapsto Ax + v$. Let $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$ and $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^d)^N$. Clearly, for each $x \in \mathbb{R}^d$ we have

$$f(u^{-1}A_iu, v_i)(x) = f(u^{-1}, 0)(f(A_i, u(v_i))(f(u, 0)(x)))$$

and $f(u^{-1}, 0)(f(u, 0)(x)) = x$. Thus, $\Phi_{u(\mathbf{A}), \mathbf{v}}$ and $\Phi_{\mathbf{A}, u(\mathbf{v})}$ are equivalent. \square

We define a partition of $GL_d(\mathbb{R})^N$ by setting an equivalence relation on $GL_d(\mathbb{R})^N$ as follows: if $\mathbf{A}, \mathbf{B} \in GL_d(\mathbb{R})^N$, then we say that $\mathbf{A} \sim \mathbf{B}$ if and only if there exists $u \in \mathcal{U}$ such that $\mathbf{B} = u(\mathbf{A})$. Since \mathcal{U} is a group \sim defines an equivalence relation. Hence

$$\mathcal{P}(\mathbf{A}) = \{\mathbf{B} \in GL_d(\mathbb{R})^N : \mathbf{A} \sim \mathbf{B}\}.$$

is a partition of $GL_d(\mathbb{R})^N$. Since $GL_d(\mathbb{R})^N$ equipped with the distance $d(\mathbf{A}, \mathbf{B}) = \max_i \|A_i - B_i\|$ is a separable metric space (matrix tuples with rational entries form a countable dense subset), the σ -algebra \mathcal{B} of Borel sets of $GL_d(\mathbb{R})^N$ is countably generated by $\{X_1, X_2, \dots\}$, where X_i is open in $GL_d(\mathbb{R})^N$. By defining $X'_i = \bigcup_{\mathbf{A} \in X_i} \mathcal{P}(\mathbf{A})$, the set $\{X'_1, X'_2, \dots\}$ generates the σ -algebra $\mathcal{B}_{\mathcal{P}} = \{X \in \mathcal{B} : X = \bigcup_{\mathbf{A} \in X} \mathcal{P}(\mathbf{A})\}$. Indeed, we clearly have $\sigma(\{X'_1, X'_2, \dots\}) \subset \mathcal{B}_{\mathcal{P}}$. On the other hand, if $X \in \mathcal{B}_{\mathcal{P}}$ is open, then $X = \bigcup_j X_{ij}$. But now $X = \bigcup_{\mathbf{A} \in X} \mathcal{P}(\mathbf{A}) = \bigcup_j \bigcup_{\mathbf{A} \in X_{ij}} \mathcal{P}(\mathbf{A}) = \bigcup_j X'_j$ and thus $\sigma(\{X'_1, X'_2, \dots\}) \supset \mathcal{B}_{\mathcal{P}}$.

By Rokhlin's disintegration theorem (see [25] and [26]), for any finite measure \mathcal{M} on $GL_d(\mathbb{R})^N$, there exists a family of measures $\{\mathcal{M}^{\mathcal{P}(\mathbf{A})}\}$ such that $\mathcal{M}^{\mathcal{P}(\mathbf{A})}$ is uniquely defined for \mathcal{M} -almost every \mathbf{A} , $\mathcal{M}^{\mathcal{P}(\mathbf{A})}$ is supported on $\mathcal{P}(\mathbf{A})$, and, moreover,

$$\mathcal{M} = \int \mathcal{M}^{\mathcal{P}(\mathbf{A})} d\mathcal{M}(\mathbf{A}).$$

The combination of Lemma 3.7 and the definition of \mathcal{P} implies that we have the same Furstenberg measures for all u on the partition elements. More precisely, for each $\mathbf{B} \in \mathcal{P}(\mathbf{A})$ there is $u \in \mathcal{U}$ such that $\Phi_{\mathbf{B}, \mathbf{v}}$ is equivalent to $\Phi_{\mathbf{A}, u(\mathbf{v})}$ whose Furstenberg measures are independent of $u \in \mathcal{U}$.

Theorem 3.8. *Let $\mathbf{v} \in (\mathbb{R}^d)^N$, $\eta > 0$, and m be a probability measure on a group \mathcal{U} contained in $\{u \in GL_d(\mathbb{R}) : \eta < \alpha_d(u) \leq \alpha_1(u) < \eta^{-1}\}$. Furthermore, let \mathcal{M} be a probability measure on*

$GL_d(\mathbb{R})^N$ and let $\xi_{\mathbf{A}}: \mathcal{U} \rightarrow GL_d(\mathbb{R})^N$, $\xi_{\mathbf{A}}(u) = u(\mathbf{A})$, be such that $\mathcal{M}^{\mathcal{P}(\mathbf{A})} = \xi_{\mathbf{A}} m$ for \mathcal{M} -almost all \mathbf{A} . Assume that ν is an ergodic measure on Σ satisfying the Ledrappier-Young formula and (\mathcal{U}, m) satisfies the modified self-affine transversality condition for \mathcal{M} -almost all tuples \mathbf{A} of contractive matrices. Then

$$\dim_{\mathbb{H}} \pi_{\mathbf{A}, \nu} \nu = \dim_{\mathbb{L}} \nu$$

for \mathcal{M} -almost all tuples $\mathbf{A} \in GL_d(\mathbb{R})^N$ of contractive matrices for which $\Phi_{\mathbf{A}, \nu}$ satisfies the strong separation condition or $\dim_{\mathbb{L}} \nu \leq d - 1$.

Proof. Observe that, by Theorem 3.5, if $\Phi_{\mathbf{A}, u(\nu)}$ satisfies the strong separation condition for all $u \in \mathcal{U}$ and \mathcal{M} -almost all tuples \mathbf{A} of contractive matrices or $\dim_{\mathbb{L}} \nu \leq d - 1$, then

$$\dim \pi_{\mathbf{A}, u(\nu)} \nu = \dim_{\mathbb{L}} \nu$$

for m -almost all $u \in \mathcal{U}$ and \mathcal{M} -almost all tuples \mathbf{A} of contractive matrices. By Lemma 3.7, we have $\dim_{\mathbb{H}} \pi_{u(\mathbf{A}), \nu} \nu = \dim_{\mathbb{H}} \pi_{\mathbf{A}, u(\nu)} \nu$ for all $u \in \mathcal{U}$. Thus, by Rokhlin's disintegration theorem,

$$\mathcal{M} = \int \xi_{\mathbf{A}} m \, d\mathcal{M}(\mathbf{A}),$$

and hence, $\dim_{\mathbb{H}} \pi_{\mathbf{A}, \nu} \nu = \dim_{\mathbb{L}} \nu$ for \mathcal{M} -almost all tuples \mathbf{A} of contractive matrices. \square

4. DIMENSION VIA SUB-SYSTEMS

Let \mathbb{A} be the collection of all tuples $\mathbf{A} \in GL_d(\mathbb{R})^N$ of contractive matrices that satisfy $\chi_i(\mathbf{A}, \mu) \neq \chi_j(\mathbf{A}, \mu)$ for $i \neq j$ where μ is an ergodic s -equilibrium state of \mathbf{A} and $s = \dim_{\text{aff}}(\mathbf{A})$. We note that the functions $\mathbf{A} \mapsto \chi_i(\mathbf{A}, \nu)$ and $\nu \mapsto \chi_i(\mathbf{A}, \nu)$ are lower semi-continuous in the usual and weak*-topology. Thus, by [19, Propositions 3.4 and 3.6] and [10, Proposition 5.2], the set \mathbb{A} is Borel measurable in $GL_d(\mathbb{R})^N$.

In this section, we show that equilibrium states can be approximated by step- n Bernoulli measure arbitrarily well on \mathbb{A} . Each affine iterated function system contains a well-approximating sub-system in which the Ledrappier-Young formula holds. In our setting, this observation yields a dimension formula for the self-affine set. Motivation for this section is purely technical: at the moment, we do not know whether ergodic equilibrium states satisfy the Ledrappier-Young formula – Lemma 3.4 gathers the current knowledge on this problem.

Proposition 4.1. *For every $\mathbf{A} \in \mathbb{A}$ and $\varepsilon > 0$ there exist $n \in \mathbb{N}$ and a step- n Bernoulli measure ν such that $\dim_{\mathbb{L}} \nu \geq \dim_{\text{aff}} \mathbf{A} - \varepsilon$.*

Proof. Fix $\mathbf{A} = (A_1, \dots, A_N) \in \mathbb{A}$, define $s = \dim_{\text{aff}} \mathbf{A}$, and let μ be an ergodic s -equilibrium state of \mathbf{A} . By Feng and Shmerkin [10, Theorem 3.3], there exist $\eta > 0$ and an infinite set $S \subset \mathbb{N}$ such that for every $n \in S$ there is $\Gamma_n \subset \Sigma_n$ with $\sum_{\mathbf{i} \in \Gamma_n} \mu([\mathbf{i}]) \geq \eta$. Moreover, since all the Lyapunov exponents are distinct it also follows that each $(A_{\mathbf{i}})_{\mathbf{i} \in \Gamma_n}$ satisfies the totally dominated splitting condition and hence, via e.g. [10, Lemma 2.1], there exists a constant $C \geq 1$ such that

$$\varphi^s(A_{\mathbf{i}}) \varphi^s(A_{\mathbf{j}}) \leq C \varphi^s(A_{\mathbf{ij}}) \tag{4.1}$$

for all $\mathbf{i}, \mathbf{j} \in \bigcup_{k=1}^{\infty} \Gamma_n^k$, $k \in \mathbb{N}$, $n \in S$, and $s \geq 0$. For each $n \in S$, let us choose s_n such that

$$\sum_{\mathbf{i} \in \Gamma_n} \varphi^{s_n}(A_{\mathbf{i}}) = 1.$$

We will show that $s_n \rightarrow s$ as $n \rightarrow \infty$.

Applying Egorov's Theorem, it follows from (2.8) and Shannon-McMillan-Breiman Theorem that there exists a set $E \subset \Sigma$ with $\sum_{\mathbf{i} \in \Gamma_n} \mu([\mathbf{i}] \cap E) \geq \eta/2$ such that for every $\mathbf{i} \in \bigcup_{\mathbf{j} \in \Gamma_n} [\mathbf{j}] \cap E$

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{\mu([\mathbf{i}]_n)}{\varphi^s(A_{\mathbf{i}|_n})} = 0$$

uniformly. This implies that there exists a sequence $(c_n)_{n \in \mathbb{N}}$ of reals such that $\lim_{n \rightarrow \infty} \log c_n^{1/n} = 0$ and

$$c_n^{-1} \varphi^s(A_{\mathbf{i}}) \leq \mu([\mathbf{i}]) \leq c_n \varphi^s(A_{\mathbf{i}})$$

for all $\mathbf{i} \in \bigcup_{\mathbf{j} \in \Gamma_n} [\mathbf{j}] \cap E$ and $n \in S$. Therefore, by (2.2), we have

$$c_n^{-1} \eta/2 \leq c_n^{-1} \sum_{\mathbf{i} \in \Gamma_n} \mu([\mathbf{i}] \cap E) \leq \sum_{\mathbf{i} \in \Gamma_n} \varphi^s(A_{\mathbf{i}}) \leq \|\mathbf{A}\|^{n(s-s_n)} \sum_{\mathbf{i} \in \Gamma_n} \varphi^{s_n}(A_{\mathbf{i}}) = \|\mathbf{A}\|^{n(s-s_n)}$$

and thus,

$$s - s_n \leq \frac{\log c_n^{-1} \eta/2}{n \log \|\mathbf{A}\|} \quad (4.2)$$

for all $n \in S$. On the other hand, again by (2.2), we have

$$\sum_{\mathbf{i} \in \Sigma_n} \varphi^s(A_{\mathbf{i}}) \geq \sum_{\mathbf{i} \in \Gamma_n} \varphi^s(A_{\mathbf{i}}) \geq \mathbf{m}(\mathbf{A})^{n(s-s_n)} \sum_{\mathbf{i} \in \Gamma_n} \varphi^{s_n}(A_{\mathbf{i}}) = \mathbf{m}(\mathbf{A})^{n(s-s_n)}$$

and hence,

$$s - s_n \geq \frac{\log \sum_{\mathbf{i} \in \Sigma_n} \varphi^s(A_{\mathbf{i}})}{n \log \mathbf{m}(\mathbf{A})} \quad (4.3)$$

for all $n \in S$. By the definition of the affinity dimension, (4.2) and (4.3) show that $s_n \rightarrow s$ as $n \rightarrow \infty$.

Let $\nu_{\mathbf{p}_n}$ be the Bernoulli measure obtained from the probability vector $\mathbf{p}_n = (\varphi^{s_n}(A_{\mathbf{i}}))_{\mathbf{i} \in \Gamma_n}$. Observe that, by (4.1),

$$\begin{aligned} h_{\mathbf{p}_n} &= - \sum_{\mathbf{i} \in \Gamma_n} \nu_{\mathbf{p}_n}([\mathbf{i}]) \log \varphi^{s_n}(A_{\mathbf{i}}) \\ &= -\frac{1}{k} \sum_{\mathbf{i}_1, \dots, \mathbf{i}_k \in \Gamma_n} \nu_{\mathbf{p}_n}([\mathbf{i}_1]) \cdots \nu_{\mathbf{p}_n}([\mathbf{i}_k]) \log \varphi^{s_n}(A_{\mathbf{i}_1}) \cdots \varphi^{s_n}(A_{\mathbf{i}_k}) \\ &\geq -\frac{1}{k} \sum_{\mathbf{i}_1, \dots, \mathbf{i}_k \in \Gamma_n} \nu_{\mathbf{p}_n}([\mathbf{i}_1]) \cdots \nu_{\mathbf{p}_n}([\mathbf{i}_k]) \log C^{k-1} \varphi^{s_n}(A_{\mathbf{i}_1 \dots \mathbf{i}_k}) \end{aligned}$$

and similarly,

$$\begin{aligned} h_{\mathbf{p}_n} &= - \sum_{\mathbf{i} \in \Gamma_n} \nu_{\mathbf{p}_n}([\mathbf{i}]) \log \varphi^{s_n}(A_{\mathbf{i}}) \\ &\leq -\frac{1}{k} \sum_{\mathbf{i}_1, \dots, \mathbf{i}_k \in \Gamma_n} \nu_{\mathbf{p}_n}([\mathbf{i}_1]) \cdots \nu_{\mathbf{p}_n}([\mathbf{i}_k]) \log \varphi^{s_n}(A_{\mathbf{i}_1 \dots \mathbf{i}_k}). \end{aligned}$$

Letting $k \rightarrow \infty$ this implies

$$\dim_{\text{L}} \nu_{\mathbf{p}_n} = \lfloor s_n \rfloor + \frac{h_{\mathbf{p}_n} - \sum_{j=1}^{\lfloor s_n \rfloor} \chi_j(\mathbf{A}^n, \mathbf{p}_n)}{\chi_{\lfloor s_n \rfloor}(\mathbf{A}^n, \mathbf{p}_n)} \geq \frac{-\log C}{\chi_{\lfloor s_n \rfloor}(\mathbf{A}^n, \mathbf{p}_n)} + s_n \geq \frac{\log C}{n \log \mathbf{m}(\mathbf{A})} + s_n,$$

where $\mathbf{A}^n = (A_{\mathbf{i}})_{|\mathbf{i}|=n}$. This is what we wanted to show. \square

We will next transfer the previous proposition in the form we can use in our setting.

Theorem 4.2. *Let $\mathbf{v} \in (\mathbb{R}^d)^N$, $\eta > 0$, and m be a probability measure on a group \mathcal{U} contained in $\{u \in GL_d(\mathbb{R}) : \eta < \alpha_d(u) \leq \alpha_1(u) < \eta^{-1}\}$. Furthermore, let \mathcal{M} be a probability measure on $GL_d(\mathbb{R})^N$ such that $\mathcal{M}(\{\mathbf{A} \in GL_d(\mathbb{R})^N : \|\mathbf{A}\| < 1\} \setminus \mathbb{A}) = 0$ and let $\xi_{\mathbf{A}} : \mathcal{U} \rightarrow GL_d(\mathbb{R})^N$, $\xi_{\mathbf{A}}(u) = u(\mathbf{A})$, be such that $\mathcal{M}^{\mathcal{P}(\mathbf{A})} = \xi_{\mathbf{A}} m$ for \mathcal{M} -almost all \mathbf{A} . Assume that (\mathcal{U}, m) satisfies the modified self-affine transversality condition for \mathcal{M} -almost all tuples \mathbf{A} of contractive matrices. Then*

$$\dim_{\text{H}} E_{\mathbf{A}, \mathbf{v}} = \overline{\dim_{\text{M}}} E_{\mathbf{A}, \mathbf{v}} = \dim_{\text{aff}} \mathbf{A}$$

for \mathcal{M} -almost all tuples $\mathbf{A} \in GL_d(\mathbb{R})^N$ of contractive matrices for which $\Phi_{\mathbf{A}, \mathbf{v}}$ satisfies the strong separation condition or $\dim_{\text{aff}} \mathbf{A} \leq d - 1$.

Proof. Let us first show that for every $\mathbf{A} \in \mathbb{A}$ for which $\Phi_{\mathbf{A}, \mathbf{v}}$ satisfies the strong separation condition or $\dim_{\text{aff}} \mathbf{A} \leq d - 1$, and for every $\varepsilon > 0$ it holds that

$$\dim_{\mathbb{H}} E_{\mathbf{A}, u(\mathbf{v})} > \dim_{\text{aff}} \mathbf{A} - \varepsilon \quad (4.4)$$

for m -almost every $u \in \mathcal{U}$. Fix such a tuple $\mathbf{A} \in \mathbb{A}$ and $\varepsilon > 0$. Then, as in the proof of Proposition 4.1, we find a finite set $\Gamma \subset \Sigma_*$ and a Bernoulli measure ν on $\Gamma^{\mathbb{N}}$ such that $(A_{\mathbf{i}})_{\mathbf{i} \in \Gamma}$ satisfies the totally dominated splitting condition and $\dim_{\mathbb{L}} \nu \geq \dim_{\text{aff}} \mathbf{A} - \varepsilon$. By Lemma 3.4, ν satisfies the Ledrappier-Young formula. Hence, by Theorem 3.5, $\dim \pi_{\mathbf{A}, u(\mathbf{v})} \nu = \dim_{\mathbb{L}} \nu$ for m -almost every $u \in \mathcal{U}$ and we have finished the proof of (4.4).

By Lemma 3.7, we have $\dim_{\mathbb{H}} E_{\mathbf{A}, u(\mathbf{v})} = \dim_{\mathbb{H}} E_{u(\mathbf{A}), \mathbf{v}}$ for all $u \in \mathcal{U}$. It is also easy to see that $\dim_{\text{aff}} \mathbf{A} = \dim_{\text{aff}} u(\mathbf{A})$ for every $u \in \mathcal{U}$. Since \mathbb{A} has full measure with respect to \mathcal{M} , by Rokhlin's disintegration theorem, we thus have $\dim_{\mathbb{H}} E_{\mathbf{A}, \mathbf{v}} > \dim_{\text{aff}} \mathbf{A} - \varepsilon$ for \mathcal{M} -almost all tuples \mathbf{A} of contractive matrices for which $\Phi_{\mathbf{A}, \mathbf{v}}$ satisfies the strong separation condition or $\dim_{\text{aff}} \mathbf{A} \leq d - 1$. Since $\varepsilon > 0$ was arbitrary, the proof is complete. \square

5. THE PLANAR CASE

In this section, we prove Theorem A as an application of Theorems 3.8 and 4.2 by showing that the modified self-affine transversality condition holds for some pair (\mathcal{U}, m) for all $\mathbf{A} \in \mathcal{A}_{\mathbf{v}}$. Recall that if $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^2)^N$ is such that $v_i \neq v_j$ for every $i \neq j$, then

$$\mathcal{A}_{\mathbf{v}} = \left\{ \mathbf{A} \in GL_2(\mathbb{R})^N : 0 < \max_{i \neq j} \frac{\|A_i\| + \|A_j\|}{|v_i - v_j|} \cdot \frac{\|\mathbf{v}\|}{1 - \|\mathbf{A}\|} < \frac{\sqrt{2}}{2} \right\}.$$

Let $SO(2)$ be the special orthogonal group of $GL_2(\mathbb{R})$. Note that $SO(2) = \{u_{\alpha}\}_{\alpha \in \mathbb{R}}$, where

$$u_{\alpha} = \begin{pmatrix} \cos(\alpha) & -\sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{pmatrix}$$

is a rotation by an angle $\alpha \in \mathbb{R}$. To simplify notation, we will denote $\pi_{\mathbf{A}, u_{\alpha}(\mathbf{v})}$ by π_{α} . Furthermore, we denote the differential with respect to α , evaluated at α_0 , by $\partial_{\alpha=\alpha_0}$. The following is a transversality lemma suitable for our purposes.

Lemma 5.1. *Let $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^2)^N$ be such that $v_i \neq v_j$ for $i \neq j$ and $\mathbf{A} = (A_1, \dots, A_N) \in \mathcal{A}_{\mathbf{v}}$. Then there exists $\delta > 0$ such that for every $\alpha_0 \in [0, 2\pi]$, for every $v \in \mathbb{R}^2$ with $|v| = 1$, and for every $\mathbf{i}, \mathbf{j} \in \Sigma$ with $|\mathbf{i}|_1 \neq |\mathbf{j}|_1$*

$$|\langle v, \pi_{\alpha_0}(\mathbf{i}) - \pi_{\alpha_0}(\mathbf{j}) \rangle| \geq \delta \quad \text{or} \quad |\partial_{\alpha=\alpha_0} \langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle| \geq \delta.$$

Proof. Let us argue by contradiction. Suppose that for every $\delta > 0$ there exists $\alpha_0 \in [0, 2\pi]$, $v \in \mathbb{R}^2$ with $|v| = 1$ and $\mathbf{i}, \mathbf{j} \in \Sigma$ with $|\mathbf{i}|_1 \neq |\mathbf{j}|_1$ such that

$$|\langle v, \pi_{\alpha_0}(\mathbf{i}) - \pi_{\alpha_0}(\mathbf{j}) \rangle| < \delta \quad \text{and} \quad |\partial_{\alpha=\alpha_0} \langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle| < \delta.$$

Let $(\delta_n)_{n \in \mathbb{N}}$ be a sequence of positive reals such that $\delta_n \rightarrow 0$ as $n \rightarrow \infty$. By compactness, there exist $\alpha_0 \in [0, 2\pi]$, $v \in \mathbb{R}^2$ with $|v| = 1$ and $\mathbf{i}, \mathbf{j} \in \Sigma$ with $|\mathbf{i}|_1 \neq |\mathbf{j}|_1$ such that

$$|\langle v, \pi_{\alpha_0}(\mathbf{i}) - \pi_{\alpha_0}(\mathbf{j}) \rangle| = 0 \quad \text{and} \quad |\partial_{\alpha=\alpha_0} \langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle| = 0.$$

Since $\partial_{\alpha} u_{\alpha}^{-1} = u_{\alpha+\pi/2}^{-1}$ we have $\partial_{\alpha} \pi_{\alpha}(\mathbf{i}) = \pi_{\alpha+\pi/2}(\mathbf{i})$ for every $\mathbf{i} \in \Sigma$. Thus,

$$\begin{aligned} 0 &= \langle v, \pi_{\alpha_0}(\mathbf{i}) - \pi_{\alpha_0}(\mathbf{j}) \rangle \\ &= \langle u_{\alpha_0} v, v_{i_1} - v_{j_1} \rangle + \langle v, A_{i_1} \pi_{\alpha_0}(\sigma(\mathbf{i})) - A_{j_1} \pi_{\alpha_0}(\sigma(\mathbf{j})) \rangle \end{aligned}$$

and

$$\begin{aligned} 0 &= \partial_{\alpha=\alpha_0} \langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle \\ &= \langle u_{\alpha_0+\pi/2} v, v_{i_1} - v_{j_1} \rangle + \langle v, A_{i_1} \pi_{\alpha_0+\pi/2}(\sigma(\mathbf{i})) - A_{j_1} \pi_{\alpha_0+\pi/2}(\sigma(\mathbf{j})) \rangle. \end{aligned}$$

Hence,

$$\begin{aligned} |v_{i_1} - v_{j_1}|^2 &= (\langle v, A_{i_1} \pi_{\alpha_0}(\sigma(\mathbf{i})) - A_{j_1} \pi_{\alpha_0}(\sigma(\mathbf{j})) \rangle)^2 + (\langle v, A_{i_1} \pi_{\alpha_0 + \pi/2}(\sigma(\mathbf{i})) - A_{j_1} \pi_{\alpha_0 + \pi/2}(\sigma(\mathbf{j})) \rangle)^2 \\ &\leq (|A_{i_1} \pi_{\alpha_0}(\sigma(\mathbf{i}))| + |A_{j_1} \pi_{\alpha_0}(\sigma(\mathbf{j}))|)^2 + (|A_{i_1} \pi_{\alpha_0 + \pi/2}(\sigma(\mathbf{i}))| + |A_{j_1} \pi_{\alpha_0 + \pi/2}(\sigma(\mathbf{j}))|)^2 \\ &\leq 2(\|A_{i_1}\| + \|A_{j_1}\|)^2 \left(\frac{\|\mathbf{v}\|}{1 - \|\mathbf{A}\|} \right)^2. \end{aligned}$$

This contradicts the condition (2.9). \square

The following proposition shows that $(SO(2), \mathcal{L}^1)$ satisfies the modified self-affine transversality condition for all $\mathbf{A} \in \mathcal{A}_{\mathbf{v}}$.

Proposition 5.2. *Let $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^2)^N$ be such that $v_i \neq v_j$ for $i \neq j$ and $\mathbf{A} = (A_1, \dots, A_N) \in \mathcal{A}_{\mathbf{v}}$. Then there is a constant $C' > 0$ such that*

$$\mathcal{L}^1(\{\alpha \in [0, 2\pi] : |\text{proj}_{\theta} \pi_{\alpha}(\mathbf{i}) - \text{proj}_{\theta} \pi_{\alpha}(\mathbf{j})| < t\}) \leq C' \min\left\{1, \frac{t}{\alpha_1(\text{proj}_{\theta} A_{\mathbf{i} \wedge \mathbf{j}})}\right\}.$$

for all $\mathbf{i}, \mathbf{j} \in \Sigma$ with $\mathbf{i} \neq \mathbf{j}$, $t > 0$, and $\theta \in V(2, 1)$.

Proof. Fix $\mathbf{i}, \mathbf{j} \in \Sigma$ with $\mathbf{i} \neq \mathbf{j}$. Observe that for every $\theta \in V(2, 1)$ and $w \in \mathbb{R}^2$ we have

$$|\text{proj}_{\theta} w| = |\langle v, w \rangle|,$$

where $v \in \theta$ is so that $|v| = 1$. Thus, writing $n = |\mathbf{i} \wedge \mathbf{j}|$ we have

$$\begin{aligned} |\text{proj}_{\theta} \pi_{\alpha}(\mathbf{i}) - \text{proj}_{\theta} \pi_{\alpha}(\mathbf{j})| &= |\langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle| = |\langle v, A_{\mathbf{i} \wedge \mathbf{j}} \pi_{\alpha}(\sigma^n \mathbf{i}) - A_{\mathbf{i} \wedge \mathbf{j}} \pi_{\alpha}(\sigma^n \mathbf{j}) \rangle| \\ &= |A_{\mathbf{i} \wedge \mathbf{j}}^T v| \left| \left\langle \frac{A_{\mathbf{i} \wedge \mathbf{j}}^T v}{|A_{\mathbf{i} \wedge \mathbf{j}}^T v|}, \pi_{\alpha}(\sigma^n \mathbf{i}) - \pi_{\alpha}(\sigma^n \mathbf{j}) \right\rangle \right|. \end{aligned}$$

On the other hand,

$$\|\text{proj}_{\theta} A_{\mathbf{i} \wedge \mathbf{j}}\| = \|A_{\mathbf{i} \wedge \mathbf{j}}^T \text{proj}_{\theta}^T\| = \sup_{w \in \mathbb{R}^2} \frac{|A_{\mathbf{i} \wedge \mathbf{j}}^T \text{proj}_{\theta}^T w|}{|w|} = \sup_{w \in \mathbb{R}^2} \frac{|A_{\mathbf{i} \wedge \mathbf{j}}^T \text{proj}_{\theta} w|}{|\text{proj}_{\theta} w|} \frac{|\text{proj}_{\theta} w|}{|w|} = \|A_{\mathbf{i} \wedge \mathbf{j}}^T | \theta \|.$$

Hence, it suffices to prove that for any $v \in \mathbb{R}^2$ with $|v| = 1$

$$\mathcal{L}^1(\{\alpha \in [0, 2\pi] : |\langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle| < t\}) \leq C' t \tag{5.1}$$

for all $\mathbf{i}, \mathbf{j} \in \Sigma$ with $|\mathbf{i}|_1 \neq |\mathbf{j}|_1$.

Fix $v \in \mathbb{R}^2$ with $|v| = 1$ and $\mathbf{i}, \mathbf{j} \in \Sigma$ with $|\mathbf{i}|_1 \neq |\mathbf{j}|_1$. Write

$$H_t = \{\alpha \in [0, 2\pi] : |\langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle| < t\}$$

and let $\delta > 0$ be as in Lemma 5.1. Define

$$\begin{aligned} I &= \{\alpha_0 \in [0, 2\pi] : |\langle v, \pi_{\alpha_0}(\mathbf{i}) - \pi_{\alpha_0}(\mathbf{j}) \rangle| \geq \delta\}, \\ J &= \{\alpha_0 \in [0, 2\pi] : |\partial_{\alpha=\alpha_0} \langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle| \geq \delta\}. \end{aligned}$$

Lemma 5.1 guarantees that $I \cup J = [0, 2\pi]$. We trivially have

$$\mathcal{L}^1(H_t \cap I) \leq \begin{cases} 2\pi \leq 2\pi \delta^{-1} t, & \text{if } t \geq \delta, \\ 0, & \text{if } t < \delta. \end{cases}$$

It remains to estimate the integral in the complement of I . Since $[0, 2\pi] \setminus I \subset J$ is open, it can be written in a unique way as a union of disjoint open intervals I_j in which the function $\alpha \mapsto \langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle$ is strictly monotone. If β_1 and β_2 are the end points of I_j , then $\langle v, \pi_{\beta_1}(\mathbf{i}) - \pi_{\beta_1}(\mathbf{j}) \rangle = \delta$ and $\langle v, \pi_{\beta_2}(\mathbf{i}) - \pi_{\beta_2}(\mathbf{j}) \rangle = -\delta$ or vice versa. Since

$$\begin{aligned} \partial_{\alpha=\alpha_0} \langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle &= \langle v, \pi_{\alpha_0 + \pi/2}(\mathbf{i}) - \pi_{\alpha_0 + \pi/2}(\mathbf{j}) \rangle \\ &\leq |\pi_{\alpha_0 + \pi/2}(\mathbf{i}) - \pi_{\alpha_0 + \pi/2}(\mathbf{j})| \leq \frac{2\|\mathbf{v}\|}{1 - \|\mathbf{A}\|} \end{aligned}$$

for all $\alpha_0 \in [0, 2\pi]$, we have

$$\begin{aligned} 2\delta &= |\langle v, \pi_{\beta_2}(\mathbf{i}) - \pi_{\beta_2}(\mathbf{j}) \rangle - \langle v, \pi_{\beta_1}(\mathbf{i}) - \pi_{\beta_1}(\mathbf{j}) \rangle| \\ &= \left| \int_{\beta_1}^{\beta_2} \partial_{\alpha=\alpha_0} \langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle d\mathcal{L}^1(\alpha) \right| \leq \frac{2\|v\|}{1 - \|\mathbf{A}\|} \mathcal{L}^1(I_j). \end{aligned}$$

Thus there are only finitely many intervals I_j . For each j , by the monotonicity and the mean value theorem, there exists $\lambda_j \in I_j$ such that

$$|\langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle| = |\langle v, \pi_{\alpha}(\mathbf{i}) - \pi_{\alpha}(\mathbf{j}) \rangle - \langle v, \pi_{\lambda_j}(\mathbf{i}) - \pi_{\lambda_j}(\mathbf{j}) \rangle| \geq \delta|\alpha - \lambda_j|$$

for all $\alpha \in I_j$. By this estimate, we have

$$\mathcal{L}^1(H_t \cap I_j) \leq \mathcal{L}^1(B(\lambda_j, \delta^{-1}t)) \leq 2\pi\delta^{-1}t.$$

Since

$$\mathcal{L}^1(H_t) \leq \mathcal{L}^1(H_t \cap I) + \sum_j \mathcal{L}^1(H_t \cap I_j) \leq C't,$$

we have shown (5.1) and therefore, finished the proof. \square

Proof of Theorem A. Let \mathcal{N} be the product of Haar measures on $GL_2(\mathbb{R})^N$. This means that

$$\mathcal{N}(\mathcal{B}) = \int_{\mathcal{B}} \prod_{i=1}^N \left(\frac{1}{\det A_i} \right)^2 d\mathcal{L}^{4N}(\mathbf{A}).$$

for all measurable $\mathcal{B} \subset GL_d(\mathbb{R})^N$. Since $\mathcal{L}^{4N} \ll \mathcal{N}$ it suffices to check that the assumptions of Theorems 3.8 and 4.2 hold for the measure \mathcal{N} . By the Haar property, for any measurable $L^1(\mathcal{N})$ -function $f: GL_2(\mathbb{R})^N \rightarrow \mathbb{R}$ and for every $\alpha \in [0, 2\pi]$, we have

$$\begin{aligned} \int f(\mathbf{A}) d\mathcal{N}(\mathbf{A}) &= \int \cdots \int f(A_1, \dots, A_N) \prod_{i=1}^N \left(\frac{1}{\det A_i} \right)^2 d\mathcal{L}^4(A_1) \cdots d\mathcal{L}^4(A_N) \\ &= \int \cdots \int f(u_{\alpha}^T A_1 u_{\alpha}, \dots, u_{\alpha}^T A_N u_{\alpha}) \prod_{i=1}^N \left(\frac{1}{\det A_i} \right)^2 d\mathcal{L}^4(A_1) \cdots d\mathcal{L}^4(A_N). \end{aligned}$$

Thus, $\mathcal{N} = \int H_{\mathbf{A}} \mathcal{L}^1 d\mathcal{N}(\mathbf{A})$, where $H_{\mathbf{A}}(\alpha) = (u_{\alpha}^T A_1 u_{\alpha}, \dots, u_{\alpha}^T A_N u_{\alpha})$. Therefore, by Proposition 5.2 and Lemma 3.4, the statement for Bernoulli measures in Theorem A follow from Theorem 3.8.

If $\chi_1(\mathbf{A}, \mu) = \chi_2(\mathbf{A}, \mu)$, where μ is an ergodic equilibrium state of \mathbf{A} , then by Lemma 2.4 and [9, Theorem 2.13 and Corollary 4.16] we have $\dim_{\text{H}} E_{\mathbf{A}, \mathbf{v}} = \dim_{\text{aff}} \mathbf{A}$. Thus, without loss of generality, we may restrict \mathcal{N} to the set $\mathbb{A} = \{\mathbf{A} : \chi_1(\mathbf{A}, \mu) > \chi_2(\mathbf{A}, \mu)\}$. Hence, the statement for self-affine sets in Theorem A follows from Lemma 2.4, Proposition 5.2, and Theorem 4.2 with the measure $\mathcal{M} = \mathcal{N}|_{\mathbb{A}}$. \square

6. THE HIGHER DIMENSIONAL CASE

In this section, we prove Theorem B as a consequence of Theorem 3.8. At first, we show that the modified self-affine transversality condition holds for some pair (\mathcal{U}, m) for all $\mathbf{A} \in \mathcal{A}'_{\mathbf{v}}$. Secondly, we show that Lebesgue typical matrix tuples have simple Lyapunov spectra for all Bernoulli measures, and observe that $\mathcal{D} \subset \mathbb{A}$, where \mathcal{D} is defined in (2.6) and \mathbb{A} is as in §4.

Recall that if $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^d)^N$ is such that $v_i \neq v_j$ for every $i \neq j$, then

$$\mathcal{A}'_{\mathbf{v}} = \left\{ \mathbf{A} \in GL_d(\mathbb{R})^N : 0 < \max_{i \neq j} \frac{\|A_i\| + \|A_j\|}{|v_i - v_j|} \cdot \frac{\|\mathbf{v}\|}{1 - \|\mathbf{A}\|} < \frac{2}{\sqrt{3}} - 1 \right\}.$$

Recall also that

$$\begin{aligned} O(d) &= \{G \in GL_d(\mathbb{R}) : G^T G = G G^T = I\}, \\ SO(d) &= \{G \in O(d) : \det(G) = 1\} \end{aligned}$$

are the *orthogonal group* and the *special orthogonal group*, respectively. We define \mathcal{U} to be $O(d)$ and choose m to be the Haar measure Θ on $O(d)$. The following proposition shows that $(O(d), \Theta)$ satisfies the modified self-affine transversality condition for all $\mathbf{A} \in \mathcal{A}'_{\mathbf{v}}$.

Proposition 6.1. *Let $d \in \mathbb{N}$ be such that $d \geq 3$, $\mathbf{v} = (v_1, \dots, v_N) \in (\mathbb{R}^2)^N$ be such that $v_i \neq v_j$ for $i \neq j$, and $\mathbf{A} = (A_1, \dots, A_N) \in \mathcal{A}'_{\mathbf{v}}$. Then there is a constant $C > 0$ such that*

$$\Theta(\{G \in O(d) : |\text{proj}_V(\pi_{\mathbf{A}, G(\mathbf{v})}(\mathbf{i}) - \pi_{\mathbf{A}, G(\mathbf{v})}(\mathbf{j}))| < t\}) \leq C \prod_{i=1}^{\dim V} \min\left\{1, \frac{t}{\alpha_i(\text{proj}_V A_{\mathbf{i} \wedge \mathbf{j}})}\right\}.$$

for all $\mathbf{i}, \mathbf{j} \in \Sigma$ with $\mathbf{i} \neq \mathbf{j}$, $t > 0$, and proper subspaces V of \mathbb{R}^d .

Proof. Fix $\mathbf{A} \in \mathcal{A}'_{\mathbf{v}}$ and $t > 0$, and let $V \subset \mathbb{R}^d$ be a proper subspace. Fix $n \in \mathbb{N}$ and let $\mathbf{i} = (i_1, i_2, \dots), \mathbf{j} = (j_1, j_2, \dots) \in \Sigma$ be such that $\mathbf{i} \neq \mathbf{j}$ and $\mathbf{i} \wedge \mathbf{j} \in \Sigma_{n-1}$. Define

$$E_{\mathbf{i}, \mathbf{j}}(G) = v_{i_n} - v_{j_n} + \sum_{k=n+1}^{\infty} G^T A_{\sigma^n(\mathbf{i}|_k)} G v_{i_{k+1}} - \sum_{k=n+1}^{\infty} G^T A_{\sigma^n(\mathbf{j}|_k)} G v_{j_{k+1}}$$

and write $w_{\mathbf{i}, \mathbf{j}}(G) = E_{\mathbf{i}, \mathbf{j}}(G)/|E_{\mathbf{i}, \mathbf{j}}(G)|$ for all $G \in O(d)$. Let $D_{\mathbf{i} \wedge \mathbf{j}} = A_{\mathbf{i} \wedge \mathbf{j}}^T \text{proj}_V^T \text{proj}_V A_{\mathbf{i} \wedge \mathbf{j}}$. The claim is equivalent to

$$\Theta(\{G \in O(d) : |\langle G E_{\mathbf{i}, \mathbf{j}}(G), D_{\mathbf{i} \wedge \mathbf{j}} G E_{\mathbf{i}, \mathbf{j}}(G) \rangle| < t^2\}) \leq C \prod_{i=1}^{\dim V} \min\left\{1, \frac{t}{\alpha_i(\text{proj}_V A_{\mathbf{i} \wedge \mathbf{j}})}\right\} \quad (6.1)$$

and therefore, our task is to show that (6.1) holds.

Defining

$$R = \frac{\|\mathbf{v}\|}{1 - \|\mathbf{A}\|} \quad \text{and} \quad r = |v_{i_n} - v_{j_n}| - (\|A_{i_n}\| + \|A_{j_n}\|)R$$

we have

$$0 < r \leq |E_{\mathbf{i}, \mathbf{j}}(G)| \leq |v_{i_n} - v_{j_n}| + (\|A_{i_n}\| + \|A_{j_n}\|)R \leq 2R. \quad (6.2)$$

Observe that we also have

$$\begin{aligned} |w_{\mathbf{i}, \mathbf{j}}(G) - w_{\mathbf{i}, \mathbf{j}}(H)| &\leq \left| E_{\mathbf{i}, \mathbf{j}}(G) \left(\frac{1}{|E_{\mathbf{i}, \mathbf{j}}(G)|} - \frac{1}{|E_{\mathbf{i}, \mathbf{j}}(H)|} \right) + \frac{E_{\mathbf{i}, \mathbf{j}}(G) - E_{\mathbf{i}, \mathbf{j}}(H)}{|E_{\mathbf{i}, \mathbf{j}}(H)|} \right| \\ &\leq r^{-1} 2 |E_{\mathbf{i}, \mathbf{j}}(G) - E_{\mathbf{i}, \mathbf{j}}(H)| \\ &\leq 4r^{-1} R (\|A_{i_n}\| + \|A_{j_n}\|) \|G - H\| \end{aligned} \quad (6.3)$$

for all $G, H \in O(d)$. Furthermore, write

$$c_1 = 4r^{-1} R (\|A_{i_n}\| + \|A_{j_n}\|)$$

and note that, by the definition of $\mathcal{A}'_{\mathbf{v}}$, we have $c_1 < 1$. Recalling that $O(d)$ is compact, let $\{B(G_k, 2^{-1}t^2r^{-2}c_1^{-1})\}_{k=1}^K$ be a finite maximal packing of $O(d)$ and let $B_k = B(G_k, t^2r^{-2}c_1^{-1})$. Notice that $\{B_k\}_{k=1}^K$ is a cover of $O(d)$. Fix $e \in S^{d-1}$ such that $\langle e, v_{i_n} - v_{j_n} \rangle = 0$ and write $w_k = w_{\mathbf{i}, \mathbf{j}}(G_k)$. Let $H_k \in SO(d)$ be such that it is a rotation in the plane spanned by w_k and e and satisfies

$$w_k = H_k e.$$

Claim 1. *It holds that*

$$\begin{aligned} \Theta(\{G \in O(d) : |\langle G E_{\mathbf{i}, \mathbf{j}}(G), D_{\mathbf{i} \wedge \mathbf{j}} E_{\mathbf{i}, \mathbf{j}}(G) \rangle| < t^2\}) \\ \leq \sum_{k=1}^K \Theta(\{G \in B_k H_k : |\langle G e, D_{\mathbf{i} \wedge \mathbf{j}} G e \rangle| < 2t^2/r^2\}). \end{aligned}$$

Proof. Note that without loss of generality, we may assume that $\|D_{i \wedge j}\| \leq 1$. Therefore,

$$|\langle Gx, D_{i \wedge j}Gx \rangle - \langle Gy, D_{i \wedge j}Gy \rangle| \leq 2|x - y|. \quad (6.4)$$

for every $x, y \in S^{d-1}$. Observe that if $G \in B_k$, then (6.3) implies $|w_{i,j}(G) - w_k| < t^2/r^2$. Thus, by (6.4) and the invariance of Θ under the action of $SO(d)$, we obtain

$$\begin{aligned} & \Theta(\{G \in O(d) : |\langle GE_{i,j}(G), D_{i \wedge j}GE_{i,j}(G) \rangle| < t^2\}) \\ & \leq \Theta(\{G \in O(d) : |\langle Gw_{i,j}(G), D_{i \wedge j}Gw_{i,j}(G) \rangle| < t^2/r^2\}) \\ & \leq \sum_{k=1}^K \Theta(\{G \in B_k : |\langle Gw_{i,j}(G), D_{i \wedge j}Gw_{i,j}(G) \rangle| < t^2/r^2\}) \\ & \leq \sum_{k=1}^K \Theta(\{G \in B_k : |\langle Gw_k, D_{i \wedge j}Gw_k \rangle| < 2t^2/r^2\}) \\ & \leq \sum_{k=1}^K \Theta(\{G \in B_k : |\langle GH_k e, D_{i \wedge j}GH_k e \rangle| < 2t^2/r^2\}) \\ & = \sum_{k=1}^K \Theta(\{G \in B_k H_k : |\langle Ge, D_{i \wedge j}Ge \rangle| < 2t^2/r^2\}) \end{aligned} \quad (6.5)$$

which is what we wanted. \square

Claim 2. *There exists a constant $0 < \varrho < 1$ such that*

$$\|H_k - H_h\| \leq \varrho \|G_k - G_h\|.$$

Proof. If $w_k = w_h$, then also $H_h = H_k$ and there is nothing to prove. Furthermore, if $w_k = e$, then

$$\|H_k - H_h\| = \|\text{Id} - H_h\| = 2|\sin(\beta/2)|,$$

where β is the angle by which H_h rotates. Since $H_h(e) = w_h$ and $2\sin(\beta/2) = |w_k - w_h|$, the claim follows from (6.3). Since the case $w_h = e$ is similar, we may assume that w_k, w_h , and e are all distinct. Define

$$W = \text{span}\{w_k, w_h, e\}.$$

We have $H_k u = H_h u = u$ for all $u \in W^\perp$ since H_k and H_h are rotations on W . Observe that

$$\begin{aligned} \|H_k - H_h\| &= \|H_k^T - H_h^T\| = \sup\{|(H_k^T - H_h^T)w| : w \in W \text{ such that } |w| = 1\} \\ &= \sup\{|(2 - 2\langle H_k^T w, H_h^T w \rangle)^{1/2}| : w \in W \text{ such that } |w| = 1\} \\ &= (2 - 2\inf\{\langle w, H_k H_h^T w \rangle : w \in W \text{ such that } |w| = 1\})^{1/2} \\ &= (2 - 2\cos\beta)^{1/2} = 2|\sin(\beta/2)|, \end{aligned} \quad (6.6)$$

where β is the angle by which $H_k H_h^T$ rotates. Observe that, by (6.2),

$$\left| \left\langle \frac{v_{i_n} - v_{j_n}}{|v_{i_n} - v_{j_n}|}, w_{i,j}(G) \right\rangle \right| \geq \frac{|v_{i_n} - v_{j_n}| - (\|A_{i_n}\| + \|A_{j_n}\|)R}{|E_{i,j}(G)|} \geq \frac{r}{|v_{i_n} - v_{j_n}| + (\|A_{i_n}\| + \|A_{j_n}\|)R}.$$

Therefore, for every $u \in S^{d-1}$ with $u \perp v_{i_n} - v_{j_n}$ we have

$$|u - w_{i,j}(G)| \geq \frac{r}{|v_{i_n} - v_{j_n}| + (\|A_{i_n}\| + \|A_{j_n}\|)R}. \quad (6.7)$$

Notice that

$$H_k^T w_k = H_h^T w_h = e, \quad H_k H_h^T w_h = w_k,$$

and further,

$$H_k^T e = 2\langle e, w_k \rangle - w_k, \quad \text{and} \quad H_h^T e = 2\langle e, w_h \rangle - w_h, \quad \text{so that} \quad \langle e, H_k H_h^T e \rangle = \langle w_k, w_h \rangle.$$




FIGURE 3. Figure (a) displays the possible relative positions of the points w_h, w_k and e . The number β is the angle by which $H_k H_h^T$ rotates. Figure (b) represents the triangle with vertices w_h, w_k and f in a plane invariant under $H_k H_h^T$.

We see that, under the rotation $H_k H_h^T$, all of the vectors e, w_k, w_h are rotated by the same angle. Furthermore, w_k and w_h are mapped to each other. This is only possible if all the points are within the same distance from the equator of the unit sphere of $\text{span}\{w_k, w_h, e\}$; see Figure 3(a). Therefore the plane of rotation for $H_k H_h^T$ is either $\text{span}\{w_k - e, w_h - e\}$ or $\text{span}\{w_k + e, w_h + e\}$, depending on whether e is on the same hemisphere of the unit sphere with w_k and w_h or not. Let $f \in \{e, -e\}$ be such that $\text{span}\{w_k - f, w_h - f\}$ is the plane of rotation.

In the plane of rotation, consider the triangle having the points w_k, w_h , and f as its vertices, and is of height m (from the vertex w_k); see Figure 3(b). The area \mathfrak{A} of this triangle satisfies

$$\frac{\sin(\beta/2)|w_k - f||w_h - f|}{2} = \mathfrak{A} = \frac{m|w_h - f|}{2} \leq \frac{|w_k - w_h||w_k - f|}{2}.$$

Therefore, we have

$$\sin(\beta/2) \leq \frac{|w_k - w_h|}{|w_k - f|}. \quad (6.8)$$

Thus, by (6.6), (6.7), (6.3), and (6.8), we get

$$\|H_k - H_h\| \leq \frac{4R(\|A_{i_n}\| + \|A_{j_n}\|)(|v_{i_n} - v_{j_n}| + (\|A_{i_n}\| + \|A_{j_n}\|)R)}{r^2} \|G_k - G_h\|.$$

Since, by the definition of \mathcal{A}'_v ,

$$\frac{4R(\|A_{i_n}\| + \|A_{j_n}\|)(|v_{i_n} - v_{j_n}| + (\|A_{i_n}\| + \|A_{j_n}\|)R)}{r^2} < 1$$

we have finished the proof of Claim 2. \square

We will now finish the proof of Proposition 6.1 by verifying (6.1) as a consequence of Claims 1 and 2. By Claim 2 and the fact that $\{B(G_k, 2^{-1}t^2r^{-2}c_1^{-1})\}_{k=1}^K$ is a maximal packing of $O(d)$, we have

$$\|G_k H_k - G_h H_h\| \geq \|G_k - G_h\| - \|H_k - H_h\| \geq (1 - \varrho)2^{-1}t^2r^{-2}c_1^{-1}$$

for $k \neq h$. Since the metric $d(G, H) = \|G - H\|$ is invariant with respect to the group action, the Haar measure Θ is equivalent to the $d(d-1)/2$ -dimensional Hausdorff measure on $O(d)$. Thus,

there exists a constant C independent of t such that any $H \in O(d)$ is contained in at most C balls from the family $\{B(G_k H_k, t^2 r^{-2} c_1^{-1})\}_{k=1}^K$. Hence, by Claim 1,

$$\begin{aligned} & \Theta(\{G \in O(d) : |\langle G E_{\mathbf{i}\mathbf{j}}(G), D_{\mathbf{i}\wedge\mathbf{j}} E_{\mathbf{i}\mathbf{j}}(G) \rangle| < t^2\}) \\ & \leq \sum_{k=1}^K \Theta(\{G \in B_k H_k : |\langle G e, D_{\mathbf{i}\wedge\mathbf{j}} G e \rangle| < 2t^2/r^2\}) \\ & \leq C \Theta(\{G \in O(d) : |\langle G e, D_{\mathbf{i}\wedge\mathbf{j}} G e \rangle| < 2t^2/r^2\}) \\ & = C \sigma^{d-1}(\{x \in S^{d-1} : |\langle x, D_{\mathbf{i}\wedge\mathbf{j}} x \rangle| < 2t^2/r^2\}), \end{aligned}$$

where σ^{d-1} is the spherical measure on the unit sphere S^{d-1} . Since $\dim V \leq d-1$, we have $\alpha_d(\text{proj}_V A_{\mathbf{i}\wedge\mathbf{j}}) = 0$, and

$$\begin{aligned} & C \sigma^{d-1}(\{x \in S^{d-1} : |\langle x, D_{\mathbf{i}\wedge\mathbf{j}} x \rangle| < 2t^2/r^2\}) \\ & \leq C \mathcal{L}^{d-1}(\{(x_1, \dots, x_{d-1}) \in B(0, 1) : \sum_{l=1}^{d-1} \alpha_l(\text{proj}_V A_{\mathbf{i}\wedge\mathbf{j}})^2 x_l^2 < 2t^2/r^2\}) \\ & \leq C \min\{1, \sqrt{2}/r\}^{d-1} \prod_{l=1}^{\dim V} \min\left\{1, \frac{t}{\alpha_l(\text{proj}_V A_{\mathbf{i}\wedge\mathbf{j}})}\right\}. \end{aligned}$$

This proves (6.1) and finishes the proof. \square

Our next lemma shows that almost every matrix tuples have simple Lyapunov spectra with respect to any Bernoulli measure. The proof of the lemma is similar to that of [27, Theorem 7.12]. We say that $\mathbf{A} = (A_1, \dots, A_N) \in GL_d(\mathbb{R})^N$ is *pinching* if there exists $\mathbf{i} \in \Sigma_*$ such that $A_{\mathbf{i}}$ has only real eigenvalues with distinct absolute value. Furthermore, we say that \mathbf{A} is *twisting* if there exists $\mathbf{j} \in \Sigma_*$ such that $A_{\mathbf{j}} E \cap F = \{0\}$ for all invariant subspaces E and F of $A_{\mathbf{j}}$ with dimension $\dim E + \dim F \leq d$. Assuming \mathbf{A} to be pinching and twisting guarantees the Lyapunov exponents to be distinct; see [1, Theorem A]. In the study of self-affine sets, this is often a useful property; for example, see [18, Theorem 3.1 and Remark 3.2].

Lemma 6.2. *Let \mathcal{M} be the product of Haar measures on $GL_d(\mathbb{R})^N$, i.e.*

$$\mathcal{M}(\mathcal{B}) = \int_{\mathcal{B}} \prod_{i=1}^N \left(\frac{1}{\det A_i} \right)^d d\mathcal{L}^{d^2 N}(\mathbf{A})$$

for all measurable $\mathcal{B} \subset GL_d(\mathbb{R})^N$. Then

$$\mathcal{M}(\{\mathbf{A} \in GL_d(\mathbb{R})^N : \text{there exist } \mathbf{p} \in (0, 1)^N \text{ and } i \neq j \text{ such that } \chi_i(\mathbf{A}, \mathbf{p}) = \chi_j(\mathbf{A}, \mathbf{p})\}) = 0.$$

Proof. By [1, Theorem A], it is sufficient to show that the set of \mathbf{A} without pinching and twisting condition is contained in countably many $d^2 N - 1$ dimensional manifolds. It is easy to see that without loss of generality, we may assume that $N = 2$.

We start by sketching the argument in the case $d = 2$. This follows from [1, Theorem A], where the assumptions of the theorem are satisfied by the following argument: After perturbing one of the maps by a small rotation, we see that the twisting condition holds. The pinching condition, in the case where A_1 or A_2 have only real eigenvalues, follows immediately by slightly perturbing the matrix having real eigenvalues. Suppose then that both A_1 and A_2 have complex eigenvalues. Then we may consider A_1 to be the composition of a rotation with a dilation. Up to perturbation, we may suppose that the rotation is irrational. Up to another perturbation, we may suppose that A_2 is not conformal: there is a cone C whose image $A_2(C)$ is strictly slimmer. Then we may find arbitrarily large values of n such that $B = A_1^n A_2(C)$ is contained in C . This implies that the eigenvalues of B are real and different in absolute value and hence, the pinching condition holds.

Henceforth we assume that $d \geq 3$. Curiously, this assumption is needed in our argument. Let $\lambda_1(A), \dots, \lambda_d(A)$ be the eigenvalues of a matrix A written in a decreasing order by the absolute

value, i.e. $|\lambda_i(A)| \geq |\lambda_{i+1}(A)|$ for all i . Let $A_1, A_2 \in GL_d(\mathbb{R})$. If $\lambda_i(A_j) \in \mathbb{R}$, then $|\lambda_i(A_j)| \neq |\lambda_k(A_j)|$ for $k \neq i$. If $\lambda_i(A_j) = \overline{\lambda_{i+1}(A_j)} \in \mathbb{C}$, then $|\lambda_i(A_j)| \neq |\lambda_k(A_j)|$ for $k \notin \{i, i+1\}$. Observe that in this case, by an arbitrary small perturbation on the matrices A_1 and A_2 , we may assume that the argument of the eigenvalue $\lambda_i(A_j)$ is irrational. Similarly, we may also assume that the eigenvalues are rationally independent, i.e. none of the eigenvalues is not rational multiple of each other.

Let us assume that A_j has r_j real and c_j complex eigenvalues. Then $r_j + 2s_j = d$ and, moreover, all eigenspaces of A_j have dimension either 1 or 2. Let us denote the eigenspaces of A_j by $\xi_{e(1)}(A_j), \dots, \xi_{e(d)}(A_j)$ corresponding to the eigenvalues $\lambda_1(A_j), \dots, \lambda_d(A_j)$. This means that if $\lambda_i(A_j) \in \mathbb{R}$, then $\dim \xi_{e(i)}(A_2) = 1$ and $\xi_{e(i)}(A_2)$ is the eigenspace of $\lambda_i(A_j) \in \mathbb{R}$. On the other hand, if $\lambda_i(A_j) = \overline{\lambda_{i+1}(A_j)} \in \mathbb{C}$, then $\xi_{e(i)}(A_2) = \xi_{e(i+1)}(A_2)$, $\dim \xi_{e(i)}(A_2) = 2$, and $\xi_{e(i)}(A_2)$ is the eigenspace of $\lambda_i(A_j)$. By an arbitrary small perturbation, we may assume that the eigenspaces are in general position: for every $i \in \{1, \dots, d\}$ there exist non-zero subspaces $E_1, E_2 \subset \xi_{e(i)}(A_1)$ and $F_1, F_2 \subset \xi_{e(i+1)}(A_2)$ such that

$$\begin{aligned} \text{span}\{\xi_{e(1)}(A_1), \dots, E_1\} \cap \text{span}\{F_1, \dots, \xi_{e(d)}(A_2)\} &= \{0\}, \\ \text{span}\{\xi_{e(1)}(A_2), \dots, E_2\} \cap \text{span}\{F_2, \dots, \xi_{e(d)}(A_1)\} &= \{0\}. \end{aligned} \quad (6.9)$$

We define a flag of cones $C_1 \times C_2 \times \dots \times C_{d-1}$ in $G(1, d) \times G(2, d) \times \dots \times G(d-1, d)$ by taking sufficiently small neighbourhoods of the flag of invariant subspaces $\text{span}\{\xi_{e(1)}(A_1), \dots, \xi_{e(i)}(A_1)\}$ and sufficiently small neighbourhoods of an i -dimensional subspace of $\text{span}\{\xi_{e(1)}, \dots, \xi_{e(i)}(A_1)\}$, which contains $\text{span}\{\xi_{e(1)}, \dots, \xi_{e(i-1)}(A_1)\}$ if $\lambda_i(A_1) = \overline{\lambda_{i+1}(A_1)} \in \mathbb{C}$ such that it satisfies the following six conditions:

- (1) every C_i is open and simply connected in $G(i, d)$,
- (2) for every $E \in C_i$ there exists $F \in C_{i+1}$ such that $E \subset F$,
- (3) if $\lambda_i(A_1) = \overline{\lambda_{i+1}(A_1)} \in \mathbb{C}$, then C_i contains $\text{span}\{\xi_{e(1)}(A_1), \dots, \xi_{e(i-1)}(A_1), E\}$, where E is as in (6.9), and $\overline{C_i}$ is transversal to $\text{span}\{E', \xi_{e(i+2)}(A_1), \dots, \xi_{e(d)}(A_1)\}$ for a subspace E' with $\text{span}\{E, E'\} = \xi_{e(i)}(A_1)$,
- (4) if $\lambda_i(A_1) \in \mathbb{R}$, then C_i contains $\text{span}\{\xi_{e(1)}(A_1), \dots, \xi_{e(i)}(A_1)\}$, and $\overline{C_i}$ is transversal to $\text{span}\{\xi_{e(i+1)}(A_1), \dots, \xi_{e(d)}(A_1)\}$,
- (5) if $\lambda_i(A_2) \in \mathbb{R}$, then $\text{span}\{\xi_{e(1)}(A_2), \dots, \xi_{e(i)}(A_2)\}$ is not contained in $\overline{C_i}$ and $\overline{C_i}$ is transversal to $\text{span}\{\xi_{e(i+1)}(A_2), \dots, \xi_{e(d)}(A_2)\}$,
- (6) if $\lambda_i(A_2) = \overline{\lambda_{i+1}(A_2)} \in \mathbb{C}$, then $\overline{C_i}$ is transversal to $\text{span}\{F, \xi_{e(i+2)}(A_2), \dots, \xi_{e(d)}(A_2)\}$ and $\text{span}\{\xi_{e(1)}(A_2), \dots, F\} \notin \overline{C_i}$ for all proper subspaces $F \subset \xi_{e(i)}(A_2)$, $\overline{C_{i+1}}$ is transversal to $\text{span}\{\xi_{e(i+2)}(A_2), \dots, \xi_{e(d)}(A_2)\}$, and $\text{span}\{\xi_{e(1)}(A_2), \dots, \xi_{e(i+1)}(A_2)\} \notin \overline{C_i}$.

Here two collections of subspaces are transversal if they form a positive angle.

Observe that, by the properties of the eigenvalues, for both $j \in \{1, 2\}$, if $\lambda_i(A_j) \in \mathbb{R}$, then

$$A_j^k(C_i) \rightarrow \text{span}\{\xi_1(A_j), \dots, \xi_{e(i)}(A_j)\}$$

uniformly. Moreover, if $\lambda_i(A_j) = \overline{\lambda_{i+1}(A_j)} \in \mathbb{C}$, then

$$A_j^k(C_{i+1}) \rightarrow \text{span}\{\xi_1(A_j), \dots, \xi_{e(i+1)}(A_j)\} \quad \text{and} \quad d(A_j^k(C_i), \text{span}\{\xi_1(A_j), \dots, \xi_{e(i)}(A_j)\}) \rightarrow 0$$

uniformly. Note that in this case, $\text{diam}(A_j^k(C_i))$ does not tend to zero. Thus, if i is an index such that either $\lambda_i(A_1) \in \mathbb{R}$, or $\lambda_i(A_2) \in \mathbb{R}$, or $\lambda_{i-1}(A_1) = \overline{\lambda_i(A_1)} \in \mathbb{C}$ and $\lambda_{i-1}(A_2) = \overline{\lambda_i(A_2)} \in \mathbb{C}$, then there exist $n_1, n_2 \in \mathbb{N}$ such that

$$A_1^{n_2} A_2^{n_1}(\overline{C_i}) \subset C_i. \quad (6.10)$$

Let i_1, \dots, i_p be the indices for which $\lambda_{i_j}(A_1) = \overline{\lambda_{i_j+1}(A_1)} \in \mathbb{C}$ and $\lambda_{i_j}(A_2) = \overline{\lambda_{i_j+1}(A_2)} \in \mathbb{C}$. Let v_1^j, \dots, v_d^j be a basis of \mathbb{R}^d such that $v_i^j \in \xi_{e(i)}(A_j)$. Observe that there exists a subsequence $(m_q)_{q \in \mathbb{N}}$ such that $(\lambda_1(A_k) \cdots \lambda_{i_j-1}(A_k) |\lambda_{i_j}(A_k)|)^{-m_q} (A_k^{\wedge i_j})^{m_q} \rightarrow P_k$, where P_k is the projection onto $V_k = \text{span}\{v_1^k \wedge \cdots \wedge v_{i_j-1}^k \wedge v_{i_j}^k, v_1^k \wedge \cdots \wedge v_{i_j-1}^k \wedge v_{i_j+1}^k\}$ along the invariant subspace W_k of $A_k^{\wedge i_j}$

transversal to V_k so that $\dim V_k + \dim W_k = \binom{d}{i_j}$. Note that if $v \in \wedge^{i_j} \mathbb{R}^d$, then $\|P_k v\| \leq \|v\|$, and if $v \notin V_k$, then $\|P_k v\| < \|v\|$. Since by a sufficiently small perturbation, we may assume that the subspaces V_1, W_1 and V_2, W_2 are in general position, there exists a unique $a_k \in V_k$ with $\|a_k\| = 1$ such that $\sup_{v \in V_k} \|P_{3-k} v\| / \|v\| = \|P_{3-k} a_k\|$. Thus, there exist cones C'_k in V_k with arbitrary large diameter such that $\text{diam}(P_{3-k}(C'_k)) < c' \text{diam } C'_k$ and $a_k \in C'_k$ for some $0 < c' < 1$. Let us denote the rotation on V_k , which maps $P_k a_{3-k}$ to a_k , by O_k . Therefore, $O_k P_k O_{3-k} P_{3-k}(\overline{C'_k}) \subset C'_k$. By the irrationality of the argument of the eigenvalues, one can choose q sufficiently large and $n_1, n_2 \in \mathbb{N}$ such that $(A_1^{\wedge i_j})^{n_1} (A_1^{\wedge i_j})^{m_q} (A_2^{\wedge i_j})^{n_2} (A_2^{\wedge i_j})^{m_q} (\overline{C'_k}) \subset C'_k$. Using the natural correspondence between $G(d, i_j)$ and $\wedge^{i_j} \mathbb{R}^d$, the set C'_k corresponds to a cone C''_k in $\text{span}\{\xi_{e(1)}(A_k), \dots, \xi_{e(i_j)}(A_k)\}$ and hence, by choosing the cone C'_{i_j} in $G(d, i_j)$ sufficiently close to C''_1 , we get

$$A_1^{n_1} A_1^{m_q} A_2^{n_2} A_2^{m_q} (\overline{C'_{i_j}}) \subset C'_{i_j} \quad (6.11)$$

for all $j \in \{1, \dots, p\}$. Thus, by irrationality and rational independence of the arguments of the complex eigenvalues of A_1 and A_2 , (6.10) and (6.11) imply that there exist $n'_1, n'_2 \in \mathbb{N}$ such that

$$A_1^{n'_1} A_2^{n'_2} (\overline{C_i}) \subset C_i$$

for all $i \in \{1, \dots, d\}$. Therefore, $A_i = A_1^{n'_1} A_2^{n'_2}$ has only real eigenvalues with distinct absolute value. This shows that (A_1, A_2) is pinching. By a similar argument, there exist $m'_1, m'_2 \in \mathbb{N}$ such that $A_j = A_2^{m'_2} A_1^{m'_1}$ has only real eigenvalues with distinct absolute value and $E \cap F = \{0\}$ for all invariant subspaces E of A_i and F of A_j with $\dim E + \dim F \leq d$. This proves the twisting property. \square

Recall that \mathbb{A} is the collection of all contractive matrix tuples having distinct Lyapunov exponents with respect to an ergodic equilibrium state, see §4, and \mathcal{D} is defined in (2.6).

Proof of Theorem B. Let \mathcal{M} be as in Lemma 6.2. Since $\mathcal{L}^{d^2 N} \ll \mathcal{M}$ it suffices to check that the assumptions of Theorem 3.8 hold for the measure \mathcal{M} . By the Haar property, for any measurable $L^1(\mathcal{M})$ -function $f: GL_d(\mathbb{R})^N \rightarrow \mathbb{R}$ and for every $G \in O(d)$, we have

$$\begin{aligned} \int f(\mathbf{A}) d\mathcal{M}(\mathbf{A}) &= \int \cdots \int f(A_1, \dots, A_N) \prod_{i=1}^N \left(\frac{1}{\det A_i} \right)^d d\mathcal{L}^{d^2}(A_1) \cdots d\mathcal{L}^{d^2}(A_N) \\ &= \int \cdots \int f(G^T A_1 G, \dots, G^T A_N G) \prod_{i=1}^N \left(\frac{1}{\det A_i} \right)^d d\mathcal{L}^{d^2}(A_1) \cdots d\mathcal{L}^{d^2}(A_N). \end{aligned}$$

Thus, $\mathcal{M} = \int H_{\mathbf{A}} \Theta d\mathcal{M}(\mathbf{A})$, where $H_{\mathbf{A}}(G) = (G^T A_1 G, \dots, G^T A_N G)$.

By Lemma 6.2, for \mathcal{M} -almost every \mathbf{A} , all the Bernoulli measures have simple Lyapunov spectra and thus, by applying Lemma 3.4, any Bernoulli measure satisfies the Ledrappier-Young formula. Hence, the first statement follows by the combination of Lemma 2.4, Proposition 6.1, and Theorem 3.8.

To turn to the set dimension statement, observe that by (2.5), for every $\mathbf{A} \in \mathcal{D}$, the s -equilibrium state $\mu_{\mathbf{A}}$ has simple Lyapunov spectrum for $s = \dim_{\text{aff}} \mathbf{A}$. Moreover, by Lemma 2.3 the $\mu_{\mathbf{A}}$ is quasi-Bernoulli. Thus, by Lemma 3.4, $\mu_{\mathbf{A}}$ satisfies the Ledrappier-Young formula. Therefore, by the combination of Lemma 2.4, Proposition 6.1, Theorem 4.2, and the fact that

$$\dim_L \mu_{\mathbf{A}} = \dim_{\text{aff}} \mathbf{A},$$

the second assertion follows. \square

7. OPEN QUESTIONS

We finish the article by posing couple of questions. An affirmative answer for any of the questions would immediately improve Theorem B.

Question 7.1. Recall that \mathbb{A} is the collection of all tuples $\mathbf{A} \in GL_d(\mathbb{R})^N$ of contractive matrices that satisfy $\chi_i(\mathbf{A}, \mu) \neq \chi_j(\mathbf{A}, \mu)$ for $i \neq j$ where μ is an ergodic s -equilibrium state of \mathbf{A} and $s = \dim_{\text{aff}}(\mathbf{A})$. Since \mathcal{D} defined in (2.6) is open, the set \mathbb{A} contains interior points; see §6. Is \mathbb{A} an open and dense subset of $GL_d(\mathbb{R})^N$ with full Lebesgue measure.

Question 7.2. Is the s -equilibrium state of \mathbf{A} for $s = \dim_{\text{aff}} \mathbf{A}$ quasi-Bernoulli for \mathcal{L}^{d^2N} -almost all $\mathbf{A} \in GL_d(\mathbb{R})^N$? Recall that, by [19, Propositions 3.4 and 3.6], it is unique and satisfies certain Gibbs property in a set of full Lebesgue measure.

Question 7.3. Can every s -equilibrium state of \mathbf{A} for $s = \dim_{\text{aff}} \mathbf{A}$ be approximated by step- n Bernoulli measures? More precisely, does there, for every $\varepsilon > 0$, exist $n \in \mathbb{N}$ and a step- n Bernoulli measure μ such that $\dim_{\text{L}} \mu < \dim_{\text{aff}} \mathbf{A} - \varepsilon$? Observe that, by [22, §3.2], this cannot be done with fully supported step- n Bernoulli measures.

Acknowledgement. The authors thank Marcelo Viana for helpful discussions related to Lyapunov exponents.

REFERENCES

- [1] A. Avila and M. Viana. Simplicity of Lyapunov spectra: a sufficient criterion. *Port. Math. (N.S.)*, 64(3):311–376, 2007.
- [2] B. Bárány. On the Ledrappier-Young formula for self-affine measures. *Math. Proc. Cambridge Philos. Soc.*, 159(3):405–432, 2015.
- [3] B. Bárány and A. Käenmäki. Ledrappier-Young formula and exact dimensionality of self-affine measures. Preprint, available at arXiv:1511.05792.
- [4] B. Bárány and M. Rams. Dimension maximizing measures for self-affine systems. Preprint, available at arXiv:1507.02829.
- [5] J. Bochi and N. Gourmelon. Some characterizations of domination. *Math. Z.*, 263(1):221–231, 2009.
- [6] K. Falconer. The Hausdorff dimension of self-affine fractals. *Math. Proc. Cambridge Philos. Soc.*, 103(2):339–350, 1988.
- [7] K. Falconer. *Techniques in fractal geometry*. John Wiley & Sons, Ltd., Chichester, 1997.
- [8] K. Falconer and T. Kempton. Planar self-affine sets with equal hausdorff, box and affinity dimensions. Preprint, available at arXiv:1503.01270.
- [9] D.-J. Feng and H. Hu. Dimension theory of iterated function systems. *Comm. Pure Appl. Math.*, 62(11):1435–1500, 2009.
- [10] D.-J. Feng and P. Shmerkin. Non-conformal repellers and the continuity of pressure for matrix cocycles. *Geom. Funct. Anal.*, 24(4):1101–1128, 2014.
- [11] A. Ferguson, J. M. Fraser, and T. Sahlsten. Scaling scenery of $(\times m, \times n)$ invariant measures. *Adv. Math.*, 268:564–602, 2015.
- [12] J. M. Fraser and P. Shmerkin. On the dimensions of a family of overlapping self-affine carpets. *Ergodic Theory and Dynamical Systems*, pages 1–19, 2014.
- [13] M. Hochman. On self-similar sets with overlaps and inverse theorems for entropy in \mathbb{R}^d . Preprint, available at arXiv:1503.09043.
- [14] M. Hochman. On self-similar sets with overlaps and inverse theorems for entropy. *Ann. of Math. (2)*, 180(2):773–822, 2014.
- [15] J. Hutchinson. Fractals and self-similarity. *Indiana Univ. Math. J.*, 30(5):713–747, 1981.
- [16] T. Jordan, M. Pollicott, and K. Simon. Hausdorff dimension for randomly perturbed self affine attractors. *Comm. Math. Phys.*, 270(2):519–544, 2007.
- [17] A. Käenmäki. On natural invariant measures on generalised iterated function systems. *Ann. Acad. Sci. Fenn. Math.*, 29(2):419–458, 2004.
- [18] A. Käenmäki, H. Koivusalo, and E. Rossi. Self-affine sets with fibred tangents. *Ergodic Theory and Dynam. Systems*, to appear, available at arXiv:1505.00958.
- [19] A. Käenmäki and B. Li. Genericity of dimension drop on self-affine sets. Preprint, available at arXiv:00496.
- [20] A. Käenmäki and I. D. Morris. Structure of equilibrium states on self-affine sets and strict monotonicity of affinity dimension. Preprint, available at arXiv:1609.07360.
- [21] A. Käenmäki, T. Ojala, and E. Rossi. Rigidity of quasisymmetric mappings on self-affine carpets. Preprint, available at arXiv:1607.02244.
- [22] I. D. Morris and P. Shmerkin. On equality of Hausdorff and affinity dimensions, via self-affine measures on positive subsystems. Preprint, available at arXiv:1602.08789.
- [23] F. Przytycki and M. Urbański. On the Hausdorff dimension of some fractal sets. *Studia Math.*, 93(2):155–186, 1989.

- [24] A. Rapaport. On self-affine measures with equal Hausdorff and Lyapunov dimensions. Preprint, available at arXiv:1511.06893.
- [25] V. A. Rohlin. On the fundamental ideas of measure theory. *Amer. Math. Soc. Translation*, 1952(71):55, 1952.
- [26] D. Simmons. Conditional measures and conditional expectation; Rokhlin's disintegration theorem. *Discrete Contin. Dyn. Syst.*, 32(7):2565–2582, 2012.
- [27] M. Viana. *Lectures on Lyapunov exponents*, volume 145 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2014.

(Balázs Bárány) MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY CV4 7AL, UNITED KINGDOM
& BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS, MTA-BME STOCHASTICS RESEARCH GROUP, P.O.
BOX 91, 1521 BUDAPEST, HUNGARY

E-mail address: balubsheep@gmail.com

(Antti Käenmäki) DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. BOX 35 (MAD), FI-40014 UNIVERSITY
OF JYVÄSKYLÄ, FINLAND

E-mail address: antti.kaenmaki@jyu.fi

(Henna Koivusalo) FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, OSKAR MORGENSTERNPLATZ 1, 1090
VIENNA, AUSTRIA

E-mail address: henna.koivusalo@univie.ac.at