

Collection of formulas for the exam
Advanced Mathematics for civil engineers

$$\sin^2 x + \cos^2 x = 1$$

$$\sin(x \pm y) = \sin x \cos y \pm \cos x \sin y$$

$$\cos(x \pm y) = \cos x \cos y \mp \sin x \sin y$$

$$\tan(x \pm y) = \frac{\tan x \pm \tan y}{1 \mp \tan x \cdot \tan y}$$

$$\sin 2x = 2 \sin x \cos x$$

$$\cos 2x = \cos^2 x - \sin^2 x$$

$$\tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$$

$$\sin^2 x = \frac{1 - \cos 2x}{2}, \quad \cos^2 x = \frac{1 + \cos 2x}{2}$$

$$\sin x + \sin y = 2 \sin \frac{x+y}{2} \cos \frac{x-y}{2}$$

$$\sin x - \sin y = 2 \cos \frac{x+y}{2} \sin \frac{x-y}{2}$$

$$\cos x + \cos y = 2 \cos \frac{x+y}{2} \cos \frac{x-y}{2}$$

$$\cos x - \cos y = -2 \sin \frac{x+y}{2} \sin \frac{x-y}{2}$$

$$\sin x \cos y = \frac{1}{2} [\sin(x+y) + \sin(x-y)]$$

$$\cos x \cos y = \frac{1}{2} [\cos(x+y) + \cos(x-y)]$$

$$\sin x \sin y = -\frac{1}{2} [\cos(x+y) - \cos(x-y)]$$

$$\sinh x = \frac{e^x - e^{-x}}{2}, \quad \cosh x = \frac{e^x + e^{-x}}{2}$$

$$\cosh^2 x - \sinh^2 x = 1$$

$$\sinh 2x = 2 \sinh x \cosh x$$

$$\cosh 2x = \cosh^2 x + \sinh^2 x$$

$$\cosh^2 x = \frac{\cosh 2x+1}{2}, \quad \sinh^2 x = \frac{\cosh 2x-1}{2}$$

Notable derivatives

$$(\sinh x)' = \cosh x$$

$$(\cosh x)' = \sinh x$$

$$(\log_a x)' = \frac{1}{x \ln a}$$

$$(x^\alpha)' = \alpha x^{\alpha-1}$$

$$(e^x)' = e^x$$

$$(a^x)' = a^x \ln(a)$$

$$(\sin x)' = \cos x$$

$$(\cos x)' = -\sin x$$

$$(\tan x)' = \frac{1}{\cos^2 x}$$

$$(\cot x)' = -\frac{1}{\sin^2 x}$$

$$(\ln x)' = \frac{1}{x}$$

$$(\arcsin x)' = \frac{1}{\sqrt{1-x^2}}$$

$$(\arctan x)' = \frac{1}{1+x^2}$$

$$(\operatorname{ar sinh} x)' = \frac{1}{\sqrt{1+x^2}}$$

$$(\operatorname{ar cosh} x)' = \frac{1}{\sqrt{x^2-1}}$$

$$(\operatorname{ar tanh} x)' = \frac{1}{1-x^2}$$

$$(\operatorname{ar coth} x)' = \frac{1}{1-x^2}$$

$$(\operatorname{arc cos} x)' = -\frac{1}{\sqrt{1-x^2}}$$

$$(\operatorname{arc cot} x)' = -\frac{1}{1+x^2}$$

Differentiation rules

$$(cu)' = cu' \quad (c \text{ constant})$$

$$(u+v)' = u'+v'$$

$$(uv)' = u'v + uv'$$

$$\left(\frac{u}{v}\right)' = \frac{u'v - uv'}{v^2}$$

$$\frac{d}{dx} f(g(x)) = \frac{df}{dg} \frac{dg}{dx}$$

Rules of Integration

$$\int cf dx = c \int f dx \quad (c \text{ constant})$$

$$\int (f+g) dx = \int f dx + \int g dx$$

$$\int f(ax+b) dx = \frac{1}{a} F(ax+b) + c, \quad \text{where } F \text{ is the primitive function of } f$$

$$\int f(g(x))g'(x) dx = F(g(x)) + c, \quad \text{where } F \text{ is the primitive function of } f$$

$$\int f^\alpha f' dx = \frac{f^{\alpha+1}}{\alpha+1} + c, \text{ ha } \alpha \neq -1$$

$$\int \frac{f'}{f} dx = \ln |f| + c$$

$$\int uv' dx = uv - \int u'v dx$$

Notable substitution of variables

$$R(e^x) \quad e^x = t$$

$$R(\sqrt{ax+b}) \quad \sqrt{ax+b} = t$$

$$R\left(\frac{\sqrt{ax+b}}{\sqrt{cx+d}}\right) \quad \frac{\sqrt{ax+b}}{\sqrt{cx+d}} = t$$

$$R(\sin x, \cos x) \quad \sin x, \cos x, \tan x, \tan \frac{x}{2} = t$$

$$R(x, \sqrt{a^2 - x^2}) \quad x = a \sin t, \quad x = a \cos t$$

$$R(x, \sqrt{a^2 + x^2}) \quad x = a \sinh t$$

$$R(x, \sqrt{x^2 - a^2}) \quad x = a \cosh t$$

Notable integrals

$$\int x^\alpha dx = \frac{x^{\alpha+1}}{\alpha+1} + c \quad (\alpha \neq -1)$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax} + c$$

$$\int a^x dx = \frac{a^x}{\ln a} + c$$

$$\int \cos x dx = \sin x + c$$

$$\int \sin x dx = -\cos x + c$$

$$\int \frac{1}{\cos^2 x} dx = \tan x + c$$

$$\int \frac{1}{\sin^2 x} dx = -\cot x + c$$

$$\int \frac{1}{x} dx = \ln |x| + c$$

$$\int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a} + c$$

$$\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \operatorname{arctan} \frac{x}{a} + c$$

$$\int \frac{dx}{\sqrt{x^2 + a^2}} = \operatorname{ar sinh} \frac{x}{a} + c$$

$$\int \frac{dx}{\sqrt{x^2 - a^2}} = \operatorname{ar cosh} \frac{x}{a} + c$$

$$\int \frac{dx}{a^2 - x^2} = \frac{1}{a} \operatorname{ar tanh} \frac{x}{a} + c, \quad \text{if } \left|\frac{x}{a}\right| < 1$$

$$\int \frac{dx}{a^2 - x^2} = \frac{1}{a} \operatorname{ar coth} \frac{x}{a} + c, \quad \text{if } \left|\frac{x}{a}\right| > 1$$

$$\int \tan x dx = -\ln |\cos x| + c$$

$$\int \cot x dx = \ln |\sin x| + c$$

1. Linear algebra

1. Gram-Schmidt orthogonalization: Let $\{\underline{w}_1, \dots, \underline{w}_k\}$ be a basis of the subspace $W \subset \mathbb{R}^d$. Then $\{\underline{v}_1, \dots, \underline{v}_k\}$ forms an orthonormal basis of W , where

$$\underline{v}_1 = \frac{\underline{w}_1}{\|\underline{w}_1\|} \text{ and for } i = 2, \dots, k \quad \underline{v}_i = \frac{\underline{w}_i - \sum_{j=1}^{i-1} (\underline{v}_j \cdot \underline{w}_i) \underline{v}_j}{\|\underline{w}_i - \sum_{j=1}^{i-1} (\underline{v}_j \cdot \underline{w}_i) \underline{v}_j\|}.$$

2. Partial Differential equations

1. The sine-Fourier series of a function $f: [0, L] \mapsto \mathbb{R}$ is:

$$\sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi}{L} \cdot x\right), \text{ where } b_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi}{L} \cdot x\right) dx.$$

$$2. \int x \sin(ax) dx = \frac{\sin(ax) - ax \cos(ax)}{a^2} + c \text{ and } \int x^2 \sin(ax) dx = \frac{2 \cos(ax) + 2ax \sin(ax) - a^2 x^2 \cos(ax)}{a^3} + c$$

3. Bernoulli's solution for the vibrating string problem:

$$\begin{cases} u''_{tt} = c^2 u''_{xx} & 0 < x < L \text{ and } 0 < t \\ u(0, t) = u(L, t) \equiv 0 & 0 < t \\ u(x, 0) = f(x) & 0 < x < L \\ u'_t(x, 0) = g(x) & 0 < x < L \end{cases}$$

then $u(x, t) = \sum_{k=1}^{\infty} \sin\left(\frac{k\pi}{L}x\right) \cdot (A_k \cos\left(\frac{k\pi c}{L}t\right) + B_k \sin\left(\frac{k\pi c}{L}t\right))$, where A_k are the coefficients of the Fourier-sine series of $f(x)$ and $\frac{k\pi c}{L}B_k$ are the coefficients of the Fourier-sine series of $g(x)$.

4. Heat equation for finite rod:

$$\begin{cases} u'_t = \alpha u''_{xx} & 0 < x < L \text{ and } 0 < t \\ u(0, t) = u(L, t) \equiv 0 & 0 < t \\ u(x, 0) = f(x) & 0 < x < L \end{cases}$$

then $u(x, t) = \sum_{k=1}^{\infty} A_k e^{-\left(\frac{k\pi}{L}\right)^2 \alpha t} \sin\left(\frac{k\pi}{L}x\right)$, where A_k are the coefficients of the Fourier-sine series of $f(x)$.

3. Vectoranalysis

1. Let \mathcal{A} be an orientable surface with parametrization $\mathbf{r}(u, v)$, where $(u, v) \in T$ for some domain T and $\vec{F}: \mathbb{R}^3 \mapsto \mathbb{R}^3$ be a vectorfield. Then

$$\iint_{\mathcal{A}} \vec{F} d\vec{A} = \pm \iint_T \vec{F}(\mathbf{r}(u, v)) \cdot (\mathbf{r}'_u \times \mathbf{r}'_v) du dv,$$

where we choose $+$ if the orientation of \mathcal{A} corresponds to $\mathbf{r}'_u \times \mathbf{r}'_v$, otherwise $-$.

2. **Gauss' Theorem:** Let $K \subset \mathbb{R}^3$ be a body with boundary ∂K oriented pointing outwards. If all the second partial derivatives of the vectorfield \vec{F} exist and continuous on K then

$$\iint_{\partial K} \vec{F} d\vec{A} = \iiint_K \operatorname{div}(\vec{F}) dx dy dz.$$

3. **Stokes' Theorem:** Let \mathcal{F} be an orientable surface and $\partial\mathcal{F}$ its boundary with coherent orientation. If all the partial derivatives of the vectorfield \vec{F} exist and continuous on \mathcal{F} then

$$\int_{\partial\mathcal{F}} \vec{F} d\mathbf{r} = \iint_{\mathcal{F}} \operatorname{curl}(\vec{F}) dA.$$

4. **Green's Theorem:** Let T be a domain on the plane such that its boundary is a γ simple closed curve. If all the partial derivatives of the vectorfield \vec{F} exist and continuous on T then

$$\int_{\gamma} \vec{F} d\mathbf{r} = \iint_T Q'_x - P'_y dx dy, \text{ where } \vec{F}(x, y) = (P(x, y), Q(x, y)).$$

5. Cylindrical substitution:

$$\begin{aligned} x &= r \cos(\varphi) \\ y &= r \sin(\varphi) \\ z &= z \end{aligned}$$

with Jacobian determinant: r .

6. Spherical substitution:

$$\begin{aligned} x &= r \sin(u) \cos(v) \\ y &= r \sin(u) \sin(v) \\ z &= r \cos(u) \end{aligned}$$

with Jacobian determinant: $r^2 \sin(u)$.

7. Polar substitution on the plane:

$$\begin{aligned} x &= r \cos(v) \\ y &= r \sin(v) \end{aligned}$$

with Jacobian determinant: r .