

Hausaufgaben 5.

Taylor-Reihen

Geben Sie die Potenzreihendarstellung um $x_0 = 0$ der folgenden Funktionen an:

1. $f(x) = \frac{7}{6-5x}$ $\left(\frac{7}{6} \sum_{n=0}^{\infty} \left(\frac{5}{6}x\right)^n, \quad |x| < \frac{6}{5}\right)$
2. $f(x) = \frac{5}{9-x^2}$ $\left(\sum_{k=0}^{\infty} \frac{5}{9^{k+1}} x^{2k}, \quad |x| < 3\right)$
3. $f(x) = \sin 2x$ $\left(\sum_{k=0}^{\infty} (-1)^k \frac{(2x)^{2k+1}}{(2k+1)!}, \quad -\infty < x < \infty\right)$
4. $f(x) = \sqrt{e^x}$ $\left(\sum_{k=0}^{\infty} \frac{x^k}{2^k k!}, \quad -\infty < x < \infty\right)$
5. $f(x) = \ln(1+x^2)$ $\left(\sum_{k=1}^{\infty} (-1)^{k+1} \frac{x^{2k}}{k}, \quad |x| < 1\right)$
6. $f(x) = \sqrt[3]{8+x}$ $\left(2 \sum_{k=0}^{\infty} \binom{\frac{1}{3}}{k} \left(\frac{x}{8}\right)^k, \quad |x| < 8\right)$
7. $f(x) = \frac{1}{\sqrt{1-x^2}}$ $\sum_{n=0}^{\infty} \binom{-\frac{1}{2}}{n} (-x^2)^n, \quad -1 < x < 1$

Fourier-Reihen

1. $f(x) = 1 - x, x \in (-\pi, \pi], f(x + 2k\pi) = f(x)$ $(f(x) \approx 1 + \sum_{k=1}^{\infty} \frac{2}{k} (-1)^k \sin kx)$
2. $f(x) = \cos x, -\frac{\pi}{2} < x \leq \frac{\pi}{2}, f(x + k\pi) = f(x)$ $(f(x) = \frac{2}{\pi} + \frac{4}{\pi} \sum_{n=1}^{\infty} (-1)^{n+1} \frac{\cos 2nx}{4n^2-1})$
3. $f(x) = \operatorname{sgn}(\cos x), -\pi < x \leq \pi, f(x + 2k\pi) = f(x),$ $(\operatorname{sgn} = \operatorname{signum} = \operatorname{Vorzeichen})$
 $(f(x) \approx \frac{4}{\pi} \sum_{n=1}^{\infty} (-1)^n \frac{\cos(2n+1)x}{2n+1})$
4. $f(x) = x, x \in (-1, 1], f(x + 2k) = f(x)$ $(f(x) \approx \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin n\pi x)$
5. $f(x) = \begin{cases} x+1, & -1 < x \leq 0 \\ -x+1, & 0 < x \leq 1 \end{cases}, f(x + 2k) = f(x)$ $(f(x) = \frac{1}{2} + \frac{4}{\pi^2} \sum_{n=1}^{\infty} \frac{\cos(2n-1)\pi x}{(2n-1)^2})$