

Correlation Decay in Certain Soft Billiards

Péter Bálint¹, Imre Péter Tóth²

¹ Alfréd Rényi Institute of the H.A.S., 1053 Reáltanoda u. 13-15, Budapest, Hungary.
E-mail: bp@renyi.hu

² Mathematical Institute, Technical University of Budapest, 1111 Egy József u. 1. Budapest, Hungary
and Research group “Stochastics” of the Hungarian Academy of Sciences, affiliated to TUB.
E-mail: mogy@math.bme.hu

Received: 10 February 2003 / Accepted: 12 May 2003
Published online: ■ ■ ■ – © Springer-Verlag 2003

Abstract: Motivated by the 2D finite horizon periodic Lorentz gas, soft planar billiard systems with axis-symmetric potentials are studied in this paper. Since Sinai’s celebrated discovery that elastic collisions of a point particle with strictly convex scatterers give rise to hyperbolic, and consequently, nice ergodic behaviour, several authors (most notably Sinai, Kubo, Knauf) have found potentials with analogous properties. These investigations concluded in the work of V. Donnay and C. Liverani who obtained general conditions for a 2-D rotationally symmetric potential to provide ergodic dynamics. Our main aim here is to understand when these potentials lead to stronger stochastic properties, in particular to exponential decay of correlations and the central limit theorem. In the main argument we work with systems in general for which the rotation function satisfies certain conditions. One of these conditions has already been used by Donnay and Liverani to obtain hyperbolicity and ergodicity. What we prove is that if, in addition, the rotation function is regular in a reasonable sense, the rate of mixing is exponential, and, consequently the central limit theorem applies. Finally, we give examples of specific potentials that fit our assumptions. This way we give a full discussion in the case of constant potentials and show potentials with any kind of power law behaviour at the origin for which the correlations decay exponentially.

1. Introduction

Consider the planar motion of a point particle in a periodic array of strictly convex scatterers. Interaction with the scatterers is in the form of elastic collisions, otherwise motion is uniform. This dynamical system, the *planar Lorentz process* is a paradigm for strongly chaotic behaviour. Among other important properties ergodicity ([Si2, SCh]) and exponential decay of correlations ([Y, Ch]) have been proven for the corresponding billiard system.

In this paper we consider the following natural modification. The scatterers are no longer hard disks, the point particle may enter them. The particle moves according to some rotation symmetric potential which vanishes identically outside the disks.

Even the issue of these softened Lorentz processes has a large amount of literature. Results point in two different directions. On the one hand, for quite general softening of the potential, the chaotic behaviour is no longer present. Stable periodic orbits and islands appear in the phase space. This is generally the case with smooth potentials, see [RT, Do2, Do1] and references therein.

However, in many cases, especially when the potential is not C^1 , the chaotic behaviour persists¹. The investigation of such soft billiards dates back to the pioneer works of Sinai ([Si1]) and Kubo et al. ([Ku] and [KM]). There are two different approaches present in the literature to this hyperbolic case. On the one hand, under conditions on the derivatives (up to the second) of the potential the Hamiltonian flow turned out to be equivalent to a geodesic flow on a negatively curved manifold. This point of view is especially suitable for potentials with Coulomb type singularities, see [Kn1] for details.

The approach we follow is to study dynamics as a hyperbolic system with singularities. [M] and, especially, [DL] – which is one of our main references – are written in the spirit of this principle. Actually, in most cases it is convenient to study the discrete time dynamics, a naturally defined Poincaré section map of the Hamiltonian flow – this is the track we are going to take.

Hyperbolicity of the system is mainly related to the properties of the so-called *rotation function* that can be calculated from the potential. Being a bit technical its definition and relevant properties are discussed in the next section. Formulation of our main theorem (Theorem 1) is likewise left to the next section as it is in terms of the rotation function. Nevertheless, it might be useful to point out that

- In case the rotation function (and the billiard configuration) satisfies some hyperbolicity condition (see Definition 2), the soft billiard system is hyperbolic and ergodic. Although somewhat otherwise stated, this fact was proved in [DL]. The condition is, essentially, necessary for ergodicity (note however Remark 1).
- In this paper we concentrate on decay of correlations. If – in addition to those needed for hyperbolicity – the rotation function satisfies further regularity conditions (see Definition 3), the rate of mixing is proved to be exponential.

In most of the paper we think of the rotation function as being fixed with the desired properties. It is only Sect. 5 when we turn to some specific potentials. Nevertheless, two technical conditions supposed to hold throughout the paper are:

- In order to be able to define a rotation function at all, we introduce $h(r) = r^2(1 - 2V(r))$ (cf. Sect. 5) and require $h'(r) > 0$ for all but finitely many r (this condition ensures the lack of trapping zones, cf. [DL]):
- The scattering occurs on rotation symmetric potentials of finite range – that is, potential for every scatterer is concentrated on a circle and depends only on the distance from the center. (Note this is the case in our references like [DL], too.)
- The horizon is finite (i.e. the maximum time between two enterings of consecutive potential disks is uniformly bounded above for any trajectory).

¹ In [DL] there is a smooth potential example with ergodic behaviour, too; however it is unstable with respect to small perturbations like varying the full energy level.

Proof of our main theorem is based on our second main reference, [Ch]. In this paper, by implementing the techniques of L. S. Young from [Y], N. Chernov showed that given any hyperbolic system with singularities for which one can show the validity of certain technical properties, correlations decay exponentially fast.

What we perform below is the proof of these technical properties for our “soft” billiard system. Even though the existence of invariant cone fields is established in [DL], the *uniformity* of hyperbolicity (Subsect. 3.2) needs detailed investigation. An even more important new difficulty that we have to overcome is the treatment of quantities connected to the second derivative of the dynamics, especially while traveling through the potential. An analysis finer than before – in this sense – of the evolution of fronts is needed. This applies especially to the self-contained proof of curvature and distortion bounds (Subsect. 3.3).

It is a key aspect of our method that arguments related to expansion and distortion can be carried out by considering motion inside and outside the potential disks separately. Actually, our choice of the outgoing phase space and the Euclidean metric (see Sect. 2) is related to this point of view and not to the tradition of [Ch]. (Using the Euclidean metric with the phase space of *incoming* particles instead of outgoing, our distortion bounds would no longer hold.) The splitting of motion into “potential” and “free” intervals is, however, slightly restrictive. Namely, certain soft billiard systems that seem ergodic and exponentially mixing are covered neither in this paper nor in [DL] (see also Sect. 6, especially Remark 1).

The paper is organized as follows. In Sect. 2 the dynamical system along with the rotation function and its properties are defined, and our main theorem is formulated. Section 3 gives a detailed geometric analysis of the system. After fixing notations and establishing some basic properties in Subsect. 3.1, Subsect. 3.2 is mainly concerned with uniform hyperbolicity (Proposition 1) and related issues. In Subsect. 3.3 important regularity properties of unstable manifolds are shown. Specifically, curvature bounds, distortion bounds and absolute continuity of the holonomy map are proven (Propositions 2, 3 and 4). As a final bit of the general proof in Sect. 4 we investigate the growth of unstable manifolds. The fact that expansion prevails the harmful effect of singularities is quantified in the growth formulas of Proposition 5. As a conclusion we refer to Theorem 2.1 from [Ch]: a hyperbolic system with singularities for which Propositions 1, 2, 3, 4 and 5 are valid enjoys exponential decay of correlations. For the reader’s convenience, we formulate the theorem of Chernov in the Appendix. Last but not least, preceding some concluding remarks, in Sect. 5 we turn to the investigation of specific potentials: as corollaries of our main theorem certain soft billiard systems are shown to exhibit exponential decay of correlations.

We note that it is not clear how sharp our results are. On the one hand, the conditions for ergodicity – which are part of our conditions – formulated by Donnay and Liverani are more or less sharp (see [Do1]). On the other hand, the conditions formulated for EDC by Chernov are sufficient, but most probably not necessary. So, although we know that Chernov’s conditions (e.g. the bounded curvature assumption and the distortion bounds) are *not* satisfied when our regularity conditions are not met, it is well possible that EDC still occurs. At some points of the paper we will point out why our regularity conditions are necessary for Chernov’s method to work.

Part of the results in this paper and a sketch of the proof can also be found in the proceedings paper [BÁTó].

2. Definition and Basic Properties of the System

The phase space. Consider finitely many disjoint circles of radius R on the unit two-dimensional flat torus T^2 . (Thinking of a periodic array of circular disks on the Euclidean plane \mathbb{R}^2 would not be very much different.) We require that the configuration has finite horizon: there is a certain constant τ_{\max} such that any straight segment longer than τ_{\max} on \mathbb{R}^2 intersects at least one of the scatterers.

Remark. As the circles are disjoint, the minimum distance between two scatterers is bigger than some positive constant τ_{\min} .

Let the Hamiltonian motion of our point particle be described by a potential which is identically zero outside and is some rotation symmetric function $V(r)$ inside the circular scatterers (here r is the distance from the center of the scatterer). For simplicity we fix the mass and the full energy of our point particle as

$$m = 1, \quad E = \frac{1}{2}.$$

This way the free flight velocity has unit length, $|v| = 1$ (in other words $v \in \mathbf{S}^1$, where \mathbf{S}^1 is the unit circle in \mathbb{R}^2).

We assume (cf. Definition 2 and the remarks following it) that the Hamiltonian flow restricted to this surface of constant full energy is ergodic (with respect to Liouville measure). Equivalently one can say that the map corresponding to the naturally defined Poincaré section of the flow (see below) is ergodic. Our aim is *to study the rate of mixing for this map*.

Following tradition we work with the Poincaré section of outgoing velocities (particles that have just left one of the scatterers).

Notation. Denote by M the Poincaré section of outgoing particles. Sometimes we will also use the notation $M_+ = M$ to stress that this is the outgoing phase space, to avoid confusion.

The phase points are the boundary points of the scatterers, equipped with unit velocities pointing outwards. The phase space M is a finite union of cylinders (each corresponding to one of the circular scatterers). Coordinates for the cylinders are:

Notations. s denotes the arclength parameter along the scatterer (starting from a point arbitrarily fixed), describing position of the outgoing particle.

φ denotes the collision angle, the angle that the outgoing velocity makes with the normal vector of the scatterer in the point s . Clearly $\varphi \in [-\frac{\pi}{2}, \frac{\pi}{2}]$.

The position can be equivalently described by another angle parameter $\Theta \in [0, 2\pi]$, for which $s = R\Theta$ (here R is the radius of the scatterer).

Note that M defined this way is a (finite union of) Riemannian manifold(s).

Notation. Let

$$|dx|_e = \sqrt{ds^2 + d\varphi^2} \tag{2.1}$$

denote the Riemannian metric on M , which will be referred to as the Euclidean metric (e-metric).

Later on we will introduce another auxiliary metric quantity very common in the billiard literature, the p-metric.

As to dynamics, let T denote the first return map onto M .

Notation for the Lebesgue measure on M is m , i.e. $dm = ds d\varphi$. Furthermore, given a curve γ in M we denote the Lebesgue measure on γ with m_γ (this is simply the length on γ).

Denote by μ the natural invariant probability measure on M . μ is absolutely continuous w.r.t Lebesgue, and the density is of the form

$$d\mu = \text{const.} \cos(\varphi) dm = \text{const.} \cos(\varphi) ds d\varphi. \quad (2.2)$$

It is this latter measure for which T is assumed to be ergodic and K-mixing and this is the one we work with as well.

Remark. In a completely similar manner we could consider the Poincaré section M_- of incoming particles. The two coordinates would be the point of income and the angle the incoming velocity makes with the (opposite) normal vector. However, in some key steps of the proof – e.g. the distortion bounds of Subsect. 3.3.2 – we heavily use that our phase space is the outgoing, and *not* the incoming Poincaré section.

With slight abuse of notation we often refer to the incoming Poincaré coordinates with the same symbols s and φ . That should cause no confusion.

Rotation function, its basic properties and formulation of the main theorem. To describe the first return map T we decouple the motion into two parts: free flight among the scatterers and flight in the potential of the scatterers. Free flight can be treated completely analogously to the billiard case. The particle leaves one of the scatterers in the point s_0 with velocity φ_0 and reaches some other scatterer in point s (or equivalently, Θ) with unit incoming velocity that makes an angle φ with the (opposite) normal vector $n(s)$ at the point of income. After some inter-potential motion the particle leaves the circle in some point $s_1 = (R\Theta_1)$ with outgoing velocity specified by φ_1 . Out of symmetry reasons $\varphi_1 = \varphi$, thus the only nontrivial quantity is the angle difference $\Delta\Theta = \Theta_1 - \Theta$. Again out of symmetry reasons $\Delta\Theta$ depends only on the angle φ .

The role in the map T played by the potential is completely described by the function $\Delta\Theta(\varphi)$.

Definition 1. From here on we will refer to this function $\Delta\Theta(\varphi)$ as **the rotation function**.

Being mainly interested in the differential aspects of T we introduce one more **Notation**

$$\kappa(\varphi) = \frac{d\Delta\Theta(\varphi)}{d\varphi}.$$

Below two important properties are defined in terms of which our main theorem is formulated.

Definition 2. The soft billiard system satisfies property H in case

1. there is some positive constant c such that $|2 + \kappa(\varphi)| > c$ for all φ ;
2. the configuration of scatterers is such that the distance of any two circles is bounded below by τ_{\min} , where

$$\tau_{\min} \geq \max_{\varphi} \left\{ -2R\kappa(\varphi) \frac{\cos \varphi}{2 + \kappa(\varphi)} \right\}.$$

Remarks. – Although a bit otherwise formulated, it was essentially proven in [DL] that soft billiard systems with property H are hyperbolic and ergodic. The mechanism of hyperbolicity is briefly explained in Sect. 3.2.

- Note that in case $\kappa > 0$ or $\kappa < -2$ for all φ , the lower bound for τ_{\min} turns out to be negative. Thus the second assumption is only restrictive in the opposite case, and the closer κ may get to -2 from above the more restrictive it is.
- In case there is some φ for which $0 > \kappa > -2$, a positive lower bound on the free path is to be assumed. Thus a planar periodic configuration of circles is needed that has finite horizon and (a possibly great) given τ_{\min} simultaneously. At first sight it seems questionable whether such configurations exist at all, nevertheless, as proven in [BöTa], this happens with positive probability in a random construction.

Definition 3. *The rotation function is termed regular in case the following properties hold:*

1. $\Delta\Theta(\varphi)$ is piecewise uniformly Hölder continuous. I.e. there are constants $C < \infty$ and $\alpha > 0$, and furthermore, $[-\frac{\pi}{2}, \frac{\pi}{2}]$ can be partitioned into finitely many intervals, such that for any φ_1 and φ_2 (from the interior of one of the intervals):

$$|\Delta\Theta(\varphi_1) - \Delta\Theta(\varphi_2)| \leq C|\varphi_1 - \varphi_2|^\alpha.$$

2. $\Delta\Theta(\varphi)$ is a piecewise C^2 function of φ on the closed interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$, in the above sense. (Note, however, that κ , in contrast to $\Delta\Theta$, can happen to have no finite one-sided limits at discontinuity points.)
3. There is some finite constant C such that

$$|\kappa'(\varphi)| \leq C|(2 + \kappa(\varphi))^3|,$$

where $\kappa'(\varphi)$ is the derivative of κ with respect to φ .

4. For the final property consider any discontinuity point φ_0 , where $\kappa(\varphi)$ (in contrast to $\Delta\Theta(\varphi)$) has no finite limit from the left. Of course, in case there is no finite limit from the right, the analogous property is similarly assumed.

Restricted to some interval $[\varphi_0 - \epsilon, \varphi_0)$; $\omega(\varphi) = \frac{2+\kappa(\varphi)}{\cos\varphi}$ is a monotonic function of φ .

Remark. Note that in case κ is C^1 (or piecewise C^1 with boundedness of itself and of κ') regularity is automatic. In case the asymptotics of κ near some discontinuity is some power law $(\varphi_0 - \varphi)^{-\xi}$ (with $\xi > 0$), regularity means $\frac{1}{2} \leq \xi < 1$.

We need two more definitions for the statement of our theorem:

Definition 4. *Consider a phase space M with a dynamics T and a T -invariant probability measure μ . We say that the dynamical system (M, T, μ) has exponential decay of correlations (EDC), if for every $f, g : M \rightarrow \mathbb{R}$ Hölder-continuous pair of functions there exist constants $C < \infty$ and $a > 0$ such that for every $n \in \mathbb{N}$,*

$$\left| \int_M f(x)g(T^n x)d\mu(x) - \int_M f(x)d\mu(x) \int_M g(T^n x)d\mu(x) \right| \leq Ce^{-an}.$$

Definition 5. We say that (M, T, μ) satisfies the central limit theorem (CLT) (for Hölder continuous functions) if for every $\eta > 0$ and every Hölder-continuous function $f : M \rightarrow \mathbb{R}$ with $\int f d\mu = 0$, there exists a $\sigma_f \geq 0$ such that

$$\frac{1}{\sqrt{n}} \sum_{i=0}^{n-1} f \circ T^i \xrightarrow{\text{distr}} \mathcal{N}(0, \sigma_f),$$

where $\mathcal{N}(0, \sigma_f)$ is the Gaussian distribution with variance σ_f^2 .

Now we are ready to formulate our main theorem.

Theorem 1. Suppose that the soft billiard system (M, T, μ) satisfies property H and the rotation function is regular. Suppose furthermore that there are no corner points and the horizon is finite ($0 < \tau_{\min}, \tau_{\max} < \infty$).

Then, dynamics enjoys, in addition to ergodicity and hyperbolicity, exponential decay of correlations and the central limit theorem for Hölder-continuous functions.

Proof. Ingredients for the proof are in Sect. 3 and 4. Actually, following tradition (e.g. [Ch]) we modify the dynamical system in several steps (Conventions 1 and 2). We will use a phase space \bar{M} , which is the original M cut into (countably many) connected components by singularities and so called “secondary singularities”. We will also use a higher iterate of the dynamics $T_1 = T^{m_0}$ with some m_0 to be found later.

It is the modified dynamical system (\bar{M}, T_1, μ) for which the conditions for EDC and LCT given in [Ch] are checked. Precisely, EDC and CLT for (\bar{M}, T_1, μ) are the consequence of Propositions 1, 2, 3, 4 and 5 and Theorem 2.1 from [Ch].

Exponential decay of correlations and the central limit theorem for (M, T, μ) follow easily from EDC and CLT for (\bar{M}, T_1, μ) . \square

For the reader’s convenience, we give a formulation of Theorem 2.1 from [Ch] in the Appendix.

Now we turn to the details of the above proof.

Some conventions. Constants that depend only on the map T itself (like $\tau_{\min}, \tau_{\max}, \dots$) will be called *global constants*.

Positive and finite global constants, whose value is otherwise not important, will be often denoted by just c or C (typically c for lower bounds and C for upper). That is, in two different lines of the same section, C can mean two different numbers.

Two quantities f and g defined on (the tangent bundle of) M (or on some subset like the unstable cone field, see Subsect. 3.1) will be called *equivalent* ($f \sim g$) if there are some global positive constants c and C such that $cf \leq g \leq Cf$.

2.1. Singularities. Just like in billiards the dynamics T is not smooth at certain one-dimensional submanifolds (curves) of M . Consider the set of tangential reflections:

$$S_0 = \left\{ (s, \varphi) \in M \mid \varphi = \pm \frac{\pi}{2} \right\}.$$

Actually $S_0 = \partial M$ (the boundary of the phase space). It is not difficult to see that T is not continuous at $S_1 = T^{-1}S_0$, i.e. at the preimages of tangential reflections. However, additional singularities appear at

$$Z_0 = \{ (s, \varphi) \in M \mid \varphi = \varphi_0 \}$$

in case φ_0 is some discontinuity point for $\Delta\Theta(\varphi)$, $\kappa(\varphi)$ or $\kappa'(\varphi)$. In such a case we will consider the phase space as if it were cut into two regions, more precisely Z_0 is treated as part of the boundary. As κ is not differentiable at Z_0 , T is not C^1 at the preimage of this set, at $Z_1 = T^{-1}Z_0$.

Furthermore we introduce the notations

$$S^{(n)} = S_1 \cup T^{-1}S_1 \cup \dots \cup T^{-n+1}S_1$$

and $Z^{(n)}$, analogously. The n^{th} iterate of the dynamics is not smooth precisely at $Z^{(n)} \cup S^{(n)}$.

The geometrical structure of $Z^{(n)}$ is much similar to that of $S^{(n)}$. Indeed, one can think of Z_1 as the set of those trajectories that would touch tangentially a smaller disk (one of radius $R \sin(|\varphi_0|)$) at the next collision. The following properties of the singularity set are of crucial importance:

- $Z^{(n)} \cup S^{(n)}$ is a finite union of C^2 curves.
- *Continuation property.* Each endpoint, x_0 , of every unextendable smooth curve $\gamma \subset Z^{(n)} \cup S^{(n)}$, lies either on the extended boundary $Z_0 \cup S_0$ or on another smooth curve $\gamma' \subset Z^{(n)} \cup S^{(n)}$ that itself does not terminate at x_0 .
- *Complexity property.* Let us denote by K_n the complexity of $Z^{(n)} \cup S^{(n)}$, i.e. the maximal number of smooth curves in $Z^{(n)} \cup S^{(n)}$ that intersect or terminate at any point of $Z^{(n)} \cup S^{(n)}$. K_n grows sub-exponentially with n .

For the proof of these properties in the billiard setting see the literature, especially [ChY], our case is analogous.

One more similarity with “hard” billiards is that for technical reasons later on we will introduce countably many secondary singularities parallel to the lines of $S^{(n)}$. Such secondary singularities are to be introduced parallel to $Z^{(n)}$ as well, in case $|\kappa|$ is unbounded as $\varphi \rightarrow \varphi_0$, at least from one side. We will turn back to this question in Subsubsection. 3.2.6.

3. Fronts, u-Manifolds and Unstable Manifolds

3.1. u-manifolds and their geometric properties.

Fronts and their geometric description. Our most important tools in describing hyperbolicity – local orthogonal manifolds or simply *fronts* – we inherit from billiard theory. A front \mathcal{W} is defined in the flow phase space rather than in the Poincaré section.

Definition 6. *Take a smooth 1-codim submanifold E of the whole configuration space, and add the unit normal vector $v(q)$ of this submanifold at every point q as a velocity, continuously. Consequently, at every point the velocity points to the same side of the submanifold E . The set*

$$\mathcal{W} = \{(q, v(q)) | q \in E\} \subset \mathcal{M}, \quad (3.1)$$

where $v : E \rightarrow \mathbf{S}^1$ is continuous (smooth) and $v \perp E$ at every point of E , is called a *front*.

Analysis of the time evolution of fronts is the key to almost all the geometric properties of the system that we need. For this reason, we first discuss time evolution of an arbitrary front. Later subsections will deal with special cases.

Consider a front with a reference point just before reaching a scatterer, and another “perturbed” point nearby. With the notations introduced before (see also Fig. 1), the




Fig. 1. Conventions for notation and signs for fronts

perturbation bringing the reference trajectory into the perturbed one is (dq_-, dv_-) just before collision, $(ds_-, d\varphi_-)$ in the incoming Poincaré section, $(ds_+, d\varphi_+)$ in the outgoing Poincaré section, (dq_+, dv_+) just after collision, and (dq'_-, dv'_-) just before the next collision. The evolution of the perturbations is:

$$\begin{aligned}
 ds_- &= \frac{dq_-}{\cos \varphi_-}, \\
 d\Theta_- &= \frac{ds_-}{R}, \\
 d\varphi_- &= dv_- + d\Theta_-, \\
 d\Theta_+ &= d\Theta_- + \kappa d\varphi_-, \\
 d\varphi &:= d\varphi_+ = d\varphi_-, \\
 ds_+ &= R d\Theta_+, \\
 dq_+ &= -\cos \varphi ds_+, \\
 dv_+ &= -d\Theta_+ - d\varphi_+,
 \end{aligned} \tag{3.2}$$

while crossing the potential. For the evolution equations of free flight, we introduce the

Notation. $\tau = \tau(x)$ will denote the length of free flight of the particle before reaching the next scatterer.

So, during free flight we have

$$\begin{aligned}
 dq'_- &= dq_+ + \tau dv_+, \\
 dv'_- &= dv_+.
 \end{aligned} \tag{3.3}$$

Note that the angles of incidence and reflection are measured in different directions – in order to keep them equal, as they traditionally are, – but dq_- and dq_+ (just like dv_- and dv_+) are measured in the same direction, unlike usually in billiards.

Based on these, we can find out about the evolution of the derivative $B = \frac{dv}{dq}$.

Notations. B will denote the derivative of the unit normal vector (velocity) $v(q)$ of a front: $dv = Bdq$ for tangent vectors (dq, dv) of the front.

$m = \frac{d\varphi}{ds}$ will denote the slope of the (trace of the) front in the Poincaré section.

B is nothing other than the curvature of the submanifold E . Yet we will prefer to call it the second fundamental form (SFF), in order to avoid confusion with other curvatures that are coming up. The term “form” refers to higher dimensional cases when B is a symmetric operator.

Equation (3.2) gives

$$\begin{aligned} m_- &= \cos \varphi B_- + \frac{1}{R}, \\ \frac{1}{m_+} &= \frac{1}{m_-} + R\kappa, \\ \cos \varphi B_+ &= m_+ + \frac{1}{R}, \end{aligned} \quad (3.4)$$

while crossing the potential, which can be summarized in

$$B_+ = \frac{2 + \kappa(\varphi) + (1 + \kappa(\varphi))R \cos \varphi B_-}{R \cos \varphi (1 + \kappa(\varphi) + \kappa(\varphi)R \cos \varphi B_-)}, \quad (3.5)$$

and (3.3) gives

$$\frac{1}{B'_-} = \frac{1}{B_+} + \tau \quad (3.6)$$

during free flight.

Notations.

$$\lambda_1 := \frac{dq_+}{dq_-}, \quad (3.7)$$

$$\lambda_2 := \frac{dq'_-}{dq_+}, \quad (3.8)$$

$$\lambda := \lambda_1 \lambda_2.$$

These are exactly the expansion factors along the front, for the respective “pieces” of the dynamics. (They are also expansion factors in the Poincaré section, but in the p-metric to be introduced later.) We have

$$\lambda_1 = 1 + \kappa + \kappa R \cos \varphi B_- = 1 + \kappa R m_- = \frac{m_-}{m_+}, \quad (3.9)$$

$$\lambda_2 = 1 + \tau B_+ = \frac{B_+}{B'_-}. \quad (3.10)$$

To study decay of correlations, we need one more derivative.

Notation. $D = \frac{dB}{dq}$.

This is exactly the *curvature* of the *front* as of a *subset of the flow phase space* (and not as of a subset of the configuration space – unlike B , cf. (3.1)).

To study the evolution of D we need to consider two small pieces of the front, one around the reference point, and one around the perturbed one. Let the change in the SFF be

$$dB_{--} = D_- dq$$

before scattering, and

$$dB_{++} = D_+ dq$$

after scattering. dB_{--} is *not* the difference of SFF-s at the points of incidence, because the perturbed point has to travel another $d\tau_- = \tan \varphi_- dq_-$ to reach the scatterer ($d\tau$ can be negative), which changes its SFF according to the rules (3.6) of free flight. Taking that into account, we have

$$dB_- = dB_{--} - B_-^2 d\tau_- = dB_{--} - B_-^2 \tan \varphi_- dq_-. \quad (3.11)$$

Similarly, for the fronts leaving the potential,

$$dB_{++} = dB_+ - B_+^2 d\tau_+ = dB_+ - B_+^2 \tan \varphi_+ dq_+. \quad (3.12)$$

(Note our convention on the signs of dq_- , dq_+ , φ_- and φ_+ .)

To follow the evolution of curvature we introduce

Notations. $D_1 = \frac{dB_-}{dq_-} (\neq D_-)$, $K_- = \frac{dm_-}{ds_-}$, $K_+ = \frac{dm_+}{ds_+}$, $D_2 = \frac{dB_+}{dq_+}$ and $\eta(\varphi) = \frac{d\kappa(\varphi)}{d\varphi}$.

With these we get from (3.2), (3.4), (3.7), (3.9), (3.11) and (3.12),

$$\begin{aligned} D_1 &= D_- - \tan \varphi B_-^2, \\ K_- &= \cos^2 \varphi D_1 - \sin \varphi B_- m_-, \\ K_+ &= \frac{1}{\lambda_1^3} K_1 - R \left(\frac{m_-}{\lambda_1} \right)^3 \eta, \\ \cos^2 \varphi D_2 &= -K_+ - \sin \varphi B_+ m_+, \\ D_+ &= D_2 - \tan \varphi B_+^2, \end{aligned} \quad (3.13)$$

while crossing the potential, and, from (3.3), (3.6) and (3.8)

$$D'_- = \frac{1}{\lambda_2^3} D_+ \quad (3.14)$$

during free flight.

3.2. Invariance of convex fronts, u-fronts and u-manifolds. In [DL] it is shown – although not explicitly stated in this integrated form – that if Property H (defined in Definition 2) is satisfied, then convex fronts with suitably small SFF-s (the upper bound may be ∞) either remain convex, or focus before reaching the next scatterer, and become convex again, with suitably small SFF. This property is called the “invariance of convex fronts”. In the present work we also require (see Theorem 1) that τ be bounded from below by some $\tau_{\min} > 0$ even in the case when [DL] did not (the “no corner points” assumption), and an upper bound τ_{\max} (the “finite horizon” assumption). In order to establish estimates that we will need later, we must repeat some steps of the argument in [DL]. We omit details of the calculations; these can be done by the reader or can be found in the above paper.

Notations.

$$\begin{aligned} \tau_1 &= \max \left\{ 0, \max_{\varphi} \left\{ -R\kappa(\varphi) \frac{\cos \varphi}{2 + \kappa(\varphi)} \right\} \right\}, \\ B^* &= \frac{1}{\tau_1} \quad (\infty \text{ if } \tau_1 = 0). \end{aligned} \quad (3.15)$$

From (3.4) we get that if $0 < B_- < B^*$ then either $m_+ > 0$ and thus $B_+ > \frac{1}{R}$ or $B_+ < -B^*$. This – by (3.6) – implies that $c < B'_- < B^{**}$ with some global constants $c > 0$ and $B^{**} < B^*$, assuming that $\tau_{\min} > 2\tau_1$, which is exactly Property H. All in all,

$$c < B_- < B^{**} \text{ implies } c < B'_- < B^{**}. \quad (3.16)$$

This motivates our

Definition 7. A *u-front* is a front with $c < B_- < B^{**}$. A *u-manifold* is the trace of a *u-front* on the Poincaré phase space.

Definition 8. An *s-front* is a front with $c < -B_+ < B^{**}$. An *s-manifold* is the trace of an *s-front* on the Poincaré phase space.

As we have seen, u-manifolds remain u-manifolds under time evolution. s-fronts are exactly the u-fronts of the inverse dynamics.

The aim of this subsection is to show important properties of u-fronts and u-manifolds, which are stronger than those shown for an arbitrary front in the previous subsection. In Subsect. 3.3 we further restrict to the case of unstable manifolds, which are special kinds of u-manifolds.

3.2.1. Expansion estimates along u-fronts. First we work out estimates for the expansion along a front from one moment of incidence to the next. We will use these estimates later to estimate expansion of our dynamics T in our outgoing Poincaré phase space M .

Consider a u-front with the earlier notations. We start with an easy observation we will often use: from (3.4) and (3.16) we get $\frac{1}{R} < m_- < \frac{1}{R} + B^{**}$, which implies

$$m_- \sim 1. \quad (3.17)$$

To get the order of magnitude for the expansion factor λ , put the formulas in (3.5) and (3.9) together, and get that

$$\frac{R \cos \varphi B_+ \lambda_1}{2 + \kappa(\varphi)} = 1 + (1 + \kappa(\varphi)) R B_- \frac{\cos \varphi}{2 + \kappa(\varphi)}.$$

The right-hand side is trivially bounded from above since B_- is bounded, and so is $\frac{1+\kappa(\varphi)}{2+\kappa(\varphi)} = 1 - \frac{1}{2+\kappa(\varphi)}$. On the other hand,

- It is greater than 1 if $\frac{1+\kappa(\varphi)}{2+\kappa(\varphi)} > 0$.
- If $\frac{1+\kappa(\varphi)}{2+\kappa(\varphi)} \leq 0$ (that is, $-2 < \kappa(\varphi) \leq -1$), then

$$\begin{aligned} 1 + (1 + \kappa(\varphi))RB_- \frac{\cos \varphi}{2 + \kappa(\varphi)} &\geq 1 + (1 + \kappa(\varphi))R \frac{\cos \varphi}{2 + \kappa(\varphi)} B^* \\ &\geq 1 + (1 + \kappa(\varphi))R \frac{\cos \varphi}{2 + \kappa(\varphi)} \frac{2 + \kappa(\varphi)}{-R\kappa(\varphi) \cos \varphi} = \frac{-1}{\kappa(\varphi)} \geq \frac{1}{2}. \end{aligned}$$

All in all, using $\lambda_2 = \frac{B_+}{B_-} \sim B_+$ (see (3.10) and (3.16)) we have

$$\lambda \sim B_+ \lambda_1 \sim \frac{2 + \kappa(\varphi)}{\cos \varphi}, \quad (3.18)$$

which is one of our key estimates. Notice that the right-hand side cannot be too small due to Property H (Definition 2).

We can also get the order of magnitude for λ_1 and λ_2 separately: (3.9) and (3.17) gives

$$|\lambda_1| \sqrt{1 + m_+^2} = \sqrt{\lambda_1^2 + m_-^2} \sim |2 + \kappa(\varphi)|. \quad (3.19)$$

(The last equivalence is true because both sides are bounded away from zero, and can only be big when they grow linearly with κ .) Notice that λ_1 can be very small (even zero), and can even change signs while $2 + \kappa(\varphi)$ remains positive. Of course, m_+ has to be infinity (and change signs) simultaneously.

Putting (3.18) and (3.19) together, we get

$$\frac{|\lambda_2|}{\sqrt{1 + m_+^2}} \sim \frac{1}{\cos \varphi}. \quad (3.20)$$

This last line can be rewritten as

$$1 \sim \frac{|\lambda_2| \cos \varphi}{\sqrt{1 + m_+^2}} \sim \frac{|B_+| \cos \varphi}{\sqrt{1 + m_+^2}} = \frac{|m_+ + \frac{1}{R}|}{\sqrt{1 + m_+^2}},$$

which implies that there is a global constant c such that

$$\left| m_+ + \frac{1}{R} \right| > c. \quad (3.21)$$

3.2.2. Expansivity. To obtain hyperbolicity, we must see that u-manifolds are expanded by the dynamics. In the first round we prove a lemma about the expansion on u-fronts from collision to collision.

Lemma 1. *There exists a global constant $\Lambda > 1$, such that for every u-front, $|\lambda| \geq \Lambda$.*

Proof. Besides $\tau > 0$ and $B_- > 0$ we will use that $\tau \geq 2\tau_1 + d$, where $d := \tau_{min} - 2\tau_1 > 0$, and $\tau_1 \geq -R\kappa(\varphi) \frac{\cos \varphi}{2 + \kappa(\varphi)}$ for every φ (see Definition 2 and (3.15)). Altogether:

$$\tau \geq d - 2R\kappa(\varphi) \frac{\cos \varphi}{2 + \kappa(\varphi)}. \quad (3.22)$$

We will also use from (3.15) and Definition 7 that

$$0 < B_- \leq \frac{2 + \kappa(\varphi)}{-R\kappa(\varphi) \cos \varphi} \quad (3.23)$$

whenever the right-hand side is positive, which is the $-2 < \kappa(\varphi) < 0$ case. Now we start by putting together (3.9), (3.10) and (3.5) to get

$$\begin{aligned} \lambda &= (1 + \kappa(\varphi)Rm_-)(1 + \tau B_+) \\ &= 1 + \kappa(\varphi) + \kappa(\varphi)R \cos \varphi B_- + \tau \left(\frac{2 + \kappa(\varphi)}{R \cos \varphi} + (1 + \kappa(\varphi))B_- \right). \end{aligned}$$

We estimate this taking care of the signs of the particular terms.

– If $\kappa(\varphi) \leq -2 - \delta$, then

$$\lambda \leq 1 + \kappa(\varphi) \leq -1 - \delta.$$

– If $-2 + \delta \leq \kappa(\varphi) \leq -1$ then both coefficients of B_- are negative, so we can use (3.23) to estimate the right-hand side from below. In the next step we find the coefficient of τ positive, so we can use (3.22). What we get is

$$\begin{aligned} \lambda &\geq 1 + \kappa(\varphi) + \kappa(\varphi)R \cos \varphi \frac{2 + \kappa(\varphi)}{-R\kappa(\varphi) \cos \varphi} \\ &\quad + \tau \left(\frac{2 + \kappa(\varphi)}{R \cos \varphi} + (1 + \kappa(\varphi)) \frac{2 + \kappa(\varphi)}{-R\kappa(\varphi) \cos \varphi} \right) \\ &= -1 + \tau \frac{2 + \kappa(\varphi)}{-\kappa(\varphi)R \cos \varphi} \\ &\geq -1 + \left(d - 2R\kappa(\varphi) \frac{\cos \varphi}{2 + \kappa(\varphi)} \right) \frac{2 + \kappa(\varphi)}{-\kappa(\varphi)R \cos \varphi} \\ &\geq 1 + \frac{d\delta}{2R}. \end{aligned}$$

– If $-1 \leq \kappa(\varphi) \leq 0$, then the coefficient of τ is positive, so we first use (3.22) to estimate the right-hand side from below. In the next step we find one coefficient of B_- positive, so we just use $B_- > 0$, and one coefficient of B_- negative, so we can use (3.23). What we get is

$$\begin{aligned} \lambda &\geq 1 + \kappa(\varphi) + \kappa(\varphi)R \cos \varphi B_- \\ &\quad + \left(d - 2R\kappa(\varphi) \frac{\cos \varphi}{2 + \kappa(\varphi)} \right) \left(\frac{2 + \kappa(\varphi)}{R \cos \varphi} + (1 + \kappa(\varphi))B_- \right) \\ &= 1 + d \frac{2 + \kappa(\varphi)}{R \cos \varphi} + d(1 + \kappa(\varphi))B_- - \kappa(\varphi) - \frac{\kappa^2(\varphi)R \cos \varphi}{2 + \kappa(\varphi)} B_- \\ &\geq 1 + d \frac{2 + \kappa(\varphi)}{R \cos \varphi} - \kappa(\varphi) - \frac{\kappa^2(\varphi)R \cos \varphi}{2 + \kappa(\varphi)} \frac{2 + \kappa(\varphi)}{-R\kappa(\varphi) \cos \varphi} \\ &\geq 1 + \frac{d}{R}. \end{aligned}$$

– If $0 < \kappa(\varphi)$, then

$$\lambda \geq 1 + \frac{2d}{R}. \quad \square$$

3.2.3. Transversality.

Lemma 2. *We will see that u- and s-manifolds are uniformly transversal. I.e. there is some global constant $\alpha_0 > 0$ such that given any two tangent vectors (in the outgoing Poincaré phase space) dx_s and dx_u of an s- and a u-manifold, respectively, we have*

$$\langle dx_u, dx_s \rangle > \alpha_0.$$

Proof. To see this, use Definition 8 and (3.4) to get $m_+^s \sim -1$ for the slope of any s-manifold. This way, it is enough to see that the slopes of u- and s-manifolds are bounded away, that is, $|m_+^u - m_+^s| > c$. To get this, use – again – Definition 8, Definition 7, the estimates before them and (3.4) to get

$$\begin{aligned} -\frac{1}{R} > m_+^s > -\frac{1}{R} - \cos \varphi B^{**}, \\ m_+^u > 0 \text{ or } m_+^u < -\frac{1}{R} - \cos \varphi B^*, \end{aligned} \quad (3.24)$$

so either $m_+^u - m_+^s > \frac{1}{R}$ or $m_+^s - m_+^u > \cos \varphi (B^* - B^{**})$. This implies the statement when $\cos \varphi$ is not too small. However, when $\cos \varphi$ is small, we have to use the estimate (3.21) and (3.24) to see also that

$$m_+^u > 0 \text{ or } m_+^u < -\frac{1}{R} - c,$$

which completes the proof. \square

3.2.4. Hyperbolicity. In what follows we will consider time evolution of vectors tangent to u-manifolds. Notation both in the incoming and the outgoing phase space will be of the type $dx = (ds, d\varphi)$. In addition to the e-metric (2.1) we will use one more metric quantity, the p-metric:

$$|dx|_p = |ds| \cos(\varphi).$$

The p-metric measures distances along the corresponding u-front. It is degenerate on the whole tangent bundle. However, when restricted a u-manifold *in the incoming phase space*, by (3.17) we have:

$$|dx|_p \sim |dx|_e \cos(\varphi).$$

According to Lemma 1, u-vectors are expanded uniformly (*from collision to collision, that is, in the incoming phase space*) in the p-metric:

$$|DT|_p = \lambda \geq \Lambda > 1.$$

To obtain expansion in the e-metric and the outgoing phase space, we look at the n^{th} iterate of the outgoing phase space dynamics the following way:

- switch to p-metric
- reach the next scatterer
- do $n - 1$ steps in the incoming phase space
- cross the potential
- switch back to Euclidean metric.

This way we get

$$|DT^n dx|_e = \frac{\sqrt{1+m_+^2(n)}}{\cos \varphi(n)} \lambda_{1(n)} \lambda_{(n-1)} \lambda_{(n-2)} \dots \lambda_{(1)} \lambda_2 \frac{\cos \varphi}{\sqrt{1+m_+^2}} |dx|_e,$$

where symbols with ()-ed subscripts mean values at the appropriate iterate of the phase point. Using (3.19), (3.20) and Lemma 1 we get

$$|DT^n dx|_e \sim \lambda_{(n-1)} \lambda_{(n-2)} \dots \lambda_{(1)} \frac{2 + \kappa(\varphi(n))}{\cos \varphi(n)} |dx|_e. \quad (3.25)$$

This way we have

$$|DT^n dx|_e > c_1 \Lambda^n |dx|_e \quad (3.26)$$

with some global constant c_1 . Again, this is for u-vectors in the outgoing phase space.

The transversality of s- and u- vectors, stated in Proposition 1 implies that the product of (length) expansion factors for s- and u- vectors is equivalent to the n -step (Lebesgue) volume expansion factor. Using (2.2), and the T -invariance of μ , we get that if dx is a u-vector and dy is an s-vector, then

$$\frac{|DT^n dx|_e}{|dx|_e} \frac{|DT^n dy|_e}{|dy|_e} \sim \frac{\cos \varphi}{\cos \varphi(n)}.$$

Combinig this with (3.25) we get

$$|DT^n dy|_e \sim \frac{\cos \varphi}{2 + \kappa(\varphi(n))} \frac{1}{\lambda_{(n-1)} \lambda_{(n-2)} \dots \lambda_{(1)}} |dy|_e,$$

which implies

$$|DT^n dy|_e < \frac{C_1}{\Lambda^n} |dx|_e \quad (3.27)$$

with some global constant C_1 . Again, this is for s-vectors in the outgoing phase space.

Convention 1. We choose a positive integer m_0 the following way. First take m_1 such that $c_1 \Lambda^{m_1} > 1$ and $\frac{C_1}{\Lambda^{m_1}} < 1$. This way any high enough power of the dynamics, T^m with $m > m_1$ is uniformly expanding along u-manifolds and uniformly contracting along s-manifolds with $\Lambda_1 = \Lambda^{m-m_1}$. Now recall the notion and the basic properties of complexity K_n from Subsect. 2.1. As K_n grows subexponentially we may choose m_2 for which we have $K_m < \Lambda^{m-m_1}$ whenever $m > m_2$. We fix $m_0 = \min(m_1, m_2) + 1$.

The advantage of this choice is that the iterate $T_1 = T^{m_0}$ is uniformly hyperbolic (see the proposition to come) with constant Λ_1 for which $\Lambda_1 > K_{m_0} + 1$. This later fact we only use in Sect. 4.

Let us summarize what we have seen so far from the hyperbolic properties in the following

Proposition 1. There exist two families of cones $C_s(x)$ and $C_u(x)$ – called stable and unstable cones – in the tangent space of M such that

$$DT(C_u(x)) \subset C_u(Tx) \text{ and } C_s(Tx) \subset DT(C_s(x)).$$

The stable/unstable cone is uniformly contracting/expanding:

$$\begin{aligned} |DT_1^{-1}(dx)| &\geq \Lambda_1 |dx| \quad \forall dx \in C_s(x), \\ |DT_1(dx)| &\geq \Lambda_1 |dx| \quad \forall dx \in C_u(x). \end{aligned}$$

Furthermore, the two cone fields are uniformly transversal in the sense above.

Vectors of the stable/unstable cone are often called *s*- and *u*-vectors.

Proof. The two cones are formed by the tangent vectors of *s*- and *u*-manifolds, respectively. Invariance is the implication (3.16), recalling Definition 7 and 8. Expansion and contraction are (3.26), (3.27) and Convention 1. Transversality is Lemma 2. \square

We note that so far we have only used that our billiard satisfies property H, which is a property already formulated in [DL], and which is known from [Do1] to be essentially necessary for ergodicity.

3.2.5. Alignment. We need to investigate the relative position of *u*-manifolds and singularities in order to find out how much of a *u*-manifold can be “close” to a singularity. Our aim is to prove the following

Lemma 3. Take any smooth component Z of $T^{-k}Z_0$ with $k \geq 0$, where

$$Z_0 = \{ (s, \varphi) \in M \mid \varphi = \varphi_0 \}$$

with any $\varphi_0 \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. Given some small positive δ let us denote the δ -neighborhood of Z by $Z^{[\delta]}$. There are global constants $C < \infty$ and $\alpha > 0$ such that for any *u*-manifold W we have

$$m_W \left(Z^{[\delta]} \cap W \right) \leq C \delta^\alpha, \quad (3.28)$$

where m_W is the Lebesgue measure – the length – on the *u*-manifold W .

Proof. If $k > 0$, then Z is an *s*-manifold, and is transversal to our *u*-manifold W according to Lemma 2, so the statement holds even with $\alpha = 1$.

So take $k = 0$, then Z is described by $m_Z = 0$. If $\kappa(\varphi)$ remains bounded near φ_0 , then for our *u*-manifold W ,

$$\frac{1}{m_+} = \frac{1}{m_-} + R\kappa(\varphi)$$

is bounded (see (3.4) and (3.17)), so the two curves are transversal again, we can choose $\alpha = 1$.

The interesting case is $k = 0$, $\kappa(\varphi) \rightarrow \infty$ as $\varphi \rightarrow \varphi_0$. In this case (3.17) ensures that $\frac{1}{m_-}$ is negligible

- say, less than the ε portion – compared to $R\kappa(\varphi)$. This – through (3.4) and the definition $m_+ = \frac{d\varphi_+}{ds_+} = \frac{d\varphi}{ds}$
- implies that for *u*-manifolds

$$(1 - \varepsilon)R\kappa(\varphi) \leq \frac{ds}{d\varphi} \leq (1 + \varepsilon)R\kappa(\varphi).$$

Integrating this with respect to φ and using the definition of $\kappa(\varphi)$, we get

$$(1 - \varepsilon)R(\Delta\Theta(\varphi) - \Delta\Theta(\bar{\varphi})) \leq s - \bar{s} \leq (1 + \varepsilon)R(\Delta\Theta(\varphi) - \Delta\Theta(\bar{\varphi}))$$

which means that, close (enough) to a $\kappa(\varphi) \rightarrow \infty$ singularity, a u -manifold is (arbitrarily) similar to the graph of the rotation function $\Delta\Theta(\varphi)$. Now the Hölder-continuity of $\Delta\Theta(\varphi)$ required in the regularity condition (Definition 3) implies the statement of the lemma. \square

We note that the proof of alignment is the only place where we use our assumption that the rotation function is Hölder-continuous. The above proof shows that Hölder-continuity is indeed a necessary condition for alignment. Alignment is *not* among the conditions of Chernov's theorem which our proof is based on, but we will use it in the proof of the growth properties (Proposition 5). At that place it seems to be unavoidable, so we think that Hölder-continuity of the rotation function is needed for Chernov's method to work. On the other hand, as already pointed out in the introduction, we do not claim that it is a necessary condition for EDC.

3.2.6. Homogeneity strips, secondary singularities and homogeneous u -manifolds.

Notation.

$$\omega(\varphi) := \frac{2 + \kappa(\varphi)}{\cos \varphi} \quad (3.29)$$

We will see that expansion in the e -metric is unbounded as $|\omega(\varphi)| \rightarrow \infty$. This certainly happens in the vicinity of $\pm\frac{\pi}{2}$, nevertheless, there can exist other discontinuity values φ_0 with the same property. Big expansion comes together with big variations of expansion (i.e. distortion) rates along u -manifolds. For that reason we need to partition the phase space into homogeneity layers in which $\omega(\varphi)$ is nearly constant. We fix a large integer k_0 (to be specified in Sect. 4) and define for $k > k_0$ the I-strips as

$$I_k = \left\{ (s, \varphi) \mid k^2 \leq |\omega(\varphi)| < (k+1)^2 \right\}. \quad (3.30)$$

Recall from Definition 3 that whenever $\lim_{\varphi \rightarrow \varphi_0} |\omega(\varphi)| = \infty$, there exists an interval $[\varphi_0 - \varepsilon, \varphi_0)$ restricted to which $|\omega(\varphi)|$ is a monotonic function of φ . We partition a subinterval of this interval into I-strips, thus k_0 is chosen accordingly large. In case there are several discontinuity points of $\omega(\varphi)$ (with unbounded one-sided limits) we may construct further I-strips, $I_k^{(s)}$, analogously. Here the index s labels the finitely many discontinuities of this kind.

Furthermore take $I_0^{(u)}$; $u = 1 \cdots U$, where the index u labels the finitely many connected components of the complement of all the above layers (that is, the “remaining part” of the phase space).

We will use the notations Γ_0 for the countably many boundary components of I-strips.

Convention 2. *From now on, Γ_0 – just like S_0 and Z_0 before – is considered as part of the boundary of the phase space. That is, we will use a modified phase space \bar{M} , whose connected components are the homogeneity strips I_k (and $I_0^{(u)}$).*

In complete analogy with primary singularities we introduce furthermore the notations Γ_1 and $\Gamma^{(n)}$ for the corresponding preimages. The geometric properties of these secondary singularity lines are analogous to those of primary ones (for example, (3.28) applies).

Definition 9. We will say that a u -manifold is homogeneous whenever it is contained in one of the homogeneity strips I_k (or $I_0^{(u)}$).

In Sects. 3.3.2 and 4 we will be concerned with u -manifolds that remain homogeneous for several steps of the dynamics.

3.3. Regularity properties of unstable manifolds.

Definition 10. An unstable manifold is a u -manifold for which all past iterates are u -manifolds as well.

Analogously, a stable manifold is an s -manifold for which all future iterates are s -manifolds as well.

From the theory of hyperbolic systems (see [Ch] and references therein) we know that there is a unique unextendable unstable (and similarly a unique unextendable stable) manifold through (μ -)almost every point of M . Thus it makes sense to talk about the (un)stable manifold through the point.

We will also refer to unstable manifolds as “local unstable manifolds” (LUMs), stressing the fact that they are (and all their past iterates as well are) contained in some homogeneity layer I_k . (Remember that our phase space ends on the boundary of I_k , so I_{k+1} is already another connected component.)

In this subsection we deal with properties of unstable manifolds which are stronger than those proved before for arbitrary u -manifolds in Subsect. 3.2.

3.3.1. Curvature bounds. In what follows we obtain bounds on unstable manifolds that will guarantee that their curvature is uniformly bounded from above.

First we look at u -fronts as submanifolds of the flow phase space.

Putting the formulas in (3.13) and (3.14) together, we get

$$D'_- = -\frac{D_-}{\lambda^3} + \frac{2 \sin \varphi B_-^2}{\lambda^3 \cos \varphi} - \frac{2 \sin \varphi B_-'^2}{\lambda_2 \cos \varphi} + \frac{\sin \varphi B_-}{R \lambda^3 \cos^2 \varphi} + \frac{\sin \varphi B_-'}{R \lambda_2^2 \cos^2 \varphi} - R m_-^3 \frac{\eta}{\lambda^3 \cos^2 \varphi}.$$

Our key estimate (3.20) implies

$$\cos \varphi |\lambda_2| \sim \sqrt{1 + m_+^2},$$

which is bounded from below. So, in the above sum, terms number 2,3,4 and 5 are all bounded in absolute value. The last term is bounded due to our assumption

$$\left| \frac{\eta(\varphi)}{(2 + \kappa(\varphi))^3} \right| < C.$$

As a consequence, we have

$$|D'_-| \leq \frac{|D_-|}{\Lambda^3} + C_2, \quad (3.31)$$

with some global constant C_2 and can state

Lemma 4. There is a global constant \hat{D} such that for almost any point of the phase space, the front corresponding to the LUM has

$$|D_-| \leq \hat{D}.$$

Proof. Choose $\hat{D} = \frac{C_2 \Lambda^3}{\Lambda^3 - 1}$. Now suppose indirectly that there is a set $H \subset M$ of positive measure, for the points of which $|D_-| > \hat{D} + \varepsilon$. Then (3.31) implies that there is a $c(\varepsilon) > 0$ such that $|D_-| > \hat{D} + \varepsilon + c$ on $T^{-1}H$. This implies that $|D_-| > \hat{D} + \varepsilon + 2c$ on $T^{-2}H$, and so on: $|D_-| > \hat{D} + \varepsilon + kc$ on $T^{-k}H$ for all $k > 0$. But the $T^{-k}H$ -s are all sets of equal positive measure, which contradicts the finiteness of the phase space. \square

As a consequence, we can give curvature bounds for local unstable manifolds in the incoming and outgoing phase spaces. Since an unstable manifold in the Poincaré section is the graph of a function $\varphi = \varphi(s)$, its curvature is given by

$$g = \frac{\varphi''(s)}{\sqrt{1 + (\varphi'(s))^2}^3} = \frac{K}{\sqrt{1 + m^2}^3}.$$

We have reached

Proposition 2. *There is a global constant C such that for almost any point of the phase space, the front corresponding to the LUM has*

$$|g_+| \leq C.$$

Proof. It can be read from (3.13) that

$$|K_-| < C, \tag{3.32}$$

thus

$$|g_-| < C.$$

To find out about g_+ , we write

$$g_+ = \frac{K_+}{\sqrt{1 + m_+^2}^3} = \left(\frac{m_+}{\sqrt{1 + m_+^2}} \right)^3 \frac{K_-}{m_-^3} - R \frac{\eta}{\sqrt{1 + (\frac{1}{m_-} + R\kappa(\varphi))^2}^3}.$$

This is also bounded in absolute value due to our assumption

$$\left| \frac{\eta(\varphi)}{(2 + \kappa(\varphi))^3} \right| < C$$

(see Definition 3). \square

We note that this proof suggests that our condition

$$|\kappa'(\varphi)| \leq C|(2 + \kappa(\varphi))^3|$$

is necessary for bounded curvature, and consequently for Chernov's method to work.

3.3.2. *Distortion bounds.* Length of a u-manifold W is expanded by T^n locally with a factor

$$J_{W,n}(x) = \frac{|DT^n dx|_e}{|dx|_e},$$

where dx is the vector tangent to the curve of W at x . The aim of this subsection is to prove

Proposition 3. *Let W be an unstable manifold on which T^n is smooth. Assume that $W_i = T^i W$ is a homogeneous unstable manifold for each $1 \leq i \leq n$. Then for all $x, \bar{x} \in W$,*

$$|\ln J_{W,n}(x) - \ln J_{W,n}(\bar{x})| \leq C [\text{dist}_{W_n}(T^n x, T^n \bar{x})]^{1/5}.$$

Proof. Note that $J_{W,n}(x) = \prod_{i=0}^{n-1} J_{W_i,1}(T^i x)$. Hence, it is enough to prove the lemma for $n = 1$, because $\text{dist}(T^i x, T^i \bar{x})$ grows uniformly exponentially in i due to (3.26). So we put $n = 1$.

Denote $x' = Tx$ and, we will use a $'$ to denote quantities related to the point x' .

Recall from Sect. 2 that the expansion factor is easily calculated in the p-metric. To obtain $J := J_{W,1}(x)$ we transform $|dx|_e$ to $|dx|_p$, take the p-expansion factor from (3.7) and (3.8) and transform back. This way:

$$J = \frac{\sqrt{1+m'^2}}{\cos \varphi'} \lambda'_1 \lambda_2 \frac{\cos \varphi}{\sqrt{1+m^2}}.$$

In order to calculate the change in the logarithm of J as we move from x to \bar{x} , it is best to write it with the help of (3.29) in the form

$$J = \omega(\varphi') J'_1 J_2 \tag{3.33}$$

with

$$J_1 = \frac{\sqrt{1+m_+^2}}{2 + \kappa(\varphi)} \lambda_1$$

and

$$J_2 = \frac{\cos \varphi}{\sqrt{1+m_+^2}} \lambda_2.$$

Equations (3.19) and (3.20) imply

$$|J_1| \sim |J_2| \sim 1. \tag{3.34}$$

The change in logarithm of the three terms can be calculated independently, moreover, J_1 and J_2 are expected to change moderately, while $\omega(\varphi')$ can be kept under good control, because it depends only on φ' . The three terms are investigated in three sublemmas. Thus Proposition 3 is the direct consequence of the three Sublemmas 1, 2 and 3. Of course, the first and third (concerning J_1 and $\omega(\varphi)$) have to be applied with $'$ -es. When applying Sublemma 3, we use the trivial fact $|\varphi - \bar{\varphi}| \leq \text{dist}(x, \bar{x})$. \square

In the arguments below, as usual, quantities with neither $+$ nor $-$ in their index are meant to have a $+$, that is, in the outgoing phase space.

Sublemma 1. *There exists a global constant C such that when a perturbation of size dx is performed on the base point, we have*

$$|d \ln J_1| \leq C|dx|.$$

Proof. In many estimates, we will use – without further mention – that m_- and K_- are bounded (see (3.17) and (3.32)).

With the help of (3.9) we choose the form

$$J_1 = \frac{\sqrt{(1 + \kappa(\varphi)Rm_-)^2 + m_-^2}}{2 + \kappa(\varphi)}.$$

When calculating the differential, we use

$$d\kappa(\varphi) = \eta(\varphi)d\varphi = \frac{\eta(\varphi)m_+}{\sqrt{1 + m_+^2}}dx$$

and

$$dm_- = K_-ds_- = K_- \frac{ds_+}{\lambda_1} = \frac{K_-}{\lambda_1} \frac{dx}{\sqrt{1 + m_+^2}} = \frac{K_-}{\sqrt{\lambda_1^2 + m_-^2}}dx.$$

Calculating the differential, we get

$$\begin{aligned} d \ln J_1 &= \frac{m_- + \kappa(\varphi)R + \kappa^2(\varphi)R^2m_-}{((1 + \kappa(\varphi)Rm_-)^2 + m_-^2)^{3/2}} K_- dx \\ &+ \frac{2Rm_- - 1 - m_-^2 + (2Rm_- - 1)Rm_- \kappa(\varphi)}{(2 + \kappa(\varphi))((1 + \kappa(\varphi)Rm_-)^2 + m_-^2)} \frac{\eta(\varphi)m_+}{\sqrt{1 + m_+^2}} dx. \end{aligned}$$

The coefficient of dx in the first term is obviously bounded since the denominator is one degree higher in $\kappa(\varphi)$ and is bounded away from zero. In the second term, we use (3.9) and (3.19) to get

$$\left| \frac{m_+}{\sqrt{1 + m_+^2}} \right| = \left| \frac{m_-}{\sqrt{1 + m_+^2}\lambda_1} \right| \sim \left| \frac{1}{2 + \kappa(\varphi)} \right|, \quad (3.35)$$

so, looking again at the degrees of polynomials (in κ) in the numerator and denominator of the second term, we have

$$\frac{2Rm_- - 1 - m_-^2 + (2Rm_- - 1)Rm_- \kappa(\varphi)}{(2 + \kappa(\varphi))(1 + \kappa(\varphi)Rm_-)^2 + m_-^2} \frac{\eta(\varphi)m_+}{\sqrt{1 + m_+^2}} \leq C \left| \frac{\eta(\varphi)}{(2 + \kappa(\varphi))^3} \right| \leq C_1. \quad \square$$

Sublemma 2. *There exists a global constant C such that when a perturbation of size dx is performed on the base point, we have*

$$|d \ln J_2| \leq C|dx'|$$

(note the ' on the right-hand side).

Proof. With the help of (3.4) and (3.10) we choose the form

$$J_2 = \frac{\cos \varphi + \tau m_+ + \frac{\tau}{R}}{\sqrt{1 + m_+^2}}.$$

When calculating the differential, we use

$$d\varphi = \frac{m_+}{\sqrt{1 + m_+^2}} dx$$

and

$$dm_+ = K_+ ds_+ = K_+ \frac{dx}{\sqrt{1 + m_+^2}} = (1 + m_+^2) g_+ dx.$$

This way we get

$$d \ln J_2 = \frac{-\sin \varphi m_+}{\cos \varphi \lambda_2 \sqrt{1 + m_+^2}} dx + B'_- d\tau - \frac{m_+ (\frac{\tau}{R} + \cos \varphi) - \tau}{\cos \varphi \lambda_2} g_+ dx.$$

Due to (3.20), the coefficient of dx in the first term is equivalent to $\frac{-\sin \varphi m_+}{1 + m_+^2}$, and in the third term to $-\frac{m_+ (\frac{\tau}{R} + \cos \varphi) - \tau}{\sqrt{1 + m_+^2}} g_+$, both of which are bounded (cf. (3.35)).

We finish by estimating dx and $d\tau$ with dx' . First,

$$dx' = J dx \sim \left| \frac{2 + \kappa(\varphi')}{\cos \varphi'} \right| dx \geq c dx. \quad (3.36)$$

Second, the triangle inequality implies $|d\tau| \leq |ds| + |ds'_-|$. On the one hand, (3.36) implies $|ds| \leq |dx| \leq C|dx'|$. On the other hand (3.19) implies,

$$dx' = \sqrt{1 + m_+^2} |\lambda'_1| ds'_- \sim |2 + \kappa(\varphi')| ds'_- \geq c ds'_-.$$

These give

$$|d\tau| \leq C|dx'|. \quad \square$$

Sublemma 3. *There exists a global constant C such that if $x = (s, \varphi)$ and $\bar{x} = (\bar{s}, \bar{\varphi})$ are in the same homogeneity layer*

$$I_k = \left\{ (s, \varphi) \mid k^2 \leq |\omega(\varphi)| < (k+1)^2 \right\},$$

then

$$|\ln |\omega(\varphi)| - \ln |\omega(\bar{\varphi})|| \leq C|\varphi - \bar{\varphi}|^{1/5}.$$

Proof. We use the notation $\omega'(\varphi) = \frac{d}{d\varphi}\omega(\varphi)$. It is easy to see that the regularity of $\kappa(\varphi)$ implies

$$\left| \frac{\omega'(\varphi)}{\omega^3(\varphi)} \right| \leq C.$$

That is, everywhere inside J_k ,

$$\left| \frac{d|\ln \omega(\varphi)|}{d\varphi} \right| = \left| \frac{\omega'(\varphi)}{\omega(\varphi)} \right| \leq C|\omega(\varphi)|^2 \leq 2Ck^4.$$

This, together with the obvious $k^2 \leq |\omega(\varphi)|, |\omega(\bar{\varphi})| < (k+1)^2$, implies

$$\begin{aligned} |\ln |\omega(\varphi)| - \ln |\omega(\bar{\varphi})|| &\leq \min \left\{ 2Ck^4|\varphi - \bar{\varphi}|, \ln(k+1)^2 - \ln k^2 \right\} \\ &\leq \min \left\{ 2Ck^4|\varphi - \bar{\varphi}|, \frac{2}{k} \right\}. \end{aligned}$$

It is easy to check that for every k and every ξ ,

$$\min \left\{ 2Ck^4|\xi|, \frac{2}{k} \right\} \leq 2C^{1/5}\xi^{1/5},$$

which completes the proof. \square

After proving that the expansion factors vary nicely between nearby points on the same u-manifold, we now investigate their behaviour at points of different u-manifolds that lie on the same s-manifold. This is the absolute continuity property. Just like it was with the distortion bounds, it is important to consider homogeneous manifolds.

We introduce the simplified notation $J_k^u(x)$ and $J_k^s(x)$ for the k -step length expansion factor at x along the unstable and the stable manifold, respectively.

Proposition 4. *Let W_s be a small s-manifold, $x, \bar{x} \in W_s$, and W_u, \bar{W}_u two u-manifolds crossing W_s at x and \bar{x} , respectively. Assume that T^k is smooth on W_s and $T^i W_s$ is a homogeneous s-manifold for each $0 \leq i \leq k$. Then*

$$|\ln J_k^u(x) - \ln J_k^u(\bar{x})| \leq C,$$

where C is a global constant.

Proof. We have bounds on the change in expansion as we move along unstable manifolds. In order to have such bounds as we move along stable manifolds, we wish to use the fact that stable manifolds are turned into unstable ones when we revert time. However, this time reflection symmetry is not complete: we always work in the outgoing Poincaré section, and reverting time turns this into the incoming one. To deal with the problem, we introduce the map P which is the dynamics through the potential, and which maps from the incoming to the outgoing Poincaré section. That is,

$$P((s_-, \varphi)) := (s_+, \varphi) = (s_- + R\delta\Theta(\varphi), \varphi).$$

We can see from (3.4) that if $dx_- = (ds_-, d\varphi)$ is a tangent vector of the incoming phase space, then

$$|DP(dx_-)|_e = \sqrt{1 + m_+^2} |\lambda_1| \frac{1}{\sqrt{1 + m_-^2}} |dx_-|_e.$$

Denote by $v(x)$ the expansion factor of DP along the unstable manifold at x , that is $v(x) = \frac{|D_x P(dx_-)|_e}{|dx_-|_e}$, where dx is an unstable vector at x . We can use (3.19) and (3.17) to get

$$v(x) \sim |2 + \kappa(\varphi)|.$$

We also introduce the “turn back” operator, which we will denote by a “ $-$ ” sign: this turns incoming phase points into outgoing phase points which corresponds to reverting the velocity. “ $-$ ” is almost the identity function from M_- to M_+ , only the collision angle is reverted (see our sign convention in Fig. 1):

$$\begin{aligned} - & : M_- \rightarrow M_+, \\ -(s, \varphi_-) & := (s_+, \varphi_+) = (s_-, -\varphi_-). \end{aligned}$$

With these notations, if $x = P(y)$, the time reflection symmetry implies

$$J_k^s(x) = \frac{v(-x)}{J_k^u(-T^k y)v(-T^k x)} \sim \frac{1}{J_k^u(-T^k y)} \frac{|2 + \kappa(\varphi)|}{|2 + \kappa(\varphi_k)|}. \quad (3.37)$$

The transversality of stable and unstable vectors, stated in Proposition 1 implies that $J_k^u(x)J_k^s(x)$ is equivalent to the k -step (Lebesgue) volume expansion factor. Using (2.2), and the T -invariance of μ , we get

$$J_k^u(x)J_k^s(x) \sim \frac{\cos \varphi}{\cos \varphi_k}. \quad (3.38)$$

Putting together (3.37) and (3.38) we get

$$J_k^u(x) \sim J_k^u(-T^k y) \frac{2 + \kappa(\varphi_k)}{2 + \kappa(\varphi)} \frac{\cos \varphi}{\cos \varphi_k} = J_k^u(-T^k y) \frac{\omega(\varphi_k)}{\omega(\varphi)}.$$

The same is true for $\bar{x} = P(\bar{y})$, so we have

$$|\ln J_k^u(x) - \ln J_k^u(\bar{x})| \leq |\ln J_k^u(-T^k y) - \ln J_k^u(-T^k \bar{y})| + \left| \ln \left| \frac{\omega(\varphi_k)}{\omega(\bar{\varphi}_k)} \right| \right| + \left| \ln \left| \frac{\omega(\varphi)}{\omega(\bar{\varphi})} \right| \right| + C.$$

To see the boundedness of the first term of the right-hand side we can apply Proposition (3), because $-T^k y$ and $-T^k \bar{y}$ are on the same local *unstable* manifold. The second and third term is bounded because W_s and $T^k W_s$ are homogeneous, see Sect. 3.2.6. Now the proof of Proposition 4 is complete. \square

4. Growth Properties of Unstable Manifolds

This last section is concerned with the growth properties of LUMs. Our aim is to show that LUMs “grow large and round, on the average”. This is expressed in the formulas of Proposition 5 below.

Recall Convention 1. Throughout the section we use the higher iterate of the dynamics, $T_1 = T^{m_0}$. This has singularity set (secondary and primary) $\Xi = \Gamma^{(m_0)}$. For the higher iterates of T_1 the singularity set is $\Xi^{(n)} = \Xi \cup T_1^{-1} \Xi \cup \dots \cup T_1^{-n+1} \Xi$.

δ_0 -LUM's. To formulate and prove further important conditions on growth of LUMs we need to recall several notions and notations from [Ch]. Let $\delta_0 > 0$. We call W a δ_0 -LUM if it is a LUM and $\text{diam } W \leq \delta_0$. For an open subset $V \subset W$ and $x \in V$ denote by $V(x)$ the connected component of V containing the point x . Let $n \geq 0$. We call an open subset $V \subset W$ a (δ_0, n) -subset if $V \cap (\Xi^{(n)}) = \emptyset$ (i.e., the map T_1^n is smooth and homogeneous on V) and $\text{diam } T_1^n V(x) \leq \delta_0$ for every $x \in V$. Note that $T_1^n V$ is then a union of δ_0 -LUM's. Define a function $r_{V,n}$ on V by

$$r_{V,n}(x) = d_{T_1^n V(x)}(T_1^n x, \partial T_1^n V(x)).$$

Note that $r_{V,n}(x)$ is the radius of the largest open ball in $T_1^n V(x)$ centered at $T_1^n x$. In particular, $r_{W,0}(x) = d_W(x, \partial W)$.

One further notation we introduce is \mathcal{U}_δ (for any $\delta > 0$), the δ -neighborhood of the closed set $\Xi \cup S_0 \cup Z_0$.

The aim of this section is to prove the proposition below.

Proposition 5. *There are constants $\alpha_0 \in (0, 1)$ and $\beta_0, D_0, \eta, \chi, \zeta > 0$ with the following property. For any sufficiently small $\delta_0, \delta > 0$ and any δ_0 -LUM W there is an open $(\delta_0, 0)$ -subset $V_\delta^0 \subset W \cap \mathcal{U}_\delta$ and an open $(\delta_0, 1)$ -subset $V_\delta^1 \subset W \setminus \mathcal{U}_\delta$ (one of these may be empty) such that $m_W(W \setminus (V_\delta^0 \cup V_\delta^1)) = 0$ and that $\forall \varepsilon > 0$,*

$$m_W(r_{V_\delta^1,1} < \varepsilon) \leq \alpha_0 \Lambda_1 \cdot m_W(r_{W,0} < \varepsilon / \Lambda_1) + \varepsilon \beta_0 \delta_0^{-1} m_W(W), \quad (4.1)$$

$$m_W(r_{V_\delta^0,0} < \varepsilon) \leq D_0 \delta^{-\eta} m_W(r_{W,0} < \varepsilon) \quad (4.2)$$

and

$$m_W(V_\delta^0) \leq D_0 m_W(r_{W,0} < \zeta \delta^\chi). \quad (4.3)$$

Proof of this Proposition goes along the lines of the arguments from [Ch]. First let us consider

Accumulation of singularity lines. There are two sources of accumulation of the components of the set Ξ that can cut LUM's into arbitrarily many pieces.

First, the set Γ_1 consists of countably many curves stretching approximately parallel to some curves in S_1 (or Z_1) and approaching them. So, each set $T^{-1}I_k$ and $k \neq 0$, is a narrow strip with curvilinear boundaries. The expansion of unstable fibers in these strips can be estimated using (3.33), (3.34) and (3.30). More precisely, let $W \subset T^{-1}I_k$ be a LUM, for some $k \neq 0$. Then the expansion factor, $J^u(x)$, on W satisfies

$$J^u(x) \sim \omega(\varphi) \sim k^2 \quad \forall x \in W. \quad (4.4)$$

Second, there might be multiple intersections of the curves in $S_1 \cup Z_1$. Recall K_n , the complexity of $S^{(n)} \cup Z^{(n)}$ and its properties from Subsect. 2.1. Specifically important for us is the choice of the higher iterate $T_1 = T^{m_0}$ with its relevant properties, see Convention 1.

Indexing system. Before proving the proposition we introduce a handy indexing system, cf. [Ch]. Let $\delta_0 > 0$ and W be a δ_0 -LUM. If δ_0 is small enough, then W crosses at most K_{m_0} curves of the set $S^{(m_0)} \cup Z^{(m_0)}$, so the set $W \setminus (S^{(m_0)} \cup Z^{(m_0)})$ consists of at most $K_{m_0} + 1$ connected curves, let us call them W_1, \dots, W_p with $p \leq K_{m_0} + 1$.

On each W_j the map T_1 (as a map on M) is smooth, but any W_j may be cut into arbitrary many (countably many) pieces by other curves in Ξ , which are the preimages of the boundaries of I_k . Let $\Delta \subset W$ be a connected component of the set $W \setminus \Xi$. It can be identified with the $(m_0 + 1)$ -tuple $(k_1, \dots, k_{m_0}; j)$ such that $\Delta \subset W_j$ and $T^i \Delta \subset I_{k_i}$ for $1 \leq i \leq m_0$. Note that this identification is almost unique. Indeed, given j , $(T^i \Delta \subset) T^i W_j$ is contained in a strip of the phase space that lies between two horizontal lines: two components of $S_0 \cup Z_0$. It might happen that expansion factors diverge – and consequently, homogeneity strips have been constructed – at both sides of the strip. Thus given the index k_i , we have $T^i \Delta \subset I_{k_i}$, where I_{k_i} can be the k_i^{th} layer from one of the two homogeneity structures. In such a case we use the following convention; the homogeneity layers at the “upper” and “lower” ends of the phase space strip (corresponding to j) are labelled by odd and even numbers, respectively. This way the indexing system is made unique and (4.4) remains true.

All in all, we will write $\Delta = \Delta(k_1, \dots, k_{m_0}; j)$. Of course, some strings $(k_1, \dots, k_{m_0}; j)$ may not correspond to any piece of W , for such strings $\Delta(k_1, \dots, k_{m_0}; j) = \emptyset$.

Denote by $J_1^u(x) = J^u(x) \cdots J^u(T^{m_0-1}x)$ the expansion factor of unstable vectors under DT_1 . Let $|\Delta| = m_\Delta(\Delta)$ be the Euclidean length of a LUM Δ . We record two important facts:

(a) For every point $x \in \Delta(k_1, \dots, k_{m_0}; j)$ we have

$$J_1^u(x) \geq L_{k_1, \dots, k_{m_0}} := \max \left\{ \Lambda_1, C_{20} \prod_{k_i \neq 0} k_i^2 \right\},$$

where C_{20} is some positive global constant. This follows from (4.4).

(b) For each $\Delta(k_1, \dots, k_{m_0}; j)$ we have

$$|\Delta(k_1, \dots, k_{m_0}; j)| \leq M_{k_1, \dots, k_{m_0}} := \min \left\{ |W|, C_{21} \prod_{k_i \neq 0} k_i^{-2} \right\},$$

where $C_{21} = C_{20}^{-1} |W|_{\max}$ and $|W|_{\max}$ is the maximal length of LUMs in M . This follows from the previous fact.

Next, put

$$\theta_0 := 2 \sum_{k=k_0}^{\infty} k^{-2} \leq 4/k_0$$

and let us turn to the proof of our growth formulas.

Let W be a δ_0 -LUM and $\delta > 0$ be small. For each connected component $\Delta \subset W \setminus \Xi$ put $\Delta^0 = \Delta \cap \mathcal{U}_\delta$ and $\Delta^1 = \text{int}(\Delta \setminus \mathcal{U}_\delta)$ (recall \mathcal{U}_δ is the δ -neighborhood of $\Xi \cup S_0 \cup Z_0$). Due to the Continuation Property (cf. Subsect. 2.1) and to Alignment (cf. Subsubsect. 3.2.5), the set Δ^0 consists of two subintervals adjacent to the endpoints of Δ (they may overlap and cover Δ , of course). The set Δ^1 is either empty or a subinterval of Δ . We put $W^1 = \cup_{\Delta \subset W \setminus \Xi} \Delta^1$.

Proof of (4.1). For each Δ^1 the set $T_1(\Delta^1 \cap \{r_{W^1,1} < \varepsilon\})$ is the union of two subintervals of $T_1\Delta^1$ of length ε adjacent to the endpoint of $T_1\Delta^1$. Using the above indexing system we get

$$\begin{aligned} m_W(r_{W^1,1} < \varepsilon) &\leq \sum_{k_1, \dots, k_{m_0}, j} 2\varepsilon L_{k_1, \dots, k_{m_0}}^{-1} \\ &\leq 2\varepsilon p \left[\Lambda_1^{-1} + C_{20}^{-1}(\theta_0 + \theta_0^2 + \dots + \theta_0^{m_0}) \right] \\ &\leq 2\varepsilon (K_{m_0} + 1) \left(\Lambda_1^{-1} + C_{20}^{-1} m_0 \theta_0 \right). \end{aligned}$$

We now assume that k_0 is large enough so that

$$\alpha_0 := (K_{m_0} + 1)(\Lambda_1^{-1} + C_{20}^{-1} m_0 \theta_0) < 1$$

and thus get

$$m_W(r_{W^1,1} < \varepsilon) \leq \min\{|W|, 2\alpha_0\varepsilon\}.$$

The first term on the right-hand side of (4.1) is equal to

$$\alpha_0 \Lambda_1 \min\{|W|, 2\varepsilon/\Lambda_1\} = \min\{\alpha_0 \Lambda_1 |W|, 2\alpha_0\varepsilon\}.$$

Since $\alpha_0 \Lambda_1 > 1$, we get

$$m_W(r_{W^1,1} < \varepsilon) \leq \alpha_0 \Lambda_1 \cdot m_W(r_{W,0} < \varepsilon/\Lambda_1). \quad \square \quad (4.5)$$

Next, to obtain an open $(\delta_0, 1)$ -subset V_δ^1 of W^1 , one needs to further subdivide the intervals $\Delta^1 \subset W$ such that $|T_1\Delta^1| > \delta_0$. Each such LUM $T_1\Delta^1$ we divide into s_Δ equal subintervals of length $\leq \delta_0$, with $s_\Delta \leq |T_1\Delta^1|/\delta_0$. If $|T_1\Delta^1| < \delta_0$, then we set $s_\Delta = 0$ and leave Δ^1 unchanged. Then the union of the preimages under T_1 of the above intervals will make V_δ^1 . Now we must estimate the measure of the ε -neighborhood of the additional endpoints of the subintervals of $T_1\Delta^1$. This gives

$$\begin{aligned} m_W(r_{V_\delta^1,1} < \varepsilon) - m_W(r_{W^1,1} < \varepsilon) &\leq \sum_{\Delta \subset W \setminus \Xi} 2s_\Delta \varepsilon |C_{22}\Delta^1|/|T_1\Delta^1| \\ &\leq \sum_{\Delta \subset W \setminus \Xi} 2C_{22}\varepsilon |\Delta^1|/\delta_0 \\ &\leq 2C_{22}\varepsilon \delta_0^{-1} |W|. \end{aligned}$$

Here $C_{22} = \exp(\text{const} \cdot |W|_{\max}^{\frac{1}{5}})$ is an upper bound on distortions on LUM's, see Proposition 3. Combining the above bound with (4.5) completes the proof of (4.1) with $\beta_0 = 2C_{22}$.

We now *prove (4.2)*. It is enough to consider $\varepsilon \leq |W|/2$, so that the right-hand side of (4.2) equals $2D_0\delta^{-\eta}\varepsilon$. We can put $V_\delta^0 = W \setminus V_\delta^1$. Then the left-hand side of (4.2) does not exceed $2J_\delta\varepsilon$, where J_δ is the number of nonempty connected components of the set V_δ^0 , which is at most the number of connected components of $W \setminus \Xi$ of length $> 2\delta$. Hence, clearly $J_\delta \leq |W|/\delta \leq \delta_0/\delta$. This proves (4.2) with $\eta = 1$.

Finally, we *prove the inequality (4.3)*. Again, let Δ be a connected component of $W \setminus \Xi$ and Δ^0, Δ^1 be defined as above, with the set Δ^0 consisting of two subintervals adjacent to the endpoints of Δ . By (3.28) – and the analogous property for the secondary

singularities, see Subsubsect. 3.2.5 and 3.2.6 – each of these subintervals has length smaller than $C\delta^\alpha$.

Now, the right-hand side of (4.3) equals $D_0 \min\{|W|, 2\zeta\delta^\chi\}$. So, it is enough to show that $m_W(V_\delta^0) \leq B\delta^\chi$ for some $B, \chi > 0$. We have

$$\begin{aligned} m_W(V_\delta^0) &\leq \sum_{\Delta \subset W \setminus \Xi} \min\{2C\delta^\alpha, |\Delta|\} \\ &\leq \sum_{k_1, \dots, k_{m_0}, j} \min\{2C\delta^\alpha, M_{k_1, \dots, k_{m_0}}\} \\ &\leq \text{const} \cdot \delta^\alpha + \text{const} \cdot \sum_{k_1, \dots, k_{m_0}}^* \min \left\{ \delta^\alpha, \prod_{k_i \neq 0} k_i^{-2} \right\}, \end{aligned}$$

where \sum^* is taken over m_0 -tuples that contain at least one nonzero index $k_i \neq 0$. The following lemma – Lemma 7.2 from [Ch], which was proved in the Appendix of that paper – completes the proof of (4.3) with $\chi = \frac{\alpha}{2m_0}$.

Lemma 5. *Let $\epsilon > 0$ and $m \geq 1$. Then*

$$\sum_{k_1, \dots, k_m \geq 2} \min \left\{ \epsilon, (k_1 \cdots k_m)^{-2} \right\} \leq B(m) \cdot \epsilon^{1/2m}.$$

With the help of this lemma Proposition 5, and consequently, Theorem 1 is proved. \square

5. Specific Potentials

In this section we would like to show that, as important corollaries of Theorem 1, exponential decay of correlations can be established for certain specific potentials. To prove such corollaries we need to calculate the rotation function $\Delta\Theta(\varphi)$ from the potential $V(r)$.

As to the detailed description of the Hamiltonian flow in a circularly symmetric potential, we refer to the literature, e.g. [DL] and references therein. Most important is that besides the full energy there is an additional integral of motion, the angular momentum l , that can be calculated for a specific trajectory as

$$l = R \sin \varphi,$$

where φ is the collision angle at income. For brevity of notation it is worth introducing the function

$$h(r) = (1 - 2V(r))r^2.$$

By the presence of the angular momentum motion is completely integrable and is described by the pair of differential equations (recall our convention that the full energy is $E = \frac{1}{2}$):

$$\begin{aligned} \dot{r}^2 &= r^{-2}(h(r) - l^2), \\ r^2 \dot{\Theta} &= l. \end{aligned}$$

Combining these we get

$$\frac{d\Theta}{dr} = \pm \frac{l}{r\sqrt{h(r) - l^2}}, \quad (5.1)$$

where the sign depends on whether r is increasing or decreasing. More precisely, there is a minimum radius

$$\hat{r} = \hat{r}(\varphi) : \quad h(\hat{r}) = l^2 = R^2 \sin^2 \varphi,$$

down to which r decreases (with negative sign in (5.1)) and from which r increases (with positive sign in (5.1)). This results in

$$\Delta\Theta(\varphi) = 2 \int_{\hat{r}}^R \frac{l}{r\sqrt{h(r) - l^2}} dr. \quad (5.2)$$

For a generic potential, the dependence of (5.2) on φ is rather implicit: φ is present both in the integrand (via l) and in the limits (via \hat{r}). One possible strategy to follow is to obtain some even more complicated formulas for the derivatives in the general case, and based on those perform estimates that guarantee the desired dynamical properties. This is possible as long as only hyperbolicity and ergodicity is treated – like in [DL] – and thus only the first derivative, $\kappa(\varphi) = \Delta\Theta'(\varphi)$ is needed. However, for rate of mixing you need one more derivative, $\kappa'(\varphi) = \Delta\Theta''(\varphi)$, cf. Definition 3. Finding good sufficient conditions on the potential $V(r)$ that guarantee the regularity of κ seems to be a very hard task, if possible at all. Thus we have chosen instead to investigate some specific cases where $\Delta\Theta$ is directly computable from (5.2). Of course, this way we could handle a much narrower class of potentials than [DL], nevertheless, the established dynamical property is stronger.

Corollary 1. *Consider the case of a constant potential,*

$$V(r) = V_0 \quad \text{for any } r \in [0, R).$$

Correlations decay with an exponential rate in case

- $V_0 > 0$ and the configuration is arbitrary,
- $V_0 < 0$ and the configuration is such that $\tau_{\min} > \frac{2R}{\sqrt{1-2V_0-1}}$.

Remarks. Actually, the analysis of this constant potential case from the point of ergodicity dates back to the late eighties, to [Kn2] and [Ba]. Rate of mixing is, to our knowledge, discussed for the first time. For potential values $V_0 > \frac{1}{2}$ the particle cannot enter the disks, the system is equivalent to the traditional dispersing billiard, thus we consider the opposite case, $V_0 < \frac{1}{2}$.

Proof. Let us introduce the quantity

$$v = \sqrt{1 - 2V_0} \quad (5.3)$$

which is less or greater than 1 depending on the sign of V_0 . Let us consider the case of positive V_0 first and introduce furthermore the angle φ_0 for which:

$$v = \sin \varphi_0.$$

In case $|\varphi| > \varphi_0$, $|l|$ is greater than the maximum value $h(r)$ can take, which indicates that the particle has too large angular momentum to enter the potential, thus $\Delta\Theta = 0$. In

the opposite case of $|\varphi| < \varphi_0$ it is easy to obtain $\hat{r} = \frac{R|\sin \varphi|}{\nu}$ and perform the integration of (5.2). All in all

$$\Delta\Theta(\varphi) = \begin{cases} 2 \arccos\left(\frac{\sin \varphi}{\nu}\right) & \text{if } |\varphi| < \varphi_0, \\ 0 & \text{if } |\varphi| > \varphi_0. \end{cases}$$

On the one hand, whatever configuration we have, the system satisfies property H (cf. Definition 2), as either $\kappa = 0$ or $\kappa \leq \frac{-2}{\nu} < -2$. On the other hand, κ is a piecewise C^1 function of φ and it behaves as $(\varphi_0 - \varphi)^{-\frac{1}{2}}$ near the discontinuity point φ_0 . Thus κ is regular (cf. Definition 3 and the remarks following it). This means that the first statement of our Corollary follows from Theorem 1.

Now let us turn to the case of $V_0 < 0$ (i.e. $\nu > 1$). It is even simpler to calculate the rotation function (5.2):

$$\Delta\Theta(\varphi) = 2 \arccos\left(\frac{\sin \varphi}{\nu}\right)$$

for all φ . As $\nu > 1$, this is a C^2 function on the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$, thus κ is definitely regular. As to property H, we have $0 > \kappa \geq -\frac{2}{\nu}$, where the minimum is obtained at $\varphi = 0$. Thus the assumption on the configuration from Definition 2 reads as $\tau_{\min} > \frac{2R}{\nu-1}$ and the second statement of the corollary follows from Theorem 1. \square

Remark. Note that motion in the constant potential is equivalent to the problem of diffraction from geometric optics. More precisely, we can think of the disks as if they were made of a material optically different from their neighborhood, where the relative diffraction coefficient is ν from (5.3). In case the disks are optically less dense than their neighborhood (i.e. $\nu < 1$, $V_0 > 0$), we may observe the phenomenon of complete reflection that corresponds to the limiting angle φ_0 .

Corollary 2. *Given constants $A > 0$ and $\beta > -2$, consider the potential*

$$V(r) = A \left(1 - \left(\frac{r}{R} \right)^\beta \right).$$

Correlations decay at an exponential rate in case:




Fig. 2. Rotation function for two examples

- $A = \frac{1}{2}$, $0 > \beta (> -2)$ and the configuration is arbitrary,
- $A = \frac{1}{2}$, $\beta > 0$ and the configuration is such that $\tau_{\min} > \frac{2R}{\beta}$.

Remark. Note that according to our construction the chosen value for the constant A , $A = \frac{1}{2}$ is exactly the full energy. If we had a different value for A , the integration in (5.2) would be much more complicated. In other words, Corollary 2, in contrast to Corollary 1 is unstable with respect to variations of the full energy (see also the discussion below, following the proof). Nevertheless it is nice to have at least one potential with exponential mixing for any kind of power law behaviour (if $\beta \leq -2$, a positive measure set of trajectories is pulled into the center of the disk, cf. [DL]).

Proof. By straightforward calculation

$$h(r) = \frac{r^{2+\beta}}{R^\beta}; \quad \text{and} \quad \hat{r} = R |\sin \varphi|^{\frac{2}{2+\beta}}.$$

Then it is not hard to integrate in (5.2):

$$\Delta\Theta(\varphi) = \frac{4}{2+\beta} \left(\frac{\pi}{2} - \varphi \right)$$

for all $\varphi \neq 0$. Thus $\Delta\Theta$ is piecewise linear (in the general case with one discontinuity of the first kind at $\varphi = 0$) and thus

$$\kappa = -\frac{4}{2+\beta}$$

identically. Regularity (in terms of Definition 3) is automatic.

Let us consider the attracting potentials, $\beta < 0$ first. In such a case the potential has a singularity at the center of the disk, resulting in the discontinuity at $\varphi = 0$ ². Nevertheless, $\kappa < -2$, thus property H (cf. Definition 2) and consequently the first statement of the corollary follows.

Now if $\beta > 0$, as $A = \frac{1}{2}$, the “top” of the potential is equal to the energy. As a consequence, for the initial value $\varphi = 0$ the flow is not uniquely defined, resulting in the discontinuity for the rotation function. However, in accordance with Definition 2, property H is satisfied if $\tau_{\min} > \frac{2R}{\beta}$. Thus the second statement of the corollary holds. \square

Discussion. As already mentioned, Corollary 2 is very sensitive to the convention $E = \frac{1}{2}$. Though very difficult to calculate, it is interesting to guess what happens if one perturbs the constant A (or equivalently, the full energy level).

Let us consider the case $\beta > 0$ first. With A either increased or decreased from the value $\frac{1}{2}$, the physical reason for the discontinuity at $\varphi = 0$ disappears and we expect smooth rotation functions. By continuity of the potential at R , $\Delta\Theta(\frac{\pi}{2}) = 0$ seems also reasonable. As to the initial value $\varphi = 0$ let us have a look at the case $A < \frac{1}{2}$ first. There is no reason for the trajectory to deviate in direction: it slows down, reaches the center and then speeds up following a linear track. Thus $\Delta\Theta(0) = \pi$. This altogether implies on the basis of Lagrange’s mean value theorem that there definitely exists at least one $\varphi \in (0, \frac{\pi}{2})$ for which $\kappa(\varphi) = -2$. In such a case, however, stable periodic orbits tend to

² However, in case $\beta = -2(1 - \frac{1}{n})$, the left and right limits coincide; this corresponds to the possibility of regularizing the flow, cf. [DL] and [Kn1].

appear and the system is most likely not even ergodic, cf. [Do1]. One can suspect that a typical repelling potential which has a maximum less than the total energy leads to non-ergodic soft billiards in a similar fashion.

In the opposite case of $A > \frac{1}{2}$ the behaviour of trajectories in the vicinity of $\varphi = 0$ is completely different. As the top of the potential is higher than the full energy, the particle cannot “climb” it thus it should “turn back”. We expect $\Delta\Theta(0) = 0$ and a smooth rotation function with $\kappa > -2$ for all φ . That would mean ergodicity and possibly exponential mixing in case of a suitable configuration (cf. Definition 2). All in all, ergodic and statistical behaviour is very sensitive to perturbation of the full energy level.

In the case of $\beta < 0$ it is not so easy to guess. Nevertheless, we can say something rather surprising in one particular case that indicates similar sensitivity. Choose $\beta = -1$ and $A = 1$. It is not difficult to obtain $h(r) = 2r - r^2$. The integral in (5.2) is a bit more complicated now, nevertheless, it is possible to evaluate:

$$\Delta\Theta(\varphi) = 2\pi - 2\varphi \tag{5.4}$$

which means $\kappa = -2$ identically. This corresponds to the least ergodic behaviour we can have. It is straightforward to obtain that an identically zero potential ($V(r) = 0$ for all r) would result in $\Delta\Theta(\varphi) = \pi - 2\varphi$. Thus by (5.4) in this particular case of $A = 1$, $\beta = -1$ trajectories evolve as if they passed on freely and were reflected when leaving the disc.

Thus if $\beta = -1$, we may have exponential mixing ($A = \frac{1}{2}$) and stability ($A = 1$). As to other values of A it is worth mentioning that ergodicity follows from [DL] in case $A < \frac{1}{2}$.

6. Outlook

In this last section we list several possible interesting directions of future research.

1. As to the possibly most direct challenge, we conjecture that there exist rapidly mixing potentials for which the condition $|\kappa + 2| > c$ (i.e. property H from Definition 2) is not satisfied for nearly tangential trajectories. Thus these systems are not covered by Theorem 1, even more, at least to our knowledge, there is no result in the literature on the ergodicity or hyperbolicity of such soft billiards either. Thus we make the following

Remark. Note that it is possible that κ tends to -2 as $\varphi \rightarrow \frac{\pi}{2}$, nevertheless, $|\frac{2+\kappa}{\cos\varphi}| > c$ and the system can be hyperbolic (possibly ergodic or exponentially mixing). We will turn back to this question in a separate paper.

The difficulty with the treatment of this case is, as already mentioned in Sect. 1, that the separate investigation of motion inside and outside the disks seems not to work at several arguments.

2. Further exciting open questions seem even more difficult. One natural direction of generalization is of course the higher dimensional case. As to softening of multi-dimensional dispersing billiards (motivated e.g. by the three dimensional Lorentz process with spherical scatterers) we are not aware of any mathematical result. Even hyperbolicity and ergodicity seem difficult, not to mention decay of correlations, especially in view of the recently observed pathological behaviour of singularity manifolds in multi-dimensional billiards (see [BChSzT]).

3. Another direction of future research, motivated mainly by applications to physics, could be the further investigation of those systems for which rapid mixing is already established. For example, as mathematical evidence on the existence of diffusion and other transport coefficients is given, it would be interesting to understand the dependence of these on certain parameters like the full energy level.
4. Last but not least, in contrast to the generality of Theorem 1, it is striking how narrow the class of specific potentials is for which we could apply the result in Sect. 5. It would be desirable to establish – at least numerically – our reasonable regularity properties for as wide a class of potentials as possible.

Acknowledgements. We are very grateful to our advisor, Domokos Szász for suggesting to this exciting topic and for lots of useful conversations. The financial support of the Hungarian National Foundation for Scientific Research (OTKA), grants T26176, T32022 and TS040719; of the Research Group Stochastics at TUB and the Research Project 2001-122 of the Hungarian Academy of Sciences, both affiliated to the Technical University of Budapest, is also acknowledged.

Appendix

Here we provide, for the reader's convenience, a very short, yet mainly self-contained formulation of Theorem 2.1 from [Ch]. For self-containedness, many notions and notations are repeatedly introduced. First we give the conditions **P0**...**P6** which are required, and then the statement of the theorem.

P0. The dynamical system is a map $T : M \setminus \Gamma \rightarrow M$, where M is an open subset in a C^∞ Riemannian manifold, \bar{M} is compact. Γ is a closed subset in \bar{M} , and T is a C^2 diffeomorphism of its range onto its image. Γ is called the singularity set.

P1. Hyperbolicity. We assume there are two families of cone fields C_x^u and C_x^s in the tangent planes $T_x M$, $x \in \bar{M}$ and there exists a constant $\Lambda > 1$ with the following properties:

- $DT(C_x^u) \subset C_{Tx}^u$ and $DT(C_x^s) \supset C_{Tx}^s$ whenever DT exists;
- $|DT(v)| \geq \Lambda|v| \quad \forall v \in C_x^u$;
- $|DT^{-1}(v)| \geq \Lambda|v| \quad \forall v \in C_x^s$;
- these families of cones are continuous on \bar{M} , their axes have the same dimensions across the entire \bar{M} which we denote by d_u and d_s , respectively;
- $d_u + d_s = \dim M$;
- the angles between C_x^u and C_x^s are uniformly bounded away from zero:

$\exists \alpha > 0$ such that $\forall x \in M$ and for any $dw_1 \in C_x^u$ and $dw_2 \in C_x^s$ one has

$$\angle(dw_1, dw_2) \geq \alpha$$

The C_x^u are called the unstable cones whereas C_x^s are called the stable ones.

The property that the angle between stable and unstable cones is uniformly bounded away from zero is called **transversality**.

Some notation and definitions. For any $\delta > 0$ denote by \mathcal{U}_δ the δ -neighborhood of the closed set $\Gamma \cup \partial M$. We denote by ρ the Riemannian metric in M and by m the Lebesgue measure (volume) in M . For any submanifold $W \subset M$ we denote by ρ_W the metric on W induced by the Riemannian metric in M , by m_W the Lebesgue measure on W generated by ρ_W , and by $\text{diam}W$ the diameter of W in the ρ_W metric.

LUM-s. To be able to formulate the further properties to be checked the reader is kindly reminded of the notion of local unstable manifolds. We call a ball-like submanifold $W^u \subset M$ a local unstable manifold (LUM) if $-\dim W^u = d_u$, $-T^{-n}$ is defined and smooth on W^u for all $n \geq 0$, $\forall x, y \in W^u$ we have $\rho(T^{-n}x, T^{-n}y) \rightarrow 0$ exponentially fast as $n \rightarrow \infty$.

We denote by $W^u(x)$ (or just $W(x)$) a local unstable manifold containing x . Similarly, local stable manifolds (LSM) are defined.

P2. SRB measure. *The dynamics T has to have an invariant ergodic Sinai-Ruelle-Bowen (SRB) measure μ . That is, there should be an ergodic probability measure μ on M such that for μ -a.e. $x \in M$ a LUM $W(x)$ exists, and the conditional measure on $W(x)$ induced by μ is absolutely continuous with respect to $m_{W(x)}$.*

Furthermore, the SRB-measure should have nice mixing properties: the system (T^n, μ) is ergodic for all finite $n \geq 0$.

In our case the SRB measure is simply the Liouville-measure defined by (2.2) in Sect. 2. Absolute continuity of μ is straightforward, while the other above required properties (invariance, ergodicity, mixing) are proved in [DL].

P3. Bounded curvature. *The tangent plane of an unstable manifold should be a Lipschitz function of the phase point. By this we mean that a base can be chosen in every tangent plane so that every base vector is a Lipschitz function of the phase point.*

Some notation. Denote by $J^u(x) = |\det(DT|_{E_x^u})|$ the Jacobian of the map T restricted to $W(x)$ at x , i.e. the factor of the volume expansion on the LUM $W(x)$ at the point x .

P4. Distortion bounds. *Let x, y be in one connected component of $W \setminus \Gamma^{(n-1)}$, which we denote by V . Then*

$$\log \prod_{i=0}^{n-1} \frac{J^u(T^i x)}{J^u(T^i y)} \leq \varphi(\rho_{T^n V}(T^n x, T^n y)),$$

where $\varphi(\cdot)$ is some function, independent of W , such that $\varphi(s) \rightarrow 0$ as $s \rightarrow 0$.

P5. Absolute continuity. *Let W_1, W_2 be two sufficiently small LUM-s, such that any LSM W^s intersects each of W_1 and W_2 in at most one point. Let $W'_1 = \{x \in W_1 : W^s(x) \cap W_2 \neq \emptyset\}$. Then we define a map $h : W'_1 \rightarrow W_2$ by sliding along stable manifolds. This map is often called a holonomy map. This has to be absolutely continuous with respect to the Lebesgue measures m_{W_1} and m_{W_2} , and its Jacobian (at any density point of W'_1) should be bounded, i.e.*

$$1/C' \leq \frac{m_{W_2}(h(W'_1))}{m_{W_1}(W'_1)} \leq C'$$

with some $C' = C'(T) > 0$.

A few words are in order to discuss how our Proposition 4 implies property (P5). Let us consider the unique ergodic SRB-measure μ for the dynamical system (in our billiard dynamics this is precisely the Liouville measure defined by (2.2)). We know that the conditional measure on any LUM induced by μ is absolutely continuous with respect to the Lebesgue measure on the unstable manifold. These conditional measures are often

referred to as u-SRB measures and their density w.r.t. the Lebesgue measure, $\rho_W(x)$ is given by the following equation (cf. [Ch]):

$$\frac{\rho_W(x)}{\rho_W(y)} = \lim_{n \rightarrow \infty} \prod_{i=1}^n \frac{J^u(T^{-i}x)}{J^u(T^{-i}y)}.$$

Actually, what directly follows from Proposition 4 is that if we consider two nearby LUM-s W and \bar{W} and points x, \bar{x} on them joint by the holonomy map along an s-manifold, then the ratio of $\rho_W(x)$ and $\rho_{\bar{W}}(\bar{x})$, the densities for the two u-SRB measures is uniformly bounded. However, taking into account the invariance of μ and the uniform contraction along s-manifolds, we may get the uniform bound on the distortion of Lebesgue measures, i.e. the property we assumed in (P5).

Some further notation. Let $\delta_0 > 0$. We call W a δ_0 -LUM if it is a LUM and $\text{diam } W \leq \delta_0$. For an open subset $V \subset W$ and $x \in V$ denote by $V(x)$ the connected component of V containing the point x . Let $n \geq 0$. We call an open subset $V \subset W$ a (δ_0, n) -subset if $V \cap \Gamma^{(n)} = \emptyset$ (i.e., the map T^n is smoothly defined on V) and $\text{diam } T^n V(x) \leq \delta_0$ for every $x \in V$. Note that $T^n V$ is then a union of δ_0 -LUM-s. Define a function $r_{V,n}$ on V by

$$r_{V,n}(x) = \rho_{T^n V(x)}(T^n x, \partial T^n V(x)).$$

Note that $r_{V,n}(x)$ is the radius of the largest open ball in $T^n V(x)$ centered at $T^n x$. In particular, $r_{W,0}(x) = \rho_W(x, \partial W)$.

Now we are able to give the last group of technical properties that have to be verified:

P6. Growth of unstable manifolds *Let us assume there is a fixed $\delta_0 > 0$. Furthermore, there exist constants $\alpha_0 \in (0, 1)$ and $\beta_0, D_0, \kappa, \sigma, \zeta > 0$ with the following property. For any sufficiently small $\delta > 0$ and any δ_0 -LUM W there is an open $(\delta_0, 0)$ -subset $V_\delta^0 \subset W \cap \mathcal{U}_\delta$ and an open $(\delta_0, 1)$ -subset $V_\delta^1 \subset W \setminus \mathcal{U}_\delta$ (one of these may be empty) such that the two sets are disjoint, $m_W(W \setminus (V_\delta^0 \cup V_\delta^1)) = 0$ and $\forall \varepsilon > 0$,*

$$\begin{aligned} m_W(r_{V_\delta^1,1} < \varepsilon) &\leq \alpha_0 \Lambda \cdot m_W(r_{W,0} < \varepsilon/\Lambda) + \varepsilon \beta_0 \delta_0^{-1} m_W(W), \\ m_W(r_{V_\delta^0,0} < \varepsilon) &\leq D_0 \delta^{-\kappa} m_W(r_{W,0} < \varepsilon), \end{aligned}$$

and

$$m_W(V_\delta^0) \leq D_0 m_W(r_{W,0} < \zeta \delta^\sigma).$$

Now we can formulate Theorem 2.1 from [Ch].

Theorem A.1 (Chernov, 1999). *Under the conditions P0 . . . P6, the dynamical system enjoys exponential decay of correlations and the central limit theorem for Hölder-continuous functions.*

The properties stated in the theorem are defined in Definitions 4 and 5.

References

- [Ba] Baldwin, P.R.: Soft billiard systems. *Physica D* **29**, 321–342 (1988)
- [BChSzT] Bálint, P., Chernov, N.I., Szász, D., Tóth, I.P.: Geometry of Multi-dimensional Dispersing Billiards. To appear in *Asterisque*
- [BáTó] Bálint, P., Tóth, I.P.: Mixing and its rate in “soft” and “hard” billiards motivated by the Lorentz process. To appear in *Physica D*
- [BöTa] Böröczky, K., Jr., Tardos, G.: The longest segment in the complement of a packing. *Matematika*, to appear
- [Ch] Chernov, N.: Decay of correlations and dispersing billiards. *J. Statist. Phys.* **94**, 513–556 (1999)
- [ChY] Chernov, N., Young, L.S.: Decay of Correlations for Lorentz gases and hard balls. In: *Hard ball systems and the Lorentz gas*, *Encyclopedia Math. Sci.* **101**, Szász, D. (ed), Berlin: Springer, 2000, pp. 89–120
- [Do1] Donnay, V.: Non-ergodicity of two particles interacting via a smooth potential. *J. Statist. Phys.* **96**(5–6), 1021–1048 (1999)
- [Do2] Donnay, V.: Elliptic islands in generalized Simai billiards. *Ergod. Th. and Dynam. Sys.* **16**(5), 975–1010 (1997)
- [DL] Donnay, V., Liverani, C.: Potentials on the two-torus for which the Hamiltonian flow is ergodic. *Commun. Math. Phys.* **135**, 267–302 (1991)
- [Kn1] Knauf, A.: Ergodic and topological properties of Coulombic periodic potentials. *Commun. Math. Phys.* **110**, 89–112 (1987)
- [Kn2] Knauf, A.: On soft billiard systems. *Physica D* **36**, 259–262 (1989)
- [Ku] Kubo, I.: Perturbed billiard systems, I. *Nagoya Math. J.* **61**, 1–57 (1976)
- [KM] Kubo, I., Murata, H.: Perturbed billiard systems II, Bernoulli properties. *Nagoya Math. J.* **81**, 1–25 (1981)
- [M] Markarian, R.: Ergodic properties of plane billiards with symmetric potentials. *Commun. Math. Phys.* **145**, 435–446 (1992)
- [RT] Rom-Kedar, V., Turaev, D.: Big islands in dispersing billiard-like potentials. *Physica D* **130**(3–4), 187–210 (1999)
- [Si1] Sinai, Ya.G.: On the foundations of the ergodic hypothesis for a dynamical system of statistical mechanics. *Dokl. Akad. Nauk SSSR* **153**, 1262–1264 (1963)
- [Si2] Sinai, Ya.G.: Dynamical systems with elastic reflections. Ergodic properties of dispersing billiards. *Russ. Math. Surv.* **25**, 137–189 (1970)
- [SCh] Sinai, Ya.G., Chernov, N.: Ergodic Properties of Certain Systems of 2–D Discs and 3–D Balls. *Russ. Math. Surv.* **42**(3), 181–201 (1987)
- [Y] Young, L.S.: Statistical properties of dynamical systems with some hyperbolicity. *Ann. Math.* **147**, 585–650 (1998)

Communicated by G. Gallavotti