

Probability 1
CEU Budapest, fall semester 2017
Imre Péter Tóth

Homework sheet 3 – due on 24.10.2017 – and exercises for practice

3.1 (**homework**) For real numbers a_1, a_2, a_3, \dots define the infinite product $\prod_{k=1}^{\infty} a_k$ as

$$\prod_{k=1}^{\infty} a_k := \lim_{n \rightarrow \infty} \prod_{k=1}^n a_k,$$

whenever this limit exists.

Let p_1, p_2, p_3, \dots satisfy $0 \leq p_k < 1$ for all k . Show that $\prod_{k=1}^{\infty} (1-p_k) > 0$ if and only if $\sum_{k=1}^{\infty} p_k < \infty$.

(Hint: estimate the logarithm of $(1-p)$ with p .)

3.2 Let X_1, X_2, \dots be independent random variables such that

$$\mathbb{P}(X_n = n^2 - 1) = \frac{1}{n^2}, \quad \mathbb{P}(X_n = -1) = 1 - \frac{1}{n^2}.$$

Show that $\mathbb{E}X_n = 0$ for every n , but

$$\lim_{n \rightarrow \infty} \frac{X_1 + \dots + X_n}{n} = -1$$

almost surely.

3.3 (**homework**) Let X_1, X_2, \dots, X_n be i.i.d. random variables. Prove that the following two statements are equivalent:

- (i) $\mathbb{E}|X_i| < \infty$.
- (ii) $\mathbb{P}(|X_n| > n \text{ for infinitely many } n\text{-s}) = 0$.

Hint: If Y is nonnegative integer valued, then $\mathbb{E}Y = \sum_{k=0}^{\infty} k\mathbb{P}(Y = k) = \sum_{n=1}^{\infty} \mathbb{P}(Y \geq n)$. (Why?)

3.4 Prove that for *any* sequence X_1, X_2, \dots of random variables (real valued, defined on the same probability space) there exists a sequence c_1, c_2, \dots of numbers such that

$$\frac{X_n}{c_n} \rightarrow 0 \text{ almost surely.}$$

3.5 Let the random variables $X_1, X_2, \dots, X_n, \dots$ and X be defined on the same probability space. Prove that the following two statements are equivalent:

- (i) $X_n \rightarrow X$ in probability as $n \rightarrow \infty$.
- (ii) From every subsequence $\{n_k\}_{k=1}^{\infty}$ a sub-subsequence $\{n_{k_j}\}_{j=1}^{\infty}$ can be chosen such that $X_{n_{k_j}} \rightarrow X$ almost surely as $j \rightarrow \infty$.

3.6 (**homework**) Let X_1, X_2, \dots be independent such that X_n has *Bernoulli*(p_n) distribution. Determine what property the sequence p_n has to satisfy so that

(a) $X_n \rightarrow 0$ in probability as $n \rightarrow \infty$

(b) $X_n \rightarrow 0$ almost surely as $n \rightarrow \infty$.

3.7 Let X_1, X_2, \dots be independent random variables. Show that $\mathbb{P}(\sup_n X_n < \infty) = 1$ if and only if there is some $A \in \mathbb{R}$ for which $\sum_{n=1}^{\infty} \mathbb{P}(X_n > A) < \infty$.

3.8 Let X_1, X_2, \dots be independent exponentially distributed random variables such that X_n has parameter λ_n . Let $S_n := \sum_{i=1}^n X_i$. Show that if $\sum_{n=1}^{\infty} \frac{1}{\lambda_n} = \infty$, then $S_n \rightarrow \infty$ almost surely, but if $\sum_{n=1}^{\infty} \frac{1}{\lambda_n} < \infty$, then $S_n \rightarrow S$ almost surely, where S is some random variable which is almost surely finite.