Probability 1 CEU Budapest, fall semester 2017

Imre Péter Tóth

Homework sheet 6 – due on 05.12.2017 – and exercises for practice

- 6.1 Durrett [1], Exercise 5.2.13
- 6.2 (homework) Let X_n be a simple random walk on \mathbb{Z} starting from $X_0 = 0$. (As before, this means that $X_n = \xi_1 + \xi_2 + \cdots + \xi_n$, where the ξ_i are i.i.d. with $\mathbb{P}(X_i = 1) = p = 1 \mathbb{P}(X_i = -1)$, and $p \in [0,1]$. (p need not be $\frac{1}{2}$, so the walk may be asymmetric.) Use the martingale convergence theorem to show that
 - a.) the walk reaches the set $\{-20, 30\}$ with probability 1.
 - b.) If $p \ge \frac{1}{2}$, then the walk reaches the point 30 with probability 1.
 - c.) If $p \leq \frac{1}{2}$, then the walk reaches the point -20 with probability 1.
- 6.3 (Pólya's urn) In an urn there is initially (at time n = 0) a black and a white ball. At each time step n = 1, 2, ...
 - we draw a ball from the urn, uniformly at random,
 - we look at its colour,
 - we put it back, and we add another ball of the same colour.

(So we add exactly one ball in each step.) Let X_n be the number of white balls in the urn after n steps, and let $M_n = \frac{X_n}{n+2}$ be the proportion of white balls after n steps.

- a.) Show that X_n is uniform on $\{1, 2, ..., n+1\}$. (Hint: a possible solution is by induction.)
- b.) Show that M_n is almost surely convergent.
- c.) What is the distribution of $M_{\infty} := \lim_{n \to \infty} M_n$?
- 6.4 (homework) In the (French style) Roulette, if you bet on "red", you lose your bet with probability $\frac{19}{37}$, and you win the amount of your bet with the remaining probability $\frac{18}{37}$. (E.g. if you bet on "red" with HUF 1 and you win, then you get your HUF 1 back, plus you get another HUF 1 as your winning.

You arrive at the casino with some money in your pocket, and keep betting on "red". At each spin, your bet may be anything between 0 and the amount of money you have. Let X_n be the amount of your money after n spins. Show that – no matter what your strategy is – X_n is convergent with probability 1.

- 6.5 Alice and Bob keep tossing a possibly biased coin. Before each toss, they agree on a stake: Alice will give this sum to Bob if the coin turns "heads", and Bob will give the (same) sum to Alice if it turns "tails". The stake has to be a non-negative multiple of 1 penny, and they are not allowed to risk more money than what they have. If they agree on a stake which is 0, then the game ends. Show that sooner or later the game will end.
- 6.6 (homework) Harry is organizing a pyramid scheme in his family.

(See http://en.wikipedia.org/wiki/Pyramid_scheme) The participants are not too persistent: every participant keeps trying to recruit new participants until the first failure (i.e. until he is first rejected). The probability of such a failure is p at every recruit attempt, independently of the history of the scheme.

The first participant is Harry, he forms the 0-th generation alone. The first generation consists of those recruited (directly) by Harry. The second generation consists of those recruited (directly) by members of the first generation, and so on.

Let Z_k denote the size of the k-th generation (k = 0, 1, 2, ...), and let N denote the total number of participants in the scheme (meaning $N = \sum_{k=0}^{\infty} Z_k$).

0-th question: What is the distribution of Z_1 (which is the same as the distribution of the number of participants recruited by any fixed member of the scheme)? This distribution has a name.

Answer the questions below

- I. for $p = \frac{2}{3}$,
- II. for $p = \frac{1}{2}$,
- III. for $p = \frac{1}{3}$:
- a.) Let r be the probability that the scheme dies out (that is, one of the generations will already be empty). Is r = 1?
- b.) What is the expectation of Z_n ?
- c.) What is the expectation of N?
- d.) In case "not dying out" has positive probability, what is the growth rate of Z_n on this event?

References

[1] Durrett, R. Probability: Theory and Examples. 4th edition, Cambridge University Press (2010)