

**Probability 1**  
**CEU Budapest, fall semester 2013**

Imre Péter Tóth

**Homework sheet 2 – due on 07.10.2013 – and exercises for practice**

2.1 (**homework**) Exercise 3 of “Homework sheet 1”, delayed from last week (unless already done)

2.2 (**homework**) Exercise 4 of “Homework sheet 1”, delayed from last week (unless already done)

2.3 *Continuity of the measure*

(a) Prove the following:

**Theorem 1** (*Continuity of the measure*)

*i. If  $(\Omega, \mathcal{F}, \mu)$  is a measure space and  $A_1, A_2, \dots$  is an increasing sequence of measurable sets (i.e.  $A_i \in \mathcal{F}$  and  $A_i \subset A_{i+1}$  for all  $i$ ), then  $\mu(\cup_{i=1}^{\infty} A_i) = \lim_{i \rightarrow \infty} \mu(A_i)$  (and both sides of the equation make sense).*

*ii. If  $(\Omega, \mathcal{F}, \mu)$  is a measure space,  $A_1, A_2, \dots$  is a decreasing sequence of measurable sets (i.e.  $A_i \in \mathcal{F}$  and  $A_i \supset A_{i+1}$  for all  $i$ ) and  $\mu(A_1) < \infty$ , then  $\mu(\cap_{i=1}^{\infty} A_i) = \lim_{i \rightarrow \infty} \mu(A_i)$  (and both sides of the equation make sense).*

(b) Show that in the second statement the condition  $\mu(A_1) < \infty$  is needed, by constructing a counterexample for the statement when this condition does not hold.

2.4 *Usefulness of the linearity of the expectation.* A building has 10 floors, not including the ground floor. On the ground floor, 10 people get into the elevator, and every one of them chooses a destination at random, uniformly out of the 10 floors, independently of the others. Let  $X$  denote the number of floors *on which the elevator stops* – i.e. the number of floors that were chosen by at least one person. Calculate the expectation and the variance of  $X$ . (*hint: First notice that the distribution of  $X$  is hard to calculate. Find a way to calculate the expectation and the variance without that.*)

2.5 Calculate the characteristic function of

(a) The Bernoulli distribution  $B(p)$  (see Homework sheet 1)

(b) The “pessimistic geometric distribution with parameter  $p$ ” – that is, the distribution  $\mu$  on  $\{0, 1, 2, \dots\}$  with weights  $\mu(\{k\}) = (1 - p)p^k$  ( $k = 0, 1, 2, \dots$ ).

(c) The “optimistic geometric distribution with parameter  $p$ ” – that is, the distribution  $\nu$  on  $\{1, 2, 3, \dots\}$  with weights  $\nu(\{k\}) = (1 - p)p^{k-1}$  ( $k = 1, 2, \dots$ ).

(d) (**homework**) The Poisson distribution with parameter  $\lambda$  – that is, the distribution  $\eta$  on  $\{0, 1, 2, \dots\}$  with weights  $\eta(\{k\}) = e^{-\lambda} \frac{\lambda^k}{k!}$  ( $k = 0, 1, 2, \dots$ ).

(e) (**homework**) The exponential distribution with parameter  $\lambda$  – that is, the distribution on  $\mathbb{R}$  with density (w.r.t. Lebesgue measure)

$$f_{\lambda}(x) = \begin{cases} \lambda e^{-\lambda x}, & \text{if } x > 0 \\ 0, & \text{if not} \end{cases} .$$

2.6 (**homework**) Calculate the characteristic function of the normal distribution  $\mathcal{N}(m, \sigma^2)$ . (Remember the definition from the old times:  $\mathcal{N}(m, \sigma^2)$  is the distribution on  $\mathbb{R}$  with density (w.r.t. Lebesgue measure)

$$f_{m, \sigma^2}(x) = \frac{1}{\sqrt{2\pi\sigma}} e^{-\frac{(x-m)^2}{2\sigma^2}}.$$

You can save yourself some paperwork if you only do the calculation for  $\mathcal{N}(0, 1)$  and reduce the general case to this using the relation between different normal distributions. You can and should use the fact that

$$\int_{-\infty}^{\infty} f_{m, \sigma^2}(x) dx = 1$$

for every  $m$  and  $\sigma$ .

2.7 *Dominated convergence and continuous differentiability of the characteristic function.*

The Lebesgue dominated convergence theorem is the following

**Theorem 2 (dominated convergence)** *Let  $(\Omega, \mathcal{F}, \mu)$  be a measure space and  $f_1, f_2, \dots$  measurable real valued functions on  $\Omega$  which converge to the limit function pointwise,  $\mu$ -almost everywhere. (That is,  $\lim_{n \rightarrow \infty} f_n(x) = f(x)$  for every  $x \in \Omega$ , except possibly for a set of  $x$ -es with  $\mu$ -measure zero.) Assume furthermore that the  $f_n$  admit a common integrable dominating function: there exists a  $g : \Omega \rightarrow \mathbb{R}$  such that  $|f_n(x)| \leq g(x)$  for every  $x \in \Omega$  and  $n \in \mathbb{N}$ , and  $\int_{\Omega} g d\mu < \infty$ . Then (all the  $f_n$  and also  $f$  are integrable and)*

$$\lim_{n \rightarrow \infty} \int_{\Omega} f_n d\mu = \int_{\Omega} f d\mu.$$

Use this theorem to prove the following

**Theorem 3 (differentiability of the characteristic function)** *Let  $X$  be a real valued random variable,  $\psi(t) = \mathbb{E}(e^{itX})$  its characteristic function and  $n \in \mathbb{N}$ . If the  $n$ -th moment of  $X$  exists and is finite (i.e.  $\mathbb{E}(|X|^n) < \infty$ ), then  $\psi$  is  $n$  times continuously differentiable and*

$$\psi^{(k)}(0) = i^k \mathbb{E}(X^k), \quad k = 0, 1, 2, \dots, n.$$