

---

## Practice exercises 11.

1. Find the following Taylor polynomials of order  $n$  and center  $x_0$ .

a)  $f(x) = \sin x, x_0 = \frac{\pi}{3}, n = 3$       b)  $f(x) = 2^x, x_0 = 1, n = 3$

c)  $f(x) = \tan x, x_0 = 0, n = 3$       d)  $f(x) = \tan x, x_0 = \frac{\pi}{4}, n = 2$

e)  $f(x) = \log(1 - x), x_0 = 1, n = 4$       f)  $f(x) = \arccos x^2, x_0 = 0, n = 2$

2. Let  $P(x)$  be a polynomial of degree  $n$ . Prove that the Taylor polynomial of order  $n$  corresponding to  $P(x)$  at any center  $x_0 \in \mathbb{R}$  is  $P(x)$  itself.

3. Estimate the value of  $\sqrt{65}$  by the Taylor polynomial of order 2 of  $f(x) = \sqrt{x}$  at center 64. Give an upper bound for the error of the approximation.

4. Estimate the value of  $\log 1.2$  by the Taylor polynomial of order 3 of  $f(x) = \log(1 + x)$  at center 0. Give an upper bound for the error of the approximation.

5. Estimate the value of  $\sinh 1$  by an appropriate Taylor polynomial with error less than  $10^{-2}$ .

6. Calculate the Taylor series of the following function with center  $x_0$  and find the radius of convergence.

a)  $f(x) = \sin 2x, x_0 = \pi$

b)  $f(x) = 3^x, x_0 = 1$

c)  $f(x) = \frac{1}{x - 2}, x_0 = 0$

d)  $f(x) = \frac{1}{x - 2}, x_0 = 5$

e)  $f(x) = \frac{1}{x^2 + 3}, x_0 = 0$

f)  $f(x) = \frac{x^5}{x^2 + 3}, x_0 = 0$

g)  $f(x) = \frac{1}{(1 - x)^2}, x_0 = 2$

h)  $f(x) = \sinh 3x^3, x_0 = 0$

i)  $f(x) = \arccos x, x_0 = 0$

j)  $f(x) = \cosh x, x_0 = -1$

k)  $f(x) = x^2 \cdot \sqrt[3]{64 - 8x^2}, x_0 = 0$

l)  $f(x) = \frac{x}{\sqrt{x - 1}}, x_0 = 3$