

12.1 (homework) *Ising model: Weak sensitivity of the canonical probability of a “+” on a single neighbouring spin at high temperature.* Consider the (nearest neighbour ferromagnetic) Ising model without external field on \mathbb{Z}^d , with interaction constant $J = 1$:

$$H(\sigma) = -\frac{1}{2} \sum_{|i-j|=1} \sigma_i \sigma_j = - \sum_{\{i,j\}: i \sim j} \sigma_i \sigma_j$$

(That is: count every pair of neighbours once with interaction 1, or count every pair twice with interaction 1/2.) Let $\gamma_k^\beta(+|\eta)$ denote the probability of finding a spin +1 at position $k \in \mathbb{Z}^d$ with respect to the canonical measure in the box $\Lambda = \{k\}$ with boundary condition $\eta \in \Omega = \{-1, 1\}^{\mathbb{Z}^d}$. For $\eta \in \Omega$ and $j \in \mathbb{Z}^d$, let η_j^+ denote the configuration which is the same as η everywhere except possibly at j , where it is +1. Define η_j^- similarly.

Show that for any $k, j \in \mathbb{Z}^d$ and any $\eta \in \Omega$,

$$|\gamma_k^\beta(+|\eta_j^+) - \gamma_k^\beta(+|\eta_j^-)| \leq \beta \mathbf{1}_{k \sim j}.$$

12.2 (homework) *Ising model: independence of the thermodynamic limit from the boundary conditions.*

(a) Consider a finite $\Lambda \subset \mathbb{Z}^d$ and the phase space $\Omega_\Lambda = \{-1, 1\}^\Lambda$ with two different Hamiltonians $H_\Lambda, H'_\Lambda : \Omega \rightarrow \mathbb{R}$. Let $\|H_\Lambda - H'_\Lambda\|$ denote the distance of H_Λ and H'_Λ in the supremum norm:

$$\|H_\Lambda - H'_\Lambda\| := \max_{\sigma \in \Omega_\Lambda} |H_\Lambda(\sigma) - H'_\Lambda(\sigma)|.$$

Consider the canonical distributions with the partition functions $Z(\Lambda, \beta)$ and $Z'(\Lambda, \beta)$ and, as usual, let $p_\Lambda(\beta)$ and $p'_\Lambda(\beta)$ denote the thermodynamic pressures for the two Hamiltonians:

$$\begin{aligned} p_\Lambda(\beta) &= \frac{1}{\beta|\Lambda|} \log Z(\Lambda, \beta), \\ p'_\Lambda(\beta) &= \frac{1}{\beta|\Lambda|} \log Z'(\Lambda, \beta). \end{aligned}$$

Estimate $|p_\Lambda(\beta) - p'_\Lambda(\beta)|$ with the help of $\|H_\Lambda - H'_\Lambda\|$.

(b) We say that $\Lambda_n \nearrow \mathbb{R}^d$ if $\Lambda_n \subset \Lambda_{n+1}$ and $\bigcup_n \Lambda_n = \mathbb{R}^d$. We say that $\Lambda_n \nearrow \mathbb{R}^d$ in the sense of van Hove if $\Lambda_n \nearrow \mathbb{R}^d$ and $\frac{|\partial\Lambda_n|}{|\Lambda_n|} \rightarrow 0$, where $\partial\Lambda$ denotes the boundary of Λ :

$$\partial\Lambda := \{k \in \Lambda : \text{dist}(k, \Lambda^c) = 1\}.$$

Consider the Ising model (with external field) for a sequence Λ_n such that $\Lambda_n \nearrow \mathbb{R}^d$ in the sense of van Hove, and with an arbitrary sequence η_n of boundary conditions. Show that the limiting thermodynamic pressure

$$p(\beta, h) := \lim_{n \rightarrow \infty} p_{\Lambda_n}(\beta, h)$$

is independent of the boundary conditions.

(Remark: We have not fully checked, but it is true that the limit exist for empty boundary conditions, and is independent of the sequence Λ_n .)

12.3 *Symmetry breaking in the 2d Ising model: Peierls argument.* For \mathbb{Z}^2 an *elementary contour* γ is a sequence of *edges* of \mathbb{Z}^2 which forms a non-self-intersecting circle of the dual graph. Such a contour has an “inside” and an “outside”, both consisting of points of \mathbb{Z}^2 . Consider the (nearest neighbour) ferromagnetic Ising model on \mathbb{Z}^2 without external field. For a configuration $\sigma \in \Omega = \{\pm 1\}^{\mathbb{Z}^d}$, an elementary contour γ is said to be *present* if all the spins adjacent to the contour on the inside are $+1$ and all the spins adjacent to the contour on the outside are -1 , or vice versa.

- (a) For $\Lambda \subset \mathbb{R}^2$ and an elementary contour γ which is “inside Λ ”, show that there is a 1 – 1 correspondance between configurations σ for which γ is present, and configurations σ' for which all the spins adjacent to γ are the same.
- (b) Estimate the probability (w.r.t. the canonical measure) of a given contour γ of length n being present.
- (c) Give a rough estimate for the number of elementary contours of length n that surround the origin. The estimate can be rough, but make sure that it's at most exponential in n .
- (d) Based on the previous two points, give the easiest possible estimate for the probability that a contour surrounding the origin is present.
- (e) Consider the ferromagnetic Ising model (without external field) in the box $\Lambda \subset \mathbb{R}^d$ with the boundary condition “ $\eta = +1$ everywhere”. Show that if β is big enough, then there is an $\alpha = \alpha(\beta) < \frac{1}{2}$ such that for *any* Λ

$$\mu_{can}^\beta(\{\sigma_0 = -1\}|\eta) \leq \alpha.$$

- (f) Conclude that for β big enough, the Gibbs measure is not unique.
- (g) Conclude that for β big enough, $p(\beta, h)$ is not differentiable in h at $h = 0$.