Probability 1

CEU Budapest, fall semester 2018

Imre Péter Tóth

Homework sheet 5 – due on 26.11.2018

5.1 Prove that

$$\lim_{n \to \infty} \int_0^1 \int_0^1 \dots \int_0^1 \frac{x_1^2 + x_2^2 + \dots + x_n^2}{x_1 + x_2 + \dots + x_n} dx_1 dx_2 \dots dx_n = \frac{2}{3}.$$

5.2 Let $f:[0;1] \to \mathbb{R}$ be a continuous function. Prove that

(a)
$$\lim_{n \to \infty} \int_0^1 \int_0^1 \dots \int_0^1 f\left(\frac{x_1 + x_2 + \dots + x_n}{n}\right) dx_1 dx_2 \dots dx_n = f\left(\frac{1}{2}\right).$$

(b)
$$\lim_{n \to \infty} \int_0^1 \int_0^1 \dots \int_0^1 f\left((x_1 x_2 \dots x_n)^{1/n}\right) dx_1 dx_2 \dots dx_n = f\left(\frac{1}{e}\right).$$

(Hint: interprete these integrals as expectations.)

- 5.3 (homework) Let $X_n \sim Bin(n, \frac{2}{3})$. Calculate $\lim_{n\to\infty} \mathbb{E}\left(\sin\left(\left(\frac{X_n}{n}\right)^4\right)\right)$.
- 5.4 Poisson approximation of the binomial distribution. Fix $0 < \lambda \in \mathbb{R}$. Show that if X_n has binomial distribution with parameters (n, p) such that $np \to \lambda$ as $n \to \infty$, then X_n converges to $Poi(\lambda)$ weakly. This can be done in a completely elementary way, using your favourite definition of weak convergence, or by using one of the stronger tools of weak convergence.
- 5.5 Continuous limit of the geometric distribution. Let X_n be geometrically distributed with parameter $p_n = \frac{1}{n}$ and let $Y_n = \frac{1}{n}X_n$. (So $\mathbb{E}Y_n = 1$.) Find the weak limit of Y_n . (Hint: you can use the method of characteristic functions, but you can also calculate the limiting distribution function directly.)
- 5.6 Continuous limit of the geometric distribution, general version. Show that if $0 \le p_n \to 0$, $0 \le a_n \to 0$, $\frac{p_n}{a_n} \to \lambda \in (0, \infty)$ and $X_n \sim Geom(p_n)$, then $a_n X_n \Rightarrow Exp(\lambda)$.
- 5.7 Let X be uniformly distributed on [-1;1], and set $Y_n = nX$.
 - a.) Calculate the characteristic function ψ_n of Y_n .
 - b.) Calculate the pointwise limit $\lim_{n\to\infty} \psi_n(t)$, if it exists.
 - c.) Does (the distribution of) Y_n have a weak limit?
 - d.) How come?
- 5.8 (homework) Show that if Ψ is the characteristic function of some random variable X, then the complex conjugate $\bar{\Psi}$ is also the characteristic function of some random variable Y. (Hint: try to find out what Y is.)
- 5.9 Durrett [1], Exercise 3.3.1 (Hint: try to find the appropriate random variables. Use Exercise 8.)
- 5.10 Durrett [1], Exercise 3.3.3
- 5.11 Durrett [1], Exercise 3.3.9
- 5.12 Durrett [1], Exercise 3.3.10. Show also that independence is needed.

- 5.13 Durrett [1], Exercise 3.3.11
- 5.14 Durrett [1], Exercise 3.3.12
- 5.15 Durrett [1], Exercise 3.3.13
- 5.16 Let X_1, X_2, \ldots be i.i.d. random variables with density (w.r.t. Lebesgue measure) $f(x) = \frac{1}{\pi} \frac{1}{1+x^2}$. (So they have the Cauchy distribution.) Find the weak limit (as $n \to \infty$) of the average

$$\frac{X_1 + \dots + X_n}{n}.$$

Warning: this is not hard, but also not as trivial as it may seem. Hint: a possible solution is using characteristic functions. Calculating the characteristic function of the Cauchy distribution is a little tricky, but you can look it up.

- 5.17 Durrett [1], Exercise 3.4.4
- 5.18 (homework) Durrett [1], Exercise 3.4.5 (Hint: Use Exercise 4.1 and Durrett [1], Exercise 3.2.14).

References

[1] Durrett, R. Probability: Theory and Examples. 4th edition, Cambridge University Press (2010)