

## Limit / large dev. thms. first midterm practice

1. (*easy*) Let  $f(x) = |x - 1| + |x + 1|$ . Find the Legendre transform of  $f$ .
2. (*easy*) Let  $Z(\lambda)$  denote the moment generating function of the r.v.  $X$ . Denote by  $X^{(\mu)}$  the exponentially tilted random variable (tilted with parameter  $\mu \in \mathbb{R}$ ). Let  $Z_\mu(\lambda) = \mathbb{E}(\exp(\lambda X^{(\mu)}))$  denote the moment generating function of  $X^{(\mu)}$ . Write down an identity between  $Z(\lambda + \mu)$ ,  $Z_\mu(\lambda)$  and  $Z(\mu)$ .
3. (*medium*) If  $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$ , denote by  $\{f, g\}^{co}$  the joint lower convex envelope of  $f$  and  $g$ , i.e., the supremum of those affine linear functions that lie below both  $f$  and  $g$ . Show that if  $f$  and  $g$  are both convex and continuous then the Legendre transform of  $\max\{f, g\}$  is  $\{\hat{f}, \hat{g}\}^{co}$  and that the Legendre transform of  $\{f, g\}^{co}$  is  $\max\{\hat{f}, \hat{g}\}$ .
4. (*easy*) We use a randomized algorithm to solve a yes/no decision problem. The algorithm gives the correct answer with probability  $p > \frac{1}{2}$ . We run the algorithm  $n$  times (where  $n$  is an odd number) and make our decision based on the majority of the results. Use the exponential Chebyshev's inequality (à la Cramér) to give a very good upper bound the probability that we make a wrong decision. Simplify the formula that you obtain as much as possible.
5. (*medium*) Let  $X$  and  $Y$  denote independent random variables. Denote by  $I_X$  and  $I_Y$  the large deviation rate function of  $X$  and  $Y$ , respectively. Show that the large deviation function  $I_{X+Y}$  of  $X + Y$  is the „infimum convolution” of  $I_X$  and  $I_Y$ .  
*Hint:* A non-rigorous proof using the heuristic meaning of Cramér's theorem (and our  $\approx$  notation) is OK. A rigorous proof is even better. You should figure out by yourselves the notion of „infimum convolution” (or Google it)
6. (*medium*) Let  $X$  denote the random variable with p.d.f.

$$f(x) = 4xe^{-2x} \mathbf{1}[x \geq 0]$$

Let  $Y$  denote the sum of 1000 i.i.d. copies of  $X$ .

- (a) Find the Legendre transform of the logarithmic mom.gen. function of  $X$ .
  - (b) Denote by  $g$  the p.d.f. of  $Y$ . Approximate  $g(1000)$ .
  - (c) Estimate the number of zeroes in the decimal expansion of  $\mathbb{P}(Y \leq 500)$ .
7. (*medium*) I roll a fair die 1000 times. Denote by  $X$  the sum of the numbers rolled.
    - (a) Estimate the probability that  $X$  is greater than or equal to 3550.
    - (b) Estimate the probability that  $X$  is exactly equal to 3550.
    - (c) Give a good lower bound on the number of zeroes in the decimal expansion of the probability that  $X$  is greater than or equal to 4500.
  8. Let  $X_1, X_2, \dots$  denote i.i.d. r.v.'s with  $\text{POI}(\lambda)$  distribution. Let  $S_n = X_1 + \dots + X_n$ .
    - (a) (*medium/hard*) Use Stirling's formula to prove the *local CLT* for  $S_n$ :

$$\lim_{n \rightarrow \infty} \sqrt{n\lambda} \mathbb{P}\left(S_n = \lfloor n\lambda + \sqrt{n\lambda}x \rfloor\right) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}.$$

- (b) (*easy*) Deduce the *global CLT* from the local CLT: show that

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(\frac{S_n - \mathbb{E}(S_n)}{\sqrt{\text{Var}(S_n)}} \leq x\right) = \int_{-\infty}^x \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy.$$

9. (*medium*) Before the national election, we want to estimate the fraction  $p$  of republican voters. We ask  $n$  random people and calculate the fraction  $p_n$  of republicans in our sample. How to choose  $n$  if we want to estimate the value of  $p$  with a margin of error 0.01 with 95% confidence? Use the CLT.

10. (*easy*) *Weibull distribution.* Let  $U_1, U_2, \dots$  denote i.i.d. random variables with  $\text{UNI}[0, 1]$  distribution. Let  $\beta > 0$ . Let

$$M_n = \min\{U_1^\beta, \dots, U_n^\beta\}.$$

Show that  $n^\alpha M_n$  converges in distribution as  $n \rightarrow \infty$  to a non-trivial probability distribution if we choose  $\alpha > 0$  correctly. Determine the cumulative distribution function (c.d.f.)  $F(x)$  of the limiting distribution.

11. (*medium/hard*) Let  $\tau_1, \tau_2, \dots$  be i.i.d. waiting times between successive events and define the *renewal process*

$$\nu_t := \max \left\{ n : \sum_{i=1}^n \tau_i < t \right\}.$$

In plain words,  $\nu_t$  is the number of events that occurred in the time interval  $[0, t]$ . Denote  $m := \mathbb{E}(\tau_j) < \infty$ ,  $\sigma^2 := \text{Var}(\tau_j) < \infty$ . Use the classic CLT for the sum of i.i.d. r.v.'s to derive a CLT for  $\nu_t$ : find  $a > 0, b > 0$  such that

$$\lim_{t \rightarrow \infty} \mathbb{P} \left( \frac{\nu_t - at}{b\sqrt{t}} < x \right) = \Phi(x),$$

where  $\Phi$  is the standard normal c.d.f. Express  $a$  and  $b$  in terms of  $m$  and  $\sigma$ .

12. (*medium/hard*) Let  $X_1, X_2, \dots$  denote i.i.d. r.v.'s and assume that  $\mathbb{E}(X_i) = 0$  and  $\mathbb{P}(|X_i| \leq K) = 1$  for some  $K \in \mathbb{R}$ . Let us define

$$Y_n = \prod_{k=1}^n \left( 1 + \frac{X_k}{\sqrt{n}} \right).$$

Use the CLT to show that  $Y_n$  converges weakly as  $n \rightarrow \infty$ . The limiting distribution is famous (e.g., in financial mathematics): name it and identify its parameter(s).

13. (*hard*) Let  $X_0^n, X_1^n, X_2^n, \dots, X_{2n}^n$  denote the conditional distribution of one dimensional simple symmetric random walk under the condition that it returns to the origin in  $2n$  steps, i.e., that  $X_{2n}^n = 0$ . Denote by  $M_n = \max\{X_0^n, X_1^n, X_2^n, \dots, X_{2n}^n\}$ . Show that  $M_n/\sqrt{n}$  converges in distribution as  $n \rightarrow \infty$  and find the c.d.f. of the limiting distribution.

14. (*medium*) For  $n, m \in \mathbb{N}$  let  $X_n, Y_m$  be independent r.v.'s with distributions  $X_n \sim \text{POI}(n), Y_m \sim \text{POI}(m)$ . Prove that

$$\frac{X_n - n - (Y_m - m)}{\sqrt{X_n + Y_m}}$$

converges in distribution as  $n, m \rightarrow \infty$ . Identify the limiting distribution.

*Hint:* This is easy if you use Slutsky in a clever way, similarly to page 54 of the scanned lecture notes.

15. (*easy/medium*) Let  $X_1, X_2, \dots$  denote i.i.d. random variables with p.d.f.  $f(x) = \frac{1}{(x-1)^2} \mathbf{1}[x \leq 0]$ . Let  $M_n = \max\{X_1, \dots, X_n\}$ . Find a sequence  $(a_n)$  such that  $M_n/a_n$  weakly converges to a non-degenerate random variable  $Z$ . Find the c.d.f. of  $Z$ . What does it mean that  $Z$  is max-stable?

16. (*medium*) At time zero a stock broker has 1000 dollars. At each time-step, three things can happen: either he gains one dollar (this happens with probability  $1/4$ ), loses one dollar (this also happens with probability  $1/4$ ) or he neither gains nor loses (this happens with probability  $1/2$ ).

How would you approximate the distribution of the time when the stock broker goes bankrupt (i.e., loses all his money)? How to make this rigorous?

17. (*easy*) Let  $Z_n$  denote an integer-valued random variable for which

$$\mathbb{P}(Z_n = k) = (k+1) \frac{1}{n^2} \left(1 - \frac{1}{n}\right)^k, \quad k = 0, 1, 2, \dots$$

Show that  $Z_n/n$  converges in distribution as  $n \rightarrow \infty$  and identify the limiting distribution.

18. (*medium*) Let  $X_1$  and  $X_2$  be i.i.d. random variables with Lévy distribution. What is the distribution of  $X_1 + 3X_2$ ?