

Limit/large dev. thms. HW assignment 2., Solutions

1. Let $\lambda \mapsto Z(\lambda)$ denote the moment generating function of the random variable X . If $Z(\mu) < +\infty$, let $X^{(\mu)}$ denote the *exponentially tilted* random variable. The cumulative distribution function of $X^{(\mu)}$ is

$$F_\mu(x) = \mathbb{P}(X^{(\mu)} \leq x) = \frac{\mathbb{E}(e^{\mu X} \mathbb{1}[X \leq x])}{Z(\mu)}.$$

We have learnt some facts about exponential tilting in class, see page 17-18 of scanned lecture notes.

- Show that $X^{(\lambda)^{(\mu)}} \sim X^{(\lambda+\mu)}$. In words: tilting the tilted random variable amounts to tilting the original random variable with the sum of the two tiltings.
- Express the logarithmic moment generating function \widehat{I}_μ of $X^{(\mu)}$ using the logarithmic moment generating function \widehat{I} of X .
- Express the Legendre transform I_μ of \widehat{I}_μ using the Legendre transform I of \widehat{I} .

Solution:

- We have learnt in class (see page 18 of scanned) that $\mathbb{E}(g(X^{(\mu)})) = \mathbb{E}(e^{\mu X} g(X)) / \mathbb{E}(e^{\mu X})$. We apply this formula in the numerator as well as the denominator in equation (*) below:

$$\begin{aligned} \mathbb{P}\left(X^{(\mu)^{(\lambda)}} \leq x\right) &= \frac{\mathbb{E}\left(e^{\lambda X^{(\mu)}} \mathbb{1}[X^{(\mu)} \leq x]\right)}{\mathbb{E}(e^{\lambda X^{(\mu)}})} \stackrel{(*)}{=} \frac{\mathbb{E}\left(e^{\mu X} e^{\lambda X} \mathbb{1}[X \leq x]\right) / \mathbb{E}(e^{\mu X})}{\mathbb{E}(e^{\mu X} e^{\lambda X}) / \mathbb{E}(e^{\mu X})} = \\ &= \frac{\mathbb{E}\left(e^{(\lambda+\mu)X} \mathbb{1}[X \leq x]\right)}{\mathbb{E}(e^{(\lambda+\mu)X})} = \mathbb{P}[X^{(\lambda+\mu)} \leq x] \end{aligned}$$

This shows that $X^{(\mu)^{(\lambda)}} \sim X^{(\mu+\lambda)}$.

- $\widehat{I}_\mu(\lambda) = \ln\left(\mathbb{E}(e^{\lambda X^{(\mu)}})\right) = \ln\left(\mathbb{E}(e^{\mu X} e^{\lambda X}) / \mathbb{E}(e^{\mu X})\right) = \widehat{I}(\mu + \lambda) - \widehat{I}(\mu)$
-

$$\begin{aligned} I_\mu(x) &= \sup_\lambda \{\lambda x - \widehat{I}_\mu(\lambda)\} = \sup_\lambda \{\lambda x - \widehat{I}(\mu + \lambda) + \widehat{I}(\mu)\} = \\ &= \widehat{I}(\mu) - \mu x + \sup_\lambda \{(\lambda + \mu)x - \widehat{I}(\mu + \lambda)\} = \widehat{I}(\mu) - \mu x + I(x) = I(x) - \mu x + \widehat{I}(\mu). \end{aligned}$$

Remark: Alternative proof of $I_\mu(x) = I(x) - \mu x + \widehat{I}(\mu)$ using our non-rigorous Cramér formalism: we know that $X_1^{(\mu)} + \dots + X_n^{(\mu)}$ has the same distribution as $S_n^{(\mu)}$, thus

$$e^{-nI_\mu(x)} \approx \mathbb{P}\left(S_n^{(\mu)} \approx nx\right) = \frac{e^{\mu nx}}{\mathbb{E}(e^{\lambda S_n})} \mathbb{P}(S_n \approx nx) \approx \frac{e^{\mu nx}}{Z(\lambda)^n} e^{-nI(x)} = e^{-n(I(x) - \mu x + \widehat{I}(\mu))}.$$

2. In this exercise $X^{(\mu)}$ denotes the random variable that we obtain by exponentially tilting the distribution of the random variable X .

- (a) Show that if $X \sim \text{BIN}(n, p)$ then $X^{(\mu)} \sim \text{BIN}(n, p')$ for some $p' = p'(p, \mu)$ and that any $p' \in (0, 1)$ can be obtained by choosing $\mu \in \mathbb{R}$ appropriately.
- (b) Show that if $X \sim \text{POI}(\lambda)$ then $X^{(\mu)} \sim \text{POI}(\lambda')$ for some $\lambda' = \lambda'(\lambda, \mu)$ and that any $\lambda' \in (0, +\infty)$ can be obtained by choosing $\mu \in \mathbb{R}$ appropriately.
- (c) If $X \sim \text{EXP}(\lambda)$, find the values of μ for which $Z(\mu) < +\infty$ (i.e., find the domain of the moment generating function $Z(\cdot)$). Show that $X^{(\mu)} \sim \text{EXP}(\lambda')$ for some $\lambda' = \lambda'(\lambda, \mu)$ and that any $\lambda' \in (0, +\infty)$ can be obtained by choosing μ from the domain of $Z(\cdot)$ appropriately.
- (d) Show that if $X \sim \mathcal{N}(m, \sigma^2)$ then $X^{(\mu)} \sim \mathcal{N}(m', \sigma^2)$ for some $m' = m'(m, \mu, \sigma)$ and that any $m' \in (-\infty, +\infty)$ can be obtained by choosing $\mu \in \mathbb{R}$ appropriately.

Solution:

- (a) If $X \sim \text{BIN}(n, p)$ then $Z(\mu) = (pe^\mu + (1-p))^n$ (since X is the sum of n i.i.d. $\text{BER}(p)$ random variables), thus

$$\mathbb{P}(X^{(\mu)} = k) = (pe^\mu + (1-p))^{-n} e^{\mu k} \binom{n}{k} p^k (1-p)^{n-k} = \binom{n}{k} \left(\frac{pe^\mu}{pe^\mu + (1-p)} \right)^k \left(\frac{(1-p)}{pe^\mu + (1-p)} \right)^{n-k}, \quad \text{thus } X^{(\mu)} \sim \text{BIN} \left(n, \frac{pe^\mu}{pe^\mu + (1-p)} \right).$$

Thus $p'(p, \mu) = \frac{pe^\mu}{pe^\mu + (1-p)}$. Clearly, $\lim_{\mu \rightarrow -\infty} p'(p, \mu) = 0$ and $\lim_{\mu \rightarrow +\infty} p'(p, \mu) = 1$, so indeed any $p' \in (0, 1)$ can be obtained by choosing $\mu \in \mathbb{R}$ appropriately.

- (b) If $X \sim \text{POI}(\lambda)$ then $Z(\mu) = e^{(e^\mu - 1)\lambda}$, thus

$$\mathbb{P}(X^{(\mu)} = k) = e^{(1-e^\mu)\lambda} e^{\mu k} e^{-\lambda} \frac{\lambda^k}{k!} = e^{-e^\mu \lambda} \frac{(e^\mu \lambda)^k}{k!}, \quad \text{thus } X^{(\mu)} \sim \text{POI}(e^\mu \lambda).$$

Thus $\lambda'(\lambda, \mu) = e^\mu \lambda$. Clearly, $\lim_{\mu \rightarrow -\infty} \lambda'(\lambda, \mu) = 0$ and $\lim_{\mu \rightarrow +\infty} \lambda'(\lambda, \mu) = +\infty$, so indeed any $\lambda' \in (0, +\infty)$ can be obtained by choosing $\mu \in \mathbb{R}$ appropriately.

- (c) If $X \sim \text{EXP}(\lambda)$ then the density function of X is $f(x) = \lambda e^{-\lambda x} \mathbf{1}[x \geq 0]$, thus

$$Z(\mu) = \mathbb{E}(e^{\mu X}) = \int_0^\infty e^{\mu x} f(x) dx = \int_0^\infty \lambda e^{(\mu - \lambda)x} dx = \frac{\lambda}{\lambda - \mu}, \quad \text{if } \mu < \lambda.$$

If $\mu \geq \lambda$ then $Z(\mu) = +\infty$.

Thus $f^{(\mu)}(x) = Z(\mu)^{-1} e^{\mu x} f(x) = (\lambda - \mu) e^{-(\lambda - \mu)x} \mathbf{1}[x \geq 0]$, thus $X^{(\mu)} \sim \text{EXP}(\lambda - \mu)$, if $\mu < \lambda$. Thus $\lambda'(\lambda, \mu) = \lambda - \mu$. Clearly, any $\lambda' \in (0, +\infty)$ can be obtained by choosing $\mu = \lambda - \lambda'$.

- (d) If $X \sim \mathcal{N}(m, \sigma^2)$ then $f(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-m)^2}{2\sigma^2}\right)$ and $Z(\mu) = \exp(\mu m + \frac{1}{2}\sigma^2\mu^2)$, thus

$$f^{(\mu)}(x) = Z(\mu)^{-1} e^{\mu x} f(x) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-m)^2}{2\sigma^2} + \mu x - \mu m - \frac{1}{2}\sigma^2\mu^2\right) = \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-m-\sigma^2\mu)^2}{2\sigma^2}\right), \quad \text{thus } X^{(\mu)} \sim \mathcal{N}(m + \sigma^2\mu, \sigma^2).$$

Thus, $m'(m, \mu, \sigma) = m + \sigma^2\mu$ and indeed any $m' \in (-\infty, +\infty)$ can be obtained by choosing $\mu \in \mathbb{R}$ appropriately.

3. Let X_1, X_2, \dots denote i.i.d. non-negative integer-valued random variables with distribution $\mathbb{P}(X_i = k) = p_k$, where $k = 0, 1, 2, \dots$. Let $\lambda \in \mathbb{R}$ such that $Z(\lambda) = \mathbb{E}[e^{\lambda X_i}] < +\infty$. Let $S_n = X_1 + \dots + X_n$. Let $X_1^{(\lambda)}, X_2^{(\lambda)}, \dots$ denote i.i.d. non-negative integer-valued random variables with distribution

$$\mathbb{P}(X_i^{(\lambda)} = k) = \frac{1}{Z(\lambda)} e^{\lambda k} p_k, \quad \text{where } k = 0, 1, 2, \dots$$

- (a) Show that we have $\mathbb{P}(X_1^{(\lambda)} + \dots + X_n^{(\lambda)} = k) = \frac{e^{\lambda k} \mathbb{P}(X_1 + \dots + X_n = k)}{Z(\lambda)^n}$.

Instruction: This could be easily derived from the Lemma on page 20 of the scanned lecture notes, but since we only proved that lemma in the absolutely continuous case, I ask you to write down a complete proof of this sub-exercise only using the basic facts about exponential tilting (page 17-18 of scanned lecture notes).

- (b) Show that $\mathbb{P}(X_1^{(\lambda)} = k \mid X_1^{(\lambda)} + \dots + X_n^{(\lambda)} = m) = \mathbb{P}(X_1 = k \mid X_1 + \dots + X_n = m)$.
(c) If $X_i \sim \text{POI}(\mu)$, what is the conditional distribution of X_1 given that $X_1 + \dots + X_n = \lfloor nx \rfloor$?
(d) If $X_i \sim \text{POI}(\mu)$, show that for any $x > 0$ we have

$$\lim_{n \rightarrow \infty} \mathbb{P}(X_1 = k \mid X_1 + \dots + X_n = \lfloor nx \rfloor) = e^{-x} \frac{x^k}{k!}.$$

Remark: This last result is a rigorous version of the result proved heuristically on page 25 of the scanned lecture notes (also note that an exponentially tilted Poisson random variable is still a Poisson random variable with a different parameter)

Solution:

- (a) Denote by Λ the set of n -tuples (k_1, \dots, k_n) of non-negative integers that satisfy $k_1 + \dots + k_n = k$.

$$\begin{aligned} \mathbb{P}(X_1^{(\lambda)} + \dots + X_n^{(\lambda)} = k) &= \sum_{(k_1, \dots, k_n) \in \Lambda} \mathbb{P}(X_1^{(\lambda)} = k_1, \dots, X_n^{(\lambda)} = k_n) = \\ &= \sum_{(k_1, \dots, k_n) \in \Lambda} \mathbb{P}(X_1^{(\lambda)} = k_1) \dots \mathbb{P}(X_n^{(\lambda)} = k_n) = \sum_{(k_1, \dots, k_n) \in \Lambda} \frac{e^{\lambda k_1} p_{k_1}}{Z(\lambda)} \dots \frac{e^{\lambda k_n} p_{k_n}}{Z(\lambda)} = \\ &= \frac{e^{\lambda k}}{Z(\lambda)^n} \sum_{(k_1, \dots, k_n) \in \Lambda} p_{k_1} \dots p_{k_n} = \frac{e^{\lambda k}}{Z(\lambda)^n} \sum_{(k_1, \dots, k_n) \in \Lambda} \mathbb{P}(X_1 = k_1, \dots, X_n = k_n) = \\ &= \frac{e^{\lambda k} \mathbb{P}(X_1 + \dots + X_n = k)}{Z(\lambda)^n} \end{aligned}$$

- (b)

$$\begin{aligned} \mathbb{P}(X_1^{(\lambda)} = k \mid X_1^{(\lambda)} + \dots + X_n^{(\lambda)} = m) &= \frac{\mathbb{P}(X_1^{(\lambda)} = k, X_2^{(\lambda)} + \dots + X_n^{(\lambda)} = m - k)}{\mathbb{P}(X_1^{(\lambda)} + \dots + X_n^{(\lambda)} = m)} = \\ &= \frac{\mathbb{P}(X_1^{(\lambda)} = k) \mathbb{P}(X_2^{(\lambda)} + \dots + X_n^{(\lambda)} = m - k)}{\mathbb{P}(X_1^{(\lambda)} + \dots + X_n^{(\lambda)} = m)} = \frac{\frac{e^{\lambda k} p_k}{Z(\lambda)} \frac{e^{\lambda(m-k)} \mathbb{P}(X_2 + \dots + X_n = m - k)}{(Z(\lambda))^{n-1}}}{\frac{e^{\lambda m} \mathbb{P}(X_1 + \dots + X_n = m)}{(Z(\lambda))^n}} = \\ &= \frac{p_k \cdot \mathbb{P}(X_2 + \dots + X_n = m - k)}{\mathbb{P}(X_1 + \dots + X_n = m)} = \mathbb{P}(X_1 = k \mid X_1 + \dots + X_n = m) \end{aligned}$$

- (c) Let $Y = X_2 + \dots + X_n$. Then $Y \sim \text{POI}((n-1)\mu)$. For any $0 \leq k \leq \lfloor nx \rfloor$ we have

$$\begin{aligned} \mathbb{P}(X_1 = k \mid X_1 + Y = \lfloor nx \rfloor) &= \frac{\mathbb{P}(X_1 = k, Y = \lfloor nx \rfloor - k)}{\mathbb{P}(X_1 + Y = \lfloor nx \rfloor)} = \frac{e^{-\mu} \frac{\mu^k}{k!} \cdot e^{-(n-1)\mu} \frac{((n-1)\mu)^{\lfloor nx \rfloor - k}}{(\lfloor nx \rfloor - k)!}}{e^{-n\mu} \frac{(n\mu)^{\lfloor nx \rfloor}}{\lfloor nx \rfloor!}} = \\ &= \frac{\frac{\mu^k}{k!} \cdot \frac{((n-1)\mu)^{\lfloor nx \rfloor - k}}{(\lfloor nx \rfloor - k)!}}{\frac{(n\mu)^{\lfloor nx \rfloor}}{\lfloor nx \rfloor!}} = \frac{\lfloor nx \rfloor!}{k! (\lfloor nx \rfloor - k)!} \frac{\mu^k \cdot ((n-1)\mu)^{\lfloor nx \rfloor - k}}{(n\mu)^{\lfloor nx \rfloor}} = \binom{\lfloor nx \rfloor}{k} \left(\frac{1}{n}\right)^k \left(1 - \frac{1}{n}\right)^{\lfloor nx \rfloor - k}, \end{aligned}$$

thus the conditional distribution of X_1 given that $X_1 + Y = \lfloor nx \rfloor$ is $\text{BIN}(\lfloor nx \rfloor, \frac{1}{n})$.

(d)

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{[nx]!}{k!([nx] - k)!} \left(\frac{1}{n}\right)^k \left(1 - \frac{1}{n}\right)^{[nx] - k} &= \\ \frac{1}{k!} \lim_{n \rightarrow \infty} \frac{[nx]}{n} \frac{[nx] - 1}{n} \cdots \frac{[nx] - (k - 1)}{n} \left(1 - \frac{1}{n}\right)^{[nx] - k} &= \frac{1}{k!} x^k e^{-x}. \end{aligned}$$